THE REVIEW OF APPLIED **ENTOMOLOGY**

SERIES A: AGRICULTURAL

ISSUED BY THE COMMONWEALTH INSTITUTE OF ENTOMOLOGY



LONDON: COMMONWEALTH INSTITUTE OF ENTOMOLOGY 56. QUEEN'S GATE, S.W.7

Price 14s. net

Commonwealth Agricultural Bureaux

EXECUTIVE COUNCIL

W. A. C. MATHIESON, C.M.G., M.B.E., Chairman, Dependent Territories

H. E. SENEVIRATNE, Vice-Chairman, Ceylon

W. G. ALEXANDER, C.B.E., United Kingdom

J. B. MARSHALL, Ph.D., Canada

P. F. BUTLER, M.Agr.Sc., Australia

V. ARMSTRONG, Ph.D., New Zealand

A. M. D'ROZARIO, Ph.D., India

C. K. REHEEM, Pakistan

J. A. AFARI. Ghana

Toh Chor Keat, Malaysia

G. A. FALASE, Federation of Nigeria

HIS EXCELLENCY THE HIGH COMMISSIONER FOR SIERRA LEONE IN THE U.K.

G. M. NHIGULA, Tanganyika

C. S. McMorris, Jamaica

E. Murray, Trinidad and Tobago

M. KAGGWA, Uganda

J. E. C. COVENTRY, Federation of Rhodesia and Nyasaland

Sir THOMAS SCRIVENOR, Secretary, Farnham House, Farnham Royal, Nr. Slough, Bucks.

COMMONWEALTH INSTITUTE OF ENTOMOLOGY

Director and Editor E. O. Pearson, M.A.

Assistant Director R. G. Fennah, M.A.

Assistant Editor R. F. Avery, M.A.

Head Office—c/o British Museum (Natural History), Cromwell Road, London, S.W.7

Publication Office and Library-56, Queen's Gate, London, S.W.7

[@] Commonwealth Agricultural Bureaux, 1963

INDEX OF AUTHORS

A reference in heavy type indicates that a paper by the author has been noticed.

Abbas, H. M., 196. Abdel-Malek, A. A., 686. Abdul Kareem, A., 424. Abrahão, J., 557. Abul-Hab, J. K., 278. Abu Yaman, I. K., 39. Acatay, A., 355. Achremowicz, J., 40. Aćimović, M., 80. Adams, J. A., 23. Adams, J. M., 88. Adkisson, P. L., 269, 417, 465, 474. Adlung, K. G., 44. Agafonova, Z. Ya., 542. Agati, J. A., 96. Agenjo, R., 659. Agosin, M., 248. Aguilar, J. d', 7. Ahmed, A., 88. Aitkenhead, P., 425. Aizawa, K., 496. Akşit, F., 82. Alam, M. Z., 88, 174, 176, 693. Alam, S. Mashhood, 91. Albrecht, F. O., 285. Alexander, R. D., 567. Ali, Hadi, 317. Alkan, B., 82, 177, 297, 609. Allan, R. E., 65. Allen, N., 465, 635. Allen, P. T., 562. Allen, T. C., 416. Aller, H. E., 271, 272. Almeida, E., 145, 646. Almeida, P. R. de, 557. Amaral, S. Franco do, 555, 644. Ambrosi, M., 183. Amitai, S., 614. Amman, G. D., 513. Ananth, K. C., 448. Anasiewicz, A., 545. Anbaroglu, M. A., 81. Andersen, A. L., 16.

Anderson, M. A., 395. Anderson, N. H., 158, 429. Anderson, N. L., 223. Anderson, W. H., 496. Andrade, A. C., 557. Andreescu, E., 56. Androić, M., 535, 536. Anglade, P., 371, 433, 666. Angus, T. A., 88, 196. Ankersmit, G. W., 205. Annamma Kurup, 117. Annecke, D. P., 490. Apostolov, L. G., 620. Appanna, M., 314. Apple, J. W., 275, 396. Arabadzhiev, D., 60. Araújo, J. B. Molinari, 555, 558. Archer, T. E., 92. Armstrong, M. T., 374, 694. Arndt, R. G., 270. Arnold, M. K., 287, 420. Arora, G. L., 174. Arrand, J. C., 394. Arru, G. M., 660. Arthur, A. P., 73, 157, 389. Arthur, B. W., 266, 475. Asatur, M. K., 360. Ascher, K. R. S., 613, 684. Asquith, D., 266. Atkins jr., E. L., 267. Atkins, J. G., 508. Atkins, M. D., 127, 458. Atwood, C. E., 163. Auclair, J. L., 453, 454. Audemard, H., 372. Auersch, O., 11. Avery, D. J., 671. Avidov, Z., 37, 611, 612, 613. Azevedo e Silva, J. de, 355.

Bache, C. A., 92. Badulin, A. V., 345. Baggiolini, M., 46, 86, 100, 552, 663. Bai, S. Godavari, 685. Bailey, J. C., 221, 406, 465.

Anderson, C. A., 88. Anderson, D. L., 475.

Anderson, L. D., 508.

Bailey, L., 617. Bailey, S. W., 632. Baker, A. H., 496. Baker, J. E., 414. Baker, J. J., 425. Baker, J. M., 496. Balachowsky, A. S., 491. Balâmír, S., 351. Balevski, A., 60, 111, 347. Ball, H. J., 518. Bally, W., 693. Balwant Singh, 88, 552. Bang (Yong-ho), 571. Banham, F. L., 424, 526. Banks, A. W., 504. Banks, C. J., 428. Baranówna, I., 365. Bardin, R., 480. Barlow, J. S., 389. Bärner, J., 248, 424. Barnes, H. F., 287, 420, 684. Barnes, O. L., 462. Baron, R. L., 566. Barrett, C. F., 524, 551. Barthel, W. F., 145, 483. Bartlett, B. R., 150, 464, 519. Bartlett, F. J., 639. Bartoloni, P., 137, 311, 535. Bates, A. N., 687. Batiashvili, I. D., 12. Batinica, J., 369. Batra, H. N., 498, 594. Bauman, L. F., 277. Baurant, R., 606. Baxter, G. D., 27. Beal jr., R. S., 17. Bean, J. L., 149, 411, 419. Beck, S. D., 4, 22, 68, 275. Becker, G., 489, 646, 647, 648, 649. Beckham, C. M., 152. Beckman, H. F., 562. Bedard, W. D., 386. Begg, J. A., 388. Beingolea G., O., 560. Béique, R., 392. Beirne, B. P., 90. Belin, M., 651. Belosel'skaya, Z. G., 359. Bénassy, C., 84, 379, 433, 434. Benkevich, V. I., 243, 597. Bennett, C. W., 17. Bennett, F. D., 289. Berck, B., 496. Berezina, V. M., 541. Berg, C. O., 91.

Bergen, P., 402. Bergman, P. W., 631. Beri, Y. P., 317, 500. Berim, N. G., 359. Berjon, J., 666. Berlin, J., 221. Berliner, E., 140. Bernard, J., 85. Bernegg, A. Sprecher von, 693. Beroza, M., 383. Bethe, J. G. C., 673. Bettai Gowder, R., 79, 594. Bevan, D., 550. Bevan, W. J., 288. Bhambhani, H. J., 118, 499. Bhatia, D. R., 172, 500. Bhatia, G. N., 116. Bhattacherjee, N. S., 315, 316, 498, 693. Bhattacherjee, S. P., 594. Bianchi, H., 379, 433, 434. Bibby, F. F., 221. Bigger, J. H., 420. Biggi, E., 643. Biliotti, E., 84, 434. Binaghi, G., 532. Binčev, B., 54. Bindra, O. S., 627, 688. Bird, F. T., 24, 25, 341. Bjegović, P., 32. Bjørnstad, A., 622. Black, L. M., 358. Blackith, R. E., 285. Blais, J. R., 25, 121, 157, 455. Blake, G. M., 249. Blanchard, E. E., 144, 552. Blanchard, R. A., 420. Bleak, E. E., 356. Blickenstaff, C. C., 277, 576, 694. Blinn, R. C., 267, 561. Blunck, H., 139, 683. Bobb, M. L., 335. Bobinskaya, S. G., 447. Boczek, J., 364. Bode, H. Maier-, **680**. Boer, J. A. de, 6. Bogavac, M., 31, 241. Bogdanov, V., 347, 541. Bognár, S., 307. Bohart, G. E., 341, 564. Böhm, H., 247, 302. Böhm, O., 184. Boldaruev, V. O., 13. Bollaerts, D., 374, 607. Bombosch, S., 376.

Bond, D. C., 686. Bond, E. J., 267, 356, 583. Bondarenko, N. V., 360. Bonfils, J., 372. Bonnemaison, L., 605, 669. Boock, O. J., 119. Booker, R. H., 531. Booth, C. O., 200. Borchardt, G., 106. Borg, A., 85, 354. Bornemissza, G. F., 452. Bosch, R. van den, 90, 151. Boselli, F., 47, 181, 182. Boswell, A. L., 264. Bottger, G. T., 478, 552. Bouček, Z., 305. Bouchard, P., 26. Bourguinet, P., 100. Bourke, T. V., 493. Bowen, M. F., 215. Bowery, T. G., 228, 331. Bowling, C. C., 329, 405. Boyce, H. R., 154. Bradley, R. H. E., 133. Brady jr., U. E., 475. Bragassa, C. B., 220. Brande, J. Van den, 607. Brann jr., J. L., 571. Brass, C. L., 470. Bravenboer, L., 682. Bray, D. F., 217, 333. Brazzel, J. R., 218, 220, 264, 330, 406, 471. Bremer, H., 425. Brénière, J., 250. Bressy, O., 86. Brett, C. H., 471. Brezner, J., 88. Briand, L. J., 162, 207. Brierley, P., 271. Briggs III, C. P., 219. Briggs, J. B., 670. Brimblecombe, A. R., 295, 589. Brindley, T. A., 329. Broadbent, L., 320. Brogden, C. A., 151. Brown, A. W. A., 128, 262, 284. Brown, E. S., 428, 484, 530, 655. Brown, L. C., 220, 632. Brown, N. R., 394, 522. Brown, R. G., 483. Browne, F. G., 296. Bruce, R. B., 483. Bruehl, G. W., 401. Bruel, W. E. van den, 374, 607, 683.

Bruggen, A. C. van, 531. Bruns, H., 141, 682. Bryant, D. G., 456. Bucher, G. E., 260, 342, 389. Buck, J., 90. Bugalho Semedo, C. M., 298. Buhr, H., 608. Bullock, J. A., 180, 686. Burbutis, P. P., 509, 511, 630. Burges, H. D., 6, 294. Burk, J. M., 24. Buro, A., 647. Burt, P. E., 320. Burton, B. T., 635. Burton, V. E., 331, 421. Busbey, R. L., 214. Busck, A., 121. Bushing, R. W., 248. Butcher, J. W., 129, 327, 410. Butler jr., G. D., 396. Butt, B. A., 338, 478, 479, 642. Büttiker, W., 262. Büttner, H., 252. Buttram, J. R., 266. Buturović, D., 240.

Cachan, P., 7. Cairaschi, E. A., 84, 435. Calica, C. A., 96. Calkins, C. O., 631. Call, F., 356. Callaham, R. Z., 642. Callahan, P. S., 563. Campbell, J. M., 410. Campbell, K. G., 32, 231. Campbell, W. V., 331, 566. Can, E., 43. Cangardel, H., 382, 383. Cann, F. R., 195. Canning, E. U., 581. Cantuel, J., 668. Capek, M., 602, 683. Caresche, L., 250. Carles, J. P., 8, 494. Carlson, R. B., 410. Carman, G. E., 561. Carne, P. B., 504. Carnegie, A. J. M., 87. Carpenter, C. P., 146. Carroll, W. J., 456. Carter, R. D., 496. Cartier, J. J., 453. Casida, J. E., 213. Casilli, O., 531.

Casimir, M., 33. Cass, L. M., 161. Castro, U. Paiva, 120. Caswell, G. H., 75. Cathey, G. W., 632. Catley, A., 235. Caussin, R., 85. Cavalli, L. M., 661. Cavanagh, L. A., 91. Čerkasov, J., 319. Cermáková, A., 196. Cessac, M., 604. Chaarawi, A. M., 180. Chaboussou, F., 8, 255, 312, 494, 665, 666, 667. Chadbourne, D. S., 629. Chadwick, P. R., 2. Chambers, H., 406. Champ, B. R., 589. Chancogne, M., 668. Chang, G. S., 624. Chang, G. Y., 428. Chang, H. G., 501. Chang (Hsueh-tsu), 592. Chang (Li-fung), 503. Chang, S. M., 501. Chant, D. A., 125, 378, 522, 528, 586. Chao (Chien-ming), 12. Chao (You-shin), 245. Chaplin, C. E., 570. Chapman, A. J., 334, 409. Chapman, J. A., 423, 424, 459. Chapman, P. J., 403. Chapman, R. K., 147, 516, 568. Charan Singh, 172. Chararas, C., 606. Chatterjee, P. N., 448. Chatterjee, S. N., 626. Chatterji, S., 196, 315, 626, 628. Chaudhuri, D. P., 315. Cheema, M. A., 175. Chen (Chun-mei), 360. Chen, K. H., 590. Cherblanc, G., 7. Cherkasskii, E. S., 675. Chiang, H. C., 398, 461. Chisholm, D., 83. Chisholm, R. D., 567, 578. Chiu (Shiu-chou), 591. Chiykowski, L. N., 686. Chokkanna, N. G., 448. Chow (Shao-lai), 592. Choyce, M. A., 53. Christensen, C. M., 414. Christidis, B. G., 352.

Chu, H. F., 624, 625, 693. Chu (Seng-fu), 502. Chung (Chi-chien), 593. Chung (Joh-min), 502. Church, N. S., 210. Chyi (Ruey-len), 593. Ciampolini, M., 309, 533. Claborn, H. V., 384. Clark, E. W., 334, 338, 475, 479, 569, 629. Clark, J. C., 263, 467. Clark, L. R., 262. Clark, R. C., 394, 522. Clark, W. H., 620. Cleveland, F. P., 384. Cloudsley-Thompson, J. L., 90. Coaker, T. H., 652. Cockbain, A. J., 203, 204, 320. Coffey, L. C., 579. Cohen, A. J., 321. Cohen, S., 614. Cohic, F., 81. Cole, J. H., 289, 553. Cole, J. R., 424. Cole, S. G., 510. Coles, L. W., 400. Colhoun, E. H., 62. Collingwood, C. A., 287, 432. Common, I. F. B., 232. Compere, H., 490, 688. Complin, J. O., 509. Concienne, E. J., 469. Condron, C. H., 573. Connell, W. A., 414, 419. Connola, D. P., 220. Converse, R. H., 132. Coon, B. F., 639. Coppel, H. C., 418, 692. Corley, C., 145. Corner, J., 394. Coscarón, S., 144. Costa, A. S., 480. Costa, J. Machado da, 357. Coudreau, J., 86. Coudriet, D. L., 408. Coulson, D. M., 91. Court, D., 668. Coutin, R., 84, 435. Coutinho, J. M., 120, 557, 558. Cowan jr., C. B., 408, 412. Cowley, G. T., 562. Cox, E. L., 463. Cox, J. A., 398. Cox, R., 384. Coxeter, W. J., 23.

Craig, C. H., 72. Craig, R., 90. Cramer, H. H., 683. Cranham, J. E., 448, 553. Creighton, C. S., 465. Cressman, A. W., 581. Crête, R., 455. Cristóbal, U. López, 299. Croix, E. A. S. la, 52. Crossan, D. F., 630. Crosskey, R. W., 293. Crow, J. F., 147. Crowe, G. B., 573. Crüger, G., 303. Cueto jr., C., 88. Cumber, R. A., 506. Cunningham, R. T., 571. Curry, A. N., 424. Curtis, R. W., 572. Cuscianna, N., 299. Cushman, H. G., 496. Cushman, R. A., 492. Cuthbert jr., F. P., 575. Cutkomp, L. K., 332, 466. Cwilich, R., 613, 680, 684. Czech, M., 10.

da Costa, J. Machado, 357.

Dadd, R. H., 5. Dafauce, C., 659. d'Aguilar, J., 7. Dahlsten, D. L., 23, 125. Dahm, P. A., 221, 225. Dale, J. E., 652. Daniélidou, R. K., 237. Daniels, N. E., 281. Das, M., 201, 202. Dastur, R. H., 139. Daubney, C. G., 177. Daugherty, D. M., 471. Dauterman, W. C., 146. Davich, T. B., 218, 224, 330, 404, 467, 472, 568. David, W. A. L., 51, 290, 488. Davidson, R. H., 470. Davies, D. E., 286. Davis, C. S., 276. Davis, D. W., 67. Davis, G. R. F., 495, 528. Davis, J. W., 408, 412. Dawsey, L. H., 479. Dawydko, B., 678. Day, M. F., 197, 357. de Almeida, P. R., 557.

Dean, H. A., 221, 406, 465, 564. Dean, R. W., 268. de Azevedo e Silva, J., 355. DeBach, P., 21, 356. de Boer, J. A., 6. Decker, G. C., 269, 301. Decker, R. W., 571. de Fluiter, H. J., 192, 608. de Gouveia, A. J. Soares, 99, 608. de Jong, D. J., 673. Delanoue, P., 435. de la Torre, O. Gamero, 560. Delmas, R., 434. de Luca, Y., 249. Delucchi, V. L., 186, 654. De Lyzer, A. J., 70, 458. Demečko, J., 678. de Mello, E. J. Rostom, 644. de Mol van oud Loosdrecht, W. E., Dempster, J. P., 30, 590, 620. Dennis, E. B., 430. Dennis, N. M., 571. Denton, R. E., 460. DePew, L. J., 468. Depner, K. R., 69. De Santis, E. Millán de, 144, 230. De Santis, L., 144, 230. Descamps, M., 620. Desmoras, J., 603. Destruel, C., 668. de Vries, J. E., 91. Dewey, J. E., 271. de Wilde, J., 6, 90, 204. Dezfulian, A., 428. Dhar, P., 499. Dias Netto, N., 555. Dickason, E. A., 211, 275, 417. Dicke, F. F., 152. Dicker, G. H. L., 287, 671. Dickson, R. C., 630. Diéguez, J. Martín, 355. Dietrick, E. J., 224. Dietvorst, F. C., 445. Dijck, W. Van, 374. di Martino, E., 299. Dimchev, V., 538. Dinther, J. B. M. van, 93. Dirimanov, M., 539. Ditman, L. P., 222, 325, 404, 630. Dittrich, V., 602. Dixon, J. C., 495. Dmitriev, G. V., 112. Doane, C. C., 641. Doane, J. F., 155, 516, 568.

Dobreanu, E., 608. Dodge, H. R., 24, 460. Dogger, J. R., 265. Dominik, J., 142, 189. Donker, J., 302. Dordević, M., 55, 368. Dorough, H. W., 266. Dorsey, C. K., 632, 636. Dosse, G., 302, 319. Doucette, C. F., 278, 517. Doull, K. M., 233. Doutt, R. L., 630. Dowden, P. B., 337, 399, 400. Downing, R. S., 395. Drew, W. A., 338. Drooz, A. T., 210, 418. Dubrovskaya, N. A., 13. Dudley, B., 496. Duffield, J. W., 478. Duffus, J. E., 131, 480. Dunn, C. L., 88. Dunning, R. A., 549. Duşchin, I., 57. Dustan, A. G., 229, 230. Dustan, G. G., 163, 404. Dutky, S. R., 324, 336. Dutt, N., 594. Düzgüneş, Z., 444. Dyer, E. D. A., 207, 424.

Earle, N. W., 407. Eastop, V. F., 357. Ebeling, W., 97, 496. Eckert, J. E., 342. Eden, W. G., 151, 227. Edgecombe, H. S., 49. Edwards, C. A., 430. Edwards, R. L., 127, 155, 575, 631. Egorov, N. N., 111, 597. Ehara, S., 452. Ehrhardt, P., 105, 443. Eisner, T., 90. Ekbom, P., 101. Eldefrawi, M. E., 227, 263. Elgee, D. E., 25. Eliseev, E. I., 423. Ellington, J. J., 226. Elmore, J. C., 636. El-Ziady, S.A.E., 30. Emden, H. F. van, 319, 320, 495. Emmel, L., 10. Engelhardt, N. T., 428. Eppley, R. K., 564. Erdös, J., 305.

Ericson, R. O., 301.
Erlich, S., 612.
Ernst, G., 552.
Erwin, D. C., 398.
Esau, K., 482.
Esenther, G. R., 416.
Evans, D. D., 140.
Evans, D. E., 429.
Evans, G. O., 41.
Evenhuis, H. H., 387, 673.
Eyles, A. C., 75, 506.
Evndhoven, G. L. van, 444.

Fadigas jr., M., 643, 645. Fahey, J. E., 570. Fahmy, H. S. M., 356. Fairchild, M. L., 225. Falanghe, O., 555. Falcon, L. A., 402. Farid-ud-Din, A., 196. Farrier, M. F., 327. Fedoseeva, L. I., 344, 564. Fenili, G. A., 310, 533. Ferguson, G. R., 149. Fernando, E. F. W., 448. Fernando, H. E., 484. Ferrière, C., 654. Ferry, G. V., 331. Ferwerda, J. D., 693. Fewkes, D. W., 248, 332. Fezer, K. D., 323. Fice, K., 369. Fieldhouse, D. J., 630. Fife, L. C., 403, 511. Filipescu, C., 56. Finlayson, T., 155, 209. Finley, L., 466. Finnegan, R. J., 457. Fiore, C., 462. Fisher, R. C., 618. Fisher, T. W., 150, 266. Fitze, K., 55. Fjelddalen, J., 622. Flanders, S. E., 150, 282. Fleming, G. A., 571. Fleschner, C. A., 378. Flock, R. A., 630. Floyd, E. H., 1, 402, 571. Fluiter, H. J. de, 192, 608. Fonseca, J. P., 559. Forbes, A. R., 692. Forbes, R., 574. Forsslund, K. H., 426. Forsythe jr., H. Y., 280, 399. Fort jr., S. W., 517. Fosset, J., 606. Fougerousse, M., 86. Fournel, J., 603. Fox, C. J. S., 73, 525. Fox, L. F., 573. Fraenkel, G., 552. France, H. L., 323. Franco do Amaral, S., 555, 644. Frank, M. B., 486. Frans, R. E., 220. Franssen, C. J. H., 192, 193. Franz, J., 184, 185, 186, 262, 377. Freeman, A. E., 225. Freeman, J. A., 236. Freeman, T. N., 121. Fremling, C. R., 262. French, R. A., 430. Frick, K. E., 299, 324, 564. Fritzsche, R., 547. Frye, R. D., 284. Fuchs, W. H., 9. Fudge, J. F., 562. Fudulov, D., 537. Fukuto, T. R., 267, 407, 634. Fulmek, L., 492. Fulton, H. G., 424. Fulton, R. A., 226, 324, 416, 418. Fye, R. E., 322.

Gabriel, W., 544. Gaines, J. C., 16, 578. Gair, R., 50. Gajewski, E., 188, 676. Galichet, P. F., 313. Gall, J. Le, 490. Galvez, G. E., 689. Gambaro, P. I., 662. Gambrell, F. L., 228. Gamero de la Torre, O., 560. Gangrade, G. A., 497. Ganhão, J. F. Pinto-, 617. Gara, R. I., 340. Garber, M. J., 398. García B., G., 559. Garcia, S. Planes, 98. Gardiner, B. O. C., 51, 290, 488. Garman, P., 694. Garner, C. F., 579. Gast, R. T., 225, 472. Gaud, S. M., 257. Gauss, R., 11, 375. Gehring, F., 254.

Gabai, N. M., 611.

Geier, P. W., 165, 262. Genung, W. G., 64. Georghiou, G. P., 515. Georgijević, E., 54, 369. Gerber, B., 664. Gerhardt, P.D., 473. Gerolt, P., 27. Gersdorf, E., 442. Gertler, S. I., 383. Gertych, Z., 676. Ghabn, A. A. A. E., 608. Ghani, M. A., 169. Giacomelli, R., 562. Giang, P. A., 196. Giannotti, O., 643. Gianotti, J. F., 144. Gibbons, C. F., 327. Giliomee, J. H., 650. Gilles, G., 445. Giray, H., 351. Giunchi, P., 182. Glass, E. H., 18, 147, 403, 633. Glasscock, H. H., 670. Glen, R., 692. Glick, P. A., 219, 334, 338, 476, 577. Glushchenko, A. F., 362. Godan, D., 442. Godavari Bai, S., 685. Goh Kee Guan, 78. Gojmerac, W. L., 573. Göksu, M. E., 177. Gołebiowska, Z., 364. González, B., J. E., 559. González Rodríguez, R. H., 495. Goode, J. P., 219. Goodhue, R. D., 495. Goos, A., 356, 599. Goossen, H., 10. Gopinath, K., 317. Gordon, H. T., 128. Gornostaev, G. N., 248. Gorz, H. J., 469. Gospodinov, G., 348. Gospodinov, M., 347. Gösswald, K., 104, 616. Gostick, K. G., 201. Gothilf, S., 612. Gotoh, A., 78, 449. Gould, G. E., 269. Gould, H. J., 431, 670. Gouveia, A. J. Soares de, 99, 608. Govind, R., 171. Govindan, M., 466. Gowder, R. Bettai, 79, 594. Graham, A. R., 693.

Graham, H. M., 219, 334, 417, 476. Graham, J. F., 530. Graham, M. W. R. de V., 493. Graham, W. M., 51. Granovsky, A. A., 552, 619. Graupner, A. J., 88. Grbić, V., 32. Grechkin, V. P., 116. Green, A. A., 28, 178. Green, N., 383. Gregory, G. E., 496. Gressitt, J. L., 166. Griffin, J. A., 567. Griffiths, K. J., 387. Grigor'eva, T. G., 363. Grigorov, S. P., 58, 59, 109, 110, 536. Grimal'skii, V. I., 598. Gris, S. Zimmermann-, 614. Grison, P., 262, 423. Grob, H., 100. Grobler, J. H., 458. Grosch, D. S., 90. Gross, A. E., 275. Groves, A. B., 20. Gruch, W., 140. Grujíc, B., 55, 370. Grujičić, G., 249. Gruszewska, M., 677. Guagliumi, P., 98. Guennelon, G., 372. Guignard, E., 664. Guimarães, J. A. Monteiro, 608. Gunn, D. L., 87, 98. Günthart, E., 682. Günthart, H., 682. Gunther, F. A., 267, 561. Günther, S., 105, 107. Guppy, J. C., 71, 129, 393. Gurland, J., 221. Gurney, B., 198. Gusev, G. V., 40. Gustafson, J., 453. Gusynina, L. M., 595. Guthrie, F. E., 152, 228, 566. Guthrie, W. D., 152, 215, 333, 476. Guyer, G., 16. Gyrisco, G. G., 92, 213, 280, 323, 330, 399, 561, 581.

Haarer, A. E., 196. Hab, J. K. Abul-, 278. Hacskaylo, J., 218, 224, 263, 467, 472. Hadi Ali, 317.

Hadžihalilović, H., 55. Hadžistević, D., 55, 138, 241. Hagel, G. T., 412. Hagen, A. F., 335. Hagen, K. S., 66, 90, 151. Hagley, E. A. C., 575. Haine, E., 444. Hall, I. M., 569. Hall, J. C., 67. Hall, V., 65. Hamilton, A. G., 40. Hamm, J. J., 284. Hamman, P. J., 406. Hammond, G. H., 619. Hammond, P. S., 651. Han, Y. F., 625. Handford, R. H., 158. Hanec, W., 4. Hans, H. (Mrs. D. Peschken), 93, 318. Haq, A., 499. Harcourt, D. G., 211, 385. Harding, J. A., 279, 477, 510. Hardman, J. A., 432. Harein, P. K., 215. Harpaz, I., 130, 349. Harper, A. M., 402, 468. Harries, F. H., 279, 480, 514. Harris, C. R., 210, 415. Harris, jr., E., 226. Harris, E. C., 489, 622. Harris jr., E. D., 275, 512. Harris, K. M., 292, 684. Harris, L. D., 330. Harris, P., **260**, **386**. Harris, R. V., 107, 546. Harris, W. V., 76. Harrison, B. D., 691. Harrow, K. M., 167. Harvey, G. T., 130, 580. Harvey, T. L., 575, 642. Harvey, W. R., 90. Harwood, R. F., 401. Hashmi, M. H. A., 317. Haskell, P. T., 423. Hassan, M. I., 180. Hastings, A. R., 513. Hastings, E., 461. Hawthorn, L. R., 341. Hay, C. J., 478. Hayman, B. I., 620. Haynes, D. L., 129, 413. Hazarika, S. H., 88. Heathcote, G. D., 320. Hedlin, A. F., 385.

Heidenreich, E., 11, 252. Heikinheimo, O., 101. Heikkenen, H. J., 481. Heimpel, A. M., 88, 161, 196, 282, 341. Heinrich, W. O., 558, 643. Heinze, K., 250, 254. Helle, W., 193, 688. Helson, G. A. H., 75. Hely, P. C., 35. Hemer, M., 45. Henneberry, T. J., 212, 264, 324, 477, 516, 633. Hensley, S. D., 469. Herbert, H. J., 122. Herdy, H., 260. Hering, E. M., 492. Hering, M., 108, 109, 255. Hernandez, D., 353. Heron, R. J., 124. Hervé, J. J., 669. Hervey, G. E. R., 633. Hetrick, L. A., 577. Hewlett, P. S., 3, 687. Heyne, E. G., 65. Hibbs, E. T., 395. Hickin, N. E., 321, 489. Hightower, B. G., 16. Hildebrand, E. M., 133. Hill, C. H., 20. Hill, E. G., 694. Hille Ris Lambers, D., 585. Hills, G. J., 690. Hills, O. A., 408. Hirano, C., 495. Hitchcock, S. W., 271, 407. Hobbs, G. A., 156, 210. Hochmut, R., 196. Hocking, B., 69. Hodek, I., 319. Hodgson, E., 640. Hodson, A. C., 212, 414, 467, 474. Hoffmann, A., 355.

Hofmaster, R. N., 335, 404.

Holdaway, F. G., 390. Hollingsworth, J. P., 219, 334.

Hopping, G. R., 390. Hoskins, W. M., 90, 128, 148, 248,

Holmes, N. D., 72, 156, 210.

Hogan, T. W., 296.

Hopkins, A. R., 322. Hopkins, T. L., 248.

Hough, W. S., 20, 637.

House, H. L., 157, 359.

263, 512.

Howden, G. F., 321. Howe, R. W., 486, 548. Howe, W. L., 631. Howitt, A. J., 510. Howland, A. F., 217. Hoy, J. M., 587. Hoyt, S. C., 466, 481. Hsu (Yu-fen), 501. Hu (Jia-qi), 501. Huddleston, E. W., 330. Hudon, M., 27, 258, 514. Huertos, M. Rubio-, 300. Huesmann, H., 304. Huffaker, C. B., 227. Huger, A., 44. Hughes, R. D., 39, 286. Hukusima, S., 35. Hunter-Jones, P., 40. Hurpin, B., 84. Husnain, A. Q., 176. Hussey, N. W., 198. Hyde, M. B., 177.

Ichinose, T., 79.
Ignoffo, C. M., 426, 427, 552, 619.
Iliev, I., 539.
Isac, G., 56.
Ishii, S., 495.
Itô, Y., 78, 449, 452.
Ivanov, S., 110.
Ivanović, J., 55, 368.
Ivany, J. J. H. Szent-, 234, 235, 300.
Ivey, M. C., 384.
Iwamura, M., 623.
Iwata, T., 78.

Jacobson, L. A., **640.** Jacobson, M., 635. Jahn, E., 100. Jamnický, J., 187. Janković, L., 55, 367, 370, 371. Janković, M., 54, 369, 370. Jannone, G., 532, 694. Janse, A. J. T., 235. Janzen, D. H., 262. Jaques, R. P., 426, 525. Jaumień, F., 545. Jayaraj, S., 424. Jen (Gai-hsin), **623.** Jennings, P. R., 689. Jensen, A., 354. Jeppson, L. R., 147, 509, 511. Jepson, W. F., 319, 320, 487.

Jermy, T., 305, 306. Jessep, C. T., 505. Jesser, M. J., 509, 511. Jester, K. G., 331. Jodko, J. Narkiewicz-, 366. Johansen, C. A., 412 Johansson, D., 101. Johnsen, R. E., 225. Johnson, C. G., 199. Johnson, E. R., 63. Johnson, J. E., 149. Johnson, N. E., 209, 259, 478. Johnson, R. B., 409. Johnson, V. A., 513. Johnston, C. J. R., 34. Johnston, C. O., 65. Johnston, H. G., 147. Joly, R., 194. Jones, D. W., 400. Jones, E. T., 65. Jones, G. D. G., 2. Jones, I. B., 286. Jones, P. A., 693. Jong, D. J. de, 673. Jørgensen, J., 608. Joseph, K. V., 499. Joshi, B. G., 117. Jotwani, M. G., 172, 317, 498, 627. Jourdheuil, P., 373. Juillet, J. A., 24, 73. Jukola, J., 194. Junnikkala, E., 194.

Kabiersch, W., 254. Kaitazov, A., 539. Kalandadze, L. P., 12. Kalandra, A., 196. Kalkat, G. S., 470. Kaloostian, G. H., 336, 424. Kalshoven, L. G. E., 42, 296, 618, 625. Kangas, E., 42. Kansu, A., 82, 300. Kanta, Shashi, 628. Kantack, E. J., 131. Karadzhov, S., 60. Karafiat, H., **184**, 185, 186. Kareem, A. Abdul, 424. Karman, M., 82. Karpova, A. I., 14. Kassab, A., 180. Kassanis, B., 690. Kasting, R., 527. Katiyar, K. P., 216.

Kazimi, S. K., 169. Kearns, C. W., 148. Keener, P. D., 408. Keller, J. C., 324, 338, 478, 479, 513. Kelsheimer, E. G., 64. Kenaga, E. E., 274. Kennedy, J. S., 200, 357. Kennett, C. E., 227. Kerr, W. E., 90. Kerrich, G. J., 52, 353. Kershaw, W. J. S., 200. Kettle, D. S., 90. Kevan, D. K. McE., 89, 686. Khan, Qadiruddin, 139. Khotyanovich, A. V., 495. Khristov, I., 109. Khristova, E., 540. Kinard, W. S., 465, 635. King, D. R., 579. Kinghorn, J. M., 207. Kinsey, M. G., 496. Kirchner, H. A., 440. Kirkov, K., 59. Kirkpatrick, R. L., 694. Kirkpatrick, T. H., 588. Kishaba, A. N., 572. Kittams, W. H., **461.** Kiyoku, M., **449.** Klee, O., 356. Kleinschmidt, R. P., 295. Klett, W. E. W., 681. Klindić, O., 240. Klingler, J., 109. Kloft, W., 44, 105. Klostermeyer, E. C., 264, 281. Klumpar, J., 444. Knight, F. B., 150. Knipling, E. F., 148. Knipling, G. D., 416, 418. Knoche, C., 494. Knorr, L. C., 80, 97. Koblitsky, L., 567, 578. Kockum, S. P. L., 51. Kodys, F., 445. Koehler, C. S., 280, 323. Koerber, T. W., 355. Kogan, M., 554. Kögler, E. Müller-, 44. Kokorin, A. N., 114. Kolb, A., **684.** Kolubajiv, S., 142. Komblas, K. N., 239. König, H., 444. Kononova, N. E., 12. Köppä, P., 41.

Korthals, A., 504. Körting, A., 253. Koślińska, M., 188, 675, 676. Kosmachevskii, A. S., 14. Kosov, V. V., 15. Koula, V., 247. Kovachevski, I., 345. Kráľovič, J., 187, 494. Kramer, J. P., 257. Krämer, K., 9, 441. Krantz, G. W., 149, 518. Krasinskaya, A. L., 343. Krause, G. F., 215. Krczal, H., 441, 546. Kress, L. M., 424. Krieg, A., 140. Kriegler, P. J., 298. Kring, J. B., 582. Krishnamurthy, K., 685. Křístek, J., 246. Krueger, H. R., 213. Krzeczkowski, K., 364. Kudler, J., 196, 600. Kuenen, D. J., 262. Kuitert, L. C., 64. Kulman, H. M., 212, 467, 474, 632. Kulunev, D., 347 Kumar, S. S., 299, 317, 500. Kuoh, J. L., 501 Kurosawa, T., **552.** Kurup, Annamma, 117. Kuschel, G., 1. Kushner, D. J., 580. Küthe, K., 679. Kuznetsov, V. I., 242. Kwei, C. M., 624.

Kopvillem, Kh. G., 244.

Laan, P. A. van der, 94. la Croix, E. A. S., 52. Lagace, C. F., 150. Laibach, E., 377. Laird jr., E. F., 630. Lajo Vasquez, L. E. R., 357. Lakocy, A., 363, 365. Lal, Rattan, 118, 316, 497, 499, 628. Lall, B. S., 173. Lamb, K. P., 587. Lambers, D. Hille Ris, 585. Lambert, J., 550. Lambremont, E. N., 407. Lamey, H. A., 132. Landis, B. J., 300, 412. Lange, R., 105.

Lange, W. H., 419, 421, 509. Large, J. R., 424. Latif, A., 175, 176, 626. Lattin, J. D., 211. Lavabre, E. M., 355. Lavery, H. J., 589. Leach, C. M., 275. Lee (Hong Suk), **464.** Lee, I., 221. Lee (Sung-yang), 328. Lee (Tsui-ying), 623. Lees, A. D., 688. Lefkovitch, L. P., 656, 658. Le Gall, J., 490. Legner, E. F., 415. Legowski, T. J., 431, 432. Leiderman, L., 120, 644. Leius, K., 208, 391. Lejeune, R. R., 126. Lekić, M., 138, 241. Lemarie, J., 246. Lenarduzzi, R., 183. Leonard, D. E., 400. Leonide, J. C., 313. LeRoux, E. J., 159, 258, 455. Leroy, P. Péreau-, 54. Łeska, W., 677. Łeski, R., 61, 188, 189, 544, 676, 677. Lespès, L., 383. Leuchs, F., 303. Levin, M. D., 341. Lhoste, J., 262, 550. Li (Lung-shu), 693. Li (Ya-tse), 593. Liao (Shun-yuan), 693. Lichtenstein, E. P., 145, 146, 272, 415, 417, 483, 562, 572. Liebermann, J., 356. Lieftinck, M. A., 496. Liesering, R., 107. Lilly, C. E., 402. Lilly, J. H., 475. Lincoln, C., 220, 632. Lindgren, D. L., 331, 574, 619. Lindquist, D. A., 218, 263, 386, 467, 472, 568. Lin Yu, 501. Lipowa, I., 365. Lisk, D. J., 424. Liu (Ching-fu), **592.** Liu (Yu-ch'iao), **623.** Llavería B., M., 559. Lloyd, C. J., 3, 687. Lloyd, D. C., 199, 586. Lloyd, E. P., 410, 473, 573.

Loan, C., 390, 414. Löcher, F. J., 440. Lodos, N., 351. Loebenstein, G., 130. Lofgren, C. S., 464, 479, 483, 639. Loher, W., 423. Long, W. H., 214, 216, 280, 469. Loosdrecht, W. E. de Mol van oud, 42. López Cristóbal, U., 299. Lord, F. T., 259, 523. Lovato, A., 182. Lovisolo, O., 80. Löw, I., 318. Lowe, A. D., 74, 505, 620. Lowry, W. L., 409, 473. Lozinskii, V. A., 598. Lu (Lien-kao), 693. Luca, Y. de, 249. Lucas, S., 86. Lüdemann, D., 304. Lüdicke, M., 43. Ludwig, D., 462. Luitjes, J., 302. Lukefahr, M. J., 517, 567. Lund, A. E., 428. Lung (Cheng-te), 591. Lupo, V., 299. Luteršek, D., 54, 369. Lyon, R. L., 248. Lyons, L. A., 459. Lyubenov, Ya., 109, 348, 537. Lyzer, A. J. De, 70, 458.

McAlpine, J. F., 128. Macan, T. T., 90. McCalley, N. F., 509. MacCarthy, H. R., 691. McClanahan, R. J., 574. McCombs, C. L., 471. McCormick, W. J., 469. McCoy, C. E., 329. MacCreary, D., 419. MacCuaig, R. D., 164, 201, 291, 657. MacDougall, D., 88. MacEachern, C. R., 83. McEnroe, W. D., 564. McEwen, F. L., 342. Macfarlane, D., 41. McGarr, R. L., 409. McGinnis, A. J., 527. McGugan, B. M., 692. Machado da Costa, J., 357. McIntosh, A. H., 202.

McIntyre, T., 336. Mack, G. L., 227, 228. Mackauer, M., 673. MacKinnon, J. P., 584, 690. McLean, D. L., 467, 496. MacLeod, C. F., 282. MacLeod, D. M., 458. McLeod, J. H., 692. McLeod, J. M., 258. McMeans, J. L., 410. McMillian, W. W., 132. McMullen, L. H., 458. McMurtry, J. A., 640. MacNay, C. G., 528. MacPhee, A. W., 83, 159, 528. Madjar, V. Melamed-, 613. Madsen, H. F., 158, 402, 411, 481. Maelzer, D. A., 165, 504. Mago Navarrete, P., 357. Magor, R. D., **636.** Maier-Bode, H., 680 Major, R. T., 577. Majumder, S. K., **685.** Makarov, M., 537, 538. Maksimović, M., 31, 54, 55, 368, 369. Maldague, M. E., **609**. Malek, A. A. Abdel-, 686. Mallis, A., 635. Mallo, R. G., 694. Malo, F., 267, 274. Manglitz, G. R., 469, 631. Mann, H. D., 384. Manninger, G. A., 306. Manolache, C., 608. Manser, P. D., 289. Manson, G. F., 574. Maramorosch, K., 96. Marble, V. L., **421.** March, R. B., 148. Marek, J., 92, 492. Mariconi, F. A. M., 120. Marikovskii, P. I., 542. Marinari, A., 310, 533. Markelova, E. M., 15. Markelova, V. P., 360. Markkula, M., 41, 102. Markov, M., 345. Marović, R., 54. Marsh, R. W., 301. Marshall, A. T., 321. Marshall, J., 394, 411. Martelli, G. M., 38. Martelli, M., 660. Martignoni, M. E., 281, 496. Martin, D. F., 517.

Martin, Hubert, 301. Martin, W. J., 131. Martín Diéguez, J., 355. Martineau, R., 455. Martinek, V., 601. Martinez, J. B., 300. Martino, E. di, 299. Martorell, L. F., 257. Martyn, É. J., 424. Marzsó, L. Szalay-, 307. Mashhood Alam, S., 91. Maskell, F. E., 50.
Masner, L., 43.
Massey, C. L., 460, 496.
Mateo-Sagasta, E., 99. Mathur, R. N., 88, 552, 620. Mathys, G., 664. Matsui, S., 622. Matsumoto, S., 552. Matthysse, J. G., 18. Mattingly, P. F., 90. Maw, M. G., 123, 130. Maxwell, C. W., 71, 273. Maxwell, F. G., 551, 629. May, A. W. S., 40. May, B. M., 231, 424, 505. Mayer, M. S., 471. Mazurek, J. H., 210. Meagher, W. R., 88. Medler, J. T., 160. Mehrotra, K. N., 40. Meiffren, M., 651. Melamed-Madjar, V., 613. Melis, A., 135, 308, 311, 532. Mello, E. J. Rostom de, 644. Mellor, F. C., 692. Meltzer, J., 445, 685. Menn, J. J., 512. Menon, M. G. Ramdas, 173, 196, 316, 317, 626, 693. Menzel, D. B., 227. Merino M., G., 353. Merker, E., 185, 252. Merkl, M. E., 410, 473, 515, 573. Merwe, J. H. van der, 694. Merzheevskaya, O. I., 620. Meshcheryakova, L. V., 192. Messenger, P. S., 563. Metcalf, R. L., 147, 407, 508, 515, 634. Metivier, J., 603. Metzer, R. B., 579. Meyer, M. K. P., 83.

Meynadier, G., 606.

Michaeli, D., 248.

Michelbacher, A. E., 276. Michener, C. D., 234. Middelem, C. H. Van, 64. Mijin, K., 31. Milaire, H., 433, 434. Miles, J. R. W., 574. Miles, J. W., 561. Milević, K. S., 55. Millán de De Santis, E., 144, 230. Miller, A. C., 635. Miller, L. A., 70. Miller, W. E., 19, 413, 481, 495. Millers, I., 281. Millington, W. F., 562. Milne, D. L., 650. Milstead, J. E., 281, 496. Miskus, R., 227, 248. Misra, C. P., 414. Mitchell, B., 286. Mitchell, R. G., 209. Mitchell, W. G., 145. Mittelstaedt, H., 90. Mittler, T. E., 322. Miyashita, D. H., 383. Miyashita, K., 78, 449. Mizuta, K., 450. Moens, R., 683. Moericke, V., 200, 300, 423. Moffitt, H. R., 271. Moiseeva, T. S., 114. Molatore, H. D., 259. Molinari Araújo, J. B., 555, 558. Mol van oud Loosdrecht, W. E. de, 42. Monro, H. A. U., 267. Monteiro Guimarães, J. A., 608. Mookherjee, P. B., 500. Moore, A. D., **577.** Moore, K. M., 232. Moore III, S., 269. Moorefield, H. H., 148. Moravskaya, A. S., 190. Moreton, B. D., 670. Morettini, A., 693. Morgan, D. G., 483. Morgan, L. W., 152. Morimoto, N., 450, 451. Morris, O. N., 519, 581. Morrison, F. O., **453.** Morrison, L., 505. Morse, R. A., 276. Moussa, M. A., 149. Mrozowska, T., 188, 676. Mueller, C. H., 572. Mueller, W. C., 132.

Muhammad, W., 626. Muir, R. C., 672. Mukerjea, T. D., 171, 288. Müller, E. W., 439, 440. Müller, F. P., 494. Müller, H. J., 423, 546. Müller, R., 552. Müller-Kögler, E., 44. Mullins, A. M., 280. Mumford, B. C., 299. Muñiz Velez, R., 356. Murphy, E. F., 228. Murphy, R. T., 145, 483. Musgrave, A. J., 128. Musil, M., 304. Muthu, M., 424. Muthuswamy, R., 297. Myrdal, G. R., 572. Myser, W. C., 478.

Naber, E. C., 326. Nachev, P., 60, 110. Nadiradze, N. V., 12. Naegele, J. A., 272. Nagarajan, K. R., 79. Nagasawa, S., 248. Nagy, B., 306. Naidu, M. B., 117. Nair, M. R. G. K., 314. Nairn, L. D., 456. Naito, A., 424, 496. Nakata, S., 463. Nallini Wickramasinghe, 484. Namba, R., 482. Narkiewicz-Jodko, J., 366. Nash, R. W., 121. Nath, S., 174. Naton, E., 184. Navarrete, P. Mago, 357. Naylor, A. F., 620. Nebieridze, É. Ya., 12. Needham, P. H., 300. Neiswander, R. B., 19, 152, 573. Nenonen, M., 194. Nerney, N. J., 224. Netto, N. Dias, 555. Neumann, H., 304. Neunzig, H. H., 323. Neury, G., 663. Newsam, A., 76. Newsom, L. D., 1, 131, 280. Newton, A. P., 449. Nicholls, C. F., 25, 551. Nicholson, A. J., 128.

Nickel, J. L., 462. Niedermeier, R. P., 146. Nielsen, A. F., 354. Nielson, M. W., 356, 462. Niemczyk, E., 676. Niklas, O. F., 140. Nikolova, V., 346. Nirula, K. K., 448. Nishijima, Y., 96. Nitsche, R., 489. Nitzany, F. E., 614. Nizamlioğlu, K., 177, 236, 237, 352, 495, 552. Noble, L. W., 334, 511, 577. Nobrega, S. A., 119. Noll, J., 139. Nonveiller, G., 54, 181, 242. Norato, G., 300. Norris jr., D. M., 418, 619. Nye, W. P., 341.

Oatman, E. R., 415, 417. O'Brien, R. D., 146, 301. Obrtel, R., 246, 445. Ochs, G., 303. Ogloblin, A., 144. Ohashi H., J., 559. Ohmori, Y., 622. Ohnesorge, B., 141, 299. Okumura, G. T., 563. Oldiges, H., **376.** Oliver, A. D., 466. Onsager, J. A., 330. Oommen, C. N., 499. Orlando, A., 119, 120, 555, 556, 557, 558, 643, 646. Orlob, G. B., 95, 133, 160, 584. Orphanidis, P. S., 237, 238, 239. Orr, C. W. M., 128. Osgood, C. E., 325. Osgood, E. A., 495. Osman, M. F. H., 30. Osmolovskii, G. E., 362. Otake, A., 451. Otero M., G., 557. Otto, D., 250. Ottolini, P., 661. Oudshoorn, W., 193. Ouye, M. T., 642. Özer, M., 551. Ozols, G. E., 115.

Pableo, G., 96. Pařkin, D. M., 543.

Paine, R. W., 353. Painter, R. H., 551, 629. Paiva Castro, U., 120. Pajni, H. R., 173. Pali, N. S., 500. Paliř, V. F., 191 Palmer, J. G., 324. Palmer, T. P., 167. Palmiter, D. H., 23. Pan (Yung-song), 214. Pant, C. P., 36. Pao, S. T., 624. Papillon, M., 285. Papp, C. S., 420. Papworth, D. S., 195. Paradis, R. O., 26. Parencia jr., C. R., 408, 412. Parent, B., 454. Parfent'ev, V. Ya., 112. Park, T., 486. Parkhe, D. P., 117. Parshad, B., 173. Pasarela, N. R., 483. Paschke, J. D., 284. Pastrana, J. A., 230. Paszek, E. C., 513. Patel, H. K., 236. Patel, N. G., 332. Patel, R. C., 236. Patel, R. M., 236. Paternotte, E., 302. Paterson, E. C., 80. Pavan, M., 615. Pavlov, I. F., 243, 596. Paylor, R. A. L., 424. Payne, D. W., 496. Pearce, G. W., 561. Pearson, A. J. A., 165, 608. Pegazzano, F., 310. Pei (Sze-wan), 501. Peiris, J. W. L., 81. Peiu, M., 56. Pen (Chun-yun), 362. Pence, R. J., 339. Pendery, W. E., 421. Pener, M. P., 142. Pengelly, D. H., 161. Peoples, S. A., 384. Pepper, J. H., 461. Péreau-Leroy, P., 54. Pereira, A. Lima G., 646. Perkins, J. F., 493. Perron, J. P., 455. Perry, A. S., 64, 148. Peschken, Mrs. D. (see Hans, H.).

Pessenda, C. E., 557. Peswani, K. M., 498, 500. Petersen, B. B., 674. Peterson, L. K., 72. Petrova, I. A., 114. Pfrimmer, T. R., 515. Phillips, A. M., 424. Phillips, D. D., 561. Phillips, J. H. H., 164, 470. Phillips, L., **466.** Phytisas, E. A., 237, 238. Pickens, L. G., 576. Pickett, A. D., 262. Pickthall, J., 496. Pielou, D. P., 388, 395, 524, 551. Pieprzyk, W., 365. Pierce, H. D., 420. Pierce, W. D., 518. Pierrard, G., 606. Pietermaat, F., 374. Pigatti, A., 145, 556, 644, 646. Piltz, H., 10. Pimentel, D., 68, 401, 427. Pinto-Ganhão, J. F., 617. Pires, J. A., 551. Planes Garcia, S., 98. Plant, H. N., 350. Pleasance, H. J., 291. Plenet, A., 88. Poinar jr., G. O., 576, 581. Pointing, P. J., 392. Pollard, G. E., 561. Pollard, H. N., 336. Polnik, A., 89. Polon, J. A., 620. Polyakov, I. Ya., 15. Pond, D. D., 69, 633. Ponsen, M. B., 691. Popov, P., 59. Popov (Popow), P. A., 355, 366. Popov, V., 537. Popov, V. I., 57. Popova, E. A., 345. Popova, I., 60. Popova, V., 58, 349. Pospelov, S. M., 362. Post, A., 444. Postner, M., 251, 375. Potter, H., 299. Povolný, D., 235. Pradhan, S., 118, 170, 172, 316, 317, 498, 499, 627. Pranter, W., 43, 356. Prasad, A. R., 119, 317. Prasad, S. K., 171, 172, 500, 627.

Prekajski, P., 55. Prentice, I. W., 107, 546. Prentice, R. M., 456, 619. Prescott, H. W., 134, 496. Prevett, P. F., 49. Price, D. W., 405. Price, R. G., 338. Prilop, H., 255. Prints, E.Ya., 543. Prints, Ya.I., 543. Pritpal Singh, 500. Proctor, J. H., 80, 486, 551. Pruess, K. P., 214. Pruthi, H. S., 594. Pschorn-Walcher, H., 184, **185**. Pussard, R., 436, 606. Putman, W. L., 521. Puttler, B., 66, 400. Puzzi, D., 120, 555, 557, 558, 646.

Qadiruddin Khan, 139. Qayyum, A., **175.** Quinn, D. O., **636.**

Raatikainen, M., 102. Rabb, R. L., 152. Race, S. R., 409. Radeleff, R. D., 384. Rademacher, B., 425. Radinovsky, S., 518. Rahalkar, G. W., 497, 628. Rai, B. K., 172, 317, 628. Raine, J., 395. Rainey, R. C., 423. Rainwater, J. C., 418. Raj, P. J. Sanjeeva, 170. Rajagopalan, K. V., 41. Rajani, V. G., 553. Ramadier, P., 666, 667. Rambier, A., 445. Ramchandra Rao, Y., 315. Ramdas Menon, M. G., 173, 196, 316, 317, 626, 693. Rammer, I. A., 572. Ramsfjell, T., 622. Randolph, N. M., 219, 578. Ranney, C. D., 218. Rao, B. R. Subba, 299, 317. Rao, D. G., 80. Rao, K. Siva, 118. Rao, V. P., 139, **169.** Rao, Y. Ramchandra, 315. Rasa, E. A., 482.

Ratcliffe, R. H., 222. Rattan Lal, 118, 316, 497, 499, 628. Raun, E. S., 284, 343. Ravelli, V., 136. Rawlins, W. A., 226, 470. Razowski, J., 189. Reed, D. K., 269. Reed, W., 53. Reeher, M. M., 134. Rehman, M. H., 169. Reid, W. W., 327. Reiner, C., 580. Rennels, R. G., 153. Rennison, B. D., 490. Reyne, A., 143. Reynolds, H. T., 148, 277, 284, 398, 508. Řezáč, M., 301. Rhynehart, T., 178. Rich, E. R., 621. Richards, O. W., 426. Richards, W. R., 584. Richardson, C. D., 517. Richardson, H. P., 524. Richmond, C. A., 334, 479, 569. Richter, G., 439. Ricou, G., 373. Ridgway, R. L., 213. Riehl, L. A., 221. Rinick jr., H. B., 639. Ripper, W. E., 148. Risbec, J., 653. Ris Lambers, D. Hille, 585. Rivard, I., 583. Rivera, C. T., 638. Rivers, C. F., 690. Riviello, M. S., 325. Rivnay, E., 351, 614. Robbins, A. J., 483. Robbs, C. F., 559. Robert, A., 455. Robert, P. C., 683. Roberts, D. W. A., **526.** Roberts, J. E., 567. Roberts, R. B., **620**. Robertson, O. T., 334, 417. Robertson jr., R. S., 264. Robinson, A. G., **525.** Robinson, J. R., 88. Rochow, W. F., 132, 134. Rodionova, L. Z., 344. Rodriguez, J. G., 273, 410, 570. Rodríguez, R. H. González, 495. Roehrich, R., 84, 256, 664. Roff, C., 295.

Roland, G., 375. Rollins, L. E., 157. Rolston, L. H., 65, 421. Romankow, W., 365. Romanyk, N., 381, 659, 660. Romashov, D. D., 355. Roonwal, M. L., 448. Ross, A. F., 132. Ross, D. A., 457. Ross, H. H., 462. Rossetti, V., 556. Rostom de Mello, E. J., 644. Rota, P., 663. Roth, L. M., 90, 520, 565. Rouse, P., 65, 421. Roussel, J. S., 147. Rout, G., 118. Rozhkov, A. S., 344. Rubenstein, D., 92. Rubio-Huertos, M., 300. Rubtsov, I. A., 596. Rubtsova, N. N., 597, 599. Rudd, R. L., 262. Rudinsky, J. A., 90, 209. Ruffinelli, A., 643. Rühm, W., 682. Rumphorst, H., 10. Rupérez Cuéllar, A., 381, 550, 660. Ruppel, R. F., 560. Rusk, H. W., 570. Russ, K., 85, 248. Russell, M. P., 337. Russo, G., 46, 438, 694. Ruszkowski, A., 364. Ryke, P. A. J., 83.

Saba, F., 428.
Sagasta, E. Mateo-, 99.
St. John jr., L. E., 424.
Saito, S., 622.
Salas, L., 560.
Salatić, S., 54.
Salazar T., J., 560.
Salibe, A. A., 556.
Salonen, K., 42.
Salt, R. W., 551.
Samšiňáková, A., 196.
Sanders, W., 255.
Sandner, H., 205.
Sands, W. A., 293, 495.
Sanford, K. H., 159.
Sanjean, J., 397.
Sanjeeva Raj, P. J., 170.

Rymashevskaya, R. S., 242.

Sans, W. W., 574. Sant, L. E. van't, 299, 608, 673. Santi, R., 562. Santis, E. Millán de De, 144, 230. Santis, L. De, 144, 230. Santoro, R., 46. Sáringer, G., 248, 307, 308. Sarma, D. S. R., 41. Sarma, P. S., 41. Sarup, P., 196, 315, 316, 498, 626, 627. Sastry, K. S. Shivashankara, 314. Satpathy, B. N., 265, 497. Saunders, G. W., 503. Saunders, J. L., 619. Săvescu, A., 56, 57. Savić, G., 32. Savoĭskaya, G. I., 243. Sawyer jr., E. W., **620**. Sawyer, K. F., 291. Saxena (Sexena), A. P., 497. Saxena, K. N., 628. Saxena, P. N., 497, 628. Saxena, S. C., 117. Sbur, D. E., 212, 270. Schaefers, G. A., 23, 567. Schäller, G., 95. Schechter, M. S., 324. Scherney, F., 494, 683. Schetters, C., 186. Schlabritzky, E., 681. Schlinger, E. I., 66, 67, 151. Schmidt, G., 104, 254. Schmidt, L., 367, 368. Schmitt, J. B., 90. Schmutterer, H., 443. Schneider, F., 90, 99. Schoch, H., 552. Schoonhoven, L. M., 205. Schopp, R., 300, 412. Schreurs, J., 75. Schulz, J. F., 330. Schulz, J. T., 395, 396. Schulz, K. R., 146, 272, 572. Schuster, M. F., 406. Schuster, R. O., 521. Schütte, F., 106. Schvester, D., 83, 84, 372. Schwartz, E., 440. Schwartz jr., P. H., 325. Schwenke, W., 251. Schwerdtfeger, F., 140, 141, 248. Scognamiglio, A., 437. Scott, H. E., 152. Scriven, G. T., 640.

Šedivý, J., 445. Seeger, J. R., 609. Seifert, G., 679. Sekhar, P. S., 37. Sellers, W. F., 229. Semedo, C. M. Bugalho, 298. Semel, M., 328. Sen, A. C., 315. Sengalevich, G., 59, 60, 539. Sengupta, G. C., 265. Sethi, G. R., 497, 628. Sevintuna, C., 82, 128. Sexena, A. P. (see Saxena, A. P.). Sferra, P. R., 339. Shaffer, C. B., 483. Shahwan, A. M., 180. Shaikh, F. R., 176. Shalapenok, E. S., 355. Shands, W. A., 460, 569. Shankland, D. L., 572. Shapiro, I. D., 495, 595. Shapiro, M., 427. Shaposhnikov, G. Kh., 423. Sharifullah, M., 576. Sharma, G. C., 500. Shashi Kanta, 628. Shaw, J. G., 279, 325. Shaw, M. W., 197. Sheals, J. G., 41. Sheets, L. W., 638. Shekhurina, T. A., 115. Shen (Kuang-pu), 623. Shenefelt, R. D., 416. Shepherd, R. F., 208. Shi, N., 265, 497. Shiller, I., 511. Shivashankara Sastry, K. S., 314. Shmal'ko, V. F., 675. Shnaider, E. V., 541. Shorey, H. H., 216, 476, 508, 569. Shotwell, R. L., 520. Shteinberg, D. M., 87. Shuel, R. W., 90. Shulov, A., 142. Shutyaev, A. M., 191. Siddiqi (Siddiqui), Z. A., 119, 497, 553. Sillar, D. I., 589. Sillibourne, J. M., 671, 672. Silva, J. de Azevedo e, 355. Silva, P., 356. Silver, G. T., 126, 395, 457. Silvestre, P., 298. Simanton, W. A., 552.

Simmonds, F. J., 229.

Simon F., J. E., 559. Simons, J. N., 635, 636. Simpson, G. W., 460, 569. Sinadskiĭ, Yu.V., 354. Sinclair, W. B., 574. Singh, Balwant, 88, 552. Singh, M. P., 315. Singh, O. P., 119, 553. Singh, Pritpal, 500. Singh, R. P., 119. Singh, S. R., 88. Sinha, R. N., 154, 424. Sinitsina, L. P., 113. Sippell, W. L., 124, 140. Sirotina, M. I., 447. Sisojević, P., 55, 370, 371. Siva Rao, K., 118. Skorikova, O. A., 355, 361. Słabecka, D., 61. Sleesman, J. P., 328, 397, 413, 463. Sloan, M. J., 226. Slonovskii, I. F., 543. Slykhuis, J. T., 70, 523, 684. Smirnoff, W. A., 53, 73, 282, 341, 382, 580. Smissman, E. E., 22, 68. Smith, C. F., 144, 152. Smith, E. H., 216, 326, 470, 631. Smith, F. F., 147, 212, 264, 271. Smith, H. S., 586. Smith, J. N., 90, 321. Smith, K. M., 581, 690. Smith, L. B., 566, 658. Smith, L. M., 384. Smith, R. F., 90. Smith, R. H., 223. Smith, R. W., 160. Smith, W. A., 503. Smithers, C. N., 38, 48, 88. Smock, R. M., 403. Smolarzowa, S., 188. Snapp, O. I., 226, 641. Snyder, T. E., 300. Soares de Gouveia, A. J., 99, 608. Soderstrom, E. L., 424. Soenen, A., 302, 445. Sokolov, A. M., 191. Solomon, J. D., 418, 620. Solomon, M. E., 91, 549. Soloway, S. B., 561. Solozhenikina, T. N., 597. Sonleitner, F. J., 621. Soueref, S. T., 239. Southwood, T. R. E., 291, 319, 320, 487.

Spaić, I., 55.

Sparks, A. N., 478, 552.

Spencer, E. Y., 88.

Spiller, D., 231.

Spink, W. T., 265.

Splittstoesser, C. M., 342.

Spradbery, J. P., 426.

Sprecher von Bernegg, A., 693.

Srinivasan, P. M., 79, 594. Sripathi Rao, B., 76.

Srivastava, B. K., 119.

Srivastava, P. D., 499.

Šrot, M., 600.

Stafford, E. M., 278, 572.

Stamato, W., 558. Stanić, V., 55, 368.

Stanković, A., 445.

Stapel, C., 139. Stapley, J. H., 651.

Stark, R. W., 125, 586.

Staub, A., 664.

Stefanović, D., 55, 370.

Steger, O., 106. Stegwee, D., 691.

Stein, W., 376, 381, 494, 680.

Steiner, H., 376, 682.

Steiner, L. F., 383, 643.

Steiner, P., 140.

Steinhauer, A. L., 638.

Steinhaus, E. A., 90. Stephen, W. A., 640.

Stephen, W. P., 140, 411, 564. Stephens, C. S., 637.

Stephens, J. M., 620. Stephens, S. G., 464.

Stephenson, H., 140.

Stern, V. M., 90.

Sternburg, J., 63.

Steudel, W., 253, 545. Steve, P. C., 475.

Stevenson, A. B., 404. Stevenson, J. O., 216.

Stewart, D. K. R., 259.

Steyvoort, L. Van, 249.

Storey, I. F., 670.

Stower, W. J., 286.

Stringer, C. E., 464, 639.

Stringfield, G. H., 215, 333.

Strong, F. E., 564.

Strong, F. M., 483.

Strong, R. G., 212, 226, 270, 331.

Stubbs, L. L., 34.

Stultz, H. T., 523.

Stürckow, B., 318.

Subba Rao, B. R., 299, 317.

Subramanian, T. R., 314.

Sullivan, C. R., 206.

Sullivan, W. N., 226, 324, 416, 418.

Summers, F. M., 405.

Sun (Yun-pei), 63, 397.

Suomalainen, E., 90.

Suplicy filho, N., 645, 646.

Suski, Z. W., 675, 677, 678. Svoboda, W. A., 227.

Swailes, G. E., 156, 210, 261.

Sweeney, R. C. H., 352. Sweet, R. C., 220.

Swirski, E., 614, 684.

Sykes, R. L., 49.

Sylvester, E. S., 322, 331, 463, 688, 691.

Syme, J., 366.

Symmons, P. M., 98.

Sytenko, L. S., 112.

Szalay-Marzsó, L., 307.

Szczygieł, A., 189, 676.

Szczypiński, W., 61.

Szelényi, G., 305.

Szent-Ivany, J. J. H., 234, 235, 300.

Szmidt, A., 377, 379. Szweda, M., 677.

Tadić, M. D., 54, 367, 682.

Tahori, A. S., 684.

Talhouk, A. S., 601, 608.

Tanada, Y., 196, 283, 428, 463, 580. Taneichi, K., 35.

Tang (Peng-gi), 591.

Tanskiĭ, V. I., 343. Tanton, M. T., 495.

Tao (Chia-hwa), 590.

Tapley, R. G., 652.

Taschenberg, E. F., 228.

Taylor, C. E., 38.

Taylor, E. A., 212, 264, 324, 477, 633.

Taylor, J. M., 489.

Taylor, K. L., 552. Taylor, L. R., 29, 199.

Telford, A. D., 123, 221.

Temlová, B., 196.

Templado, J., 300, 355.

Teofilović, Z., 32.

Terëkhin, É. S., 363.

Terriere, L. C., 417. Terrosi, U., 616.

Tešić, T., 32.

Tetefort, J., 297. Tew, R. P., 671, 672.

Thakur, C., 119, 171.

Thalenhorst, W., 107. Thapa, R. S., 448. Theden, G., 489. Thiele, H. U., 377. Thielemann, R., 545. Thomas, C. R., 421. Thomas, J. B., 122, 260. Thomas, J. G., 264. Thomas, R. T. S., 75, 76, 196. Thompson, L. S., 470. Thorsteinson, A. J., 318. Thresh, J. M., 651 Thurston, H. D., 689. Thurston, R., 405. Tietz, H. J., 577. Tiittanen, K., 103, 674. Timmerman jr., J. A., 266. Tinnilä, A., 102. Tjoa Tjien Mo, 143. Tokmakoglu, C., 81. Tomaszewski, W., 356. Tominić, A., 239, 609. Tomlinson jr., W. E., 337. Tonks, N. V., 395. Toole, E. H., 341. Torre, O. Gamero de la, 560. Torrent, J. A., 659. Torres, S. C. A., 557. Touhey, J. G., 217. Townsend, G. F., 90. Travis, R. V., 264. Treece, R. E., 18, 323. Trehan, K. N., 173. Tremblay, E., 436, 437. Treon, J. F., 384. Tsai (Pang-hua), 593, 623. Tsakmakis, A. A., 237, 238. Tsao, C. H., 473, 475. Tsitsin, N. V., 675. Tsugawa, C., 35. Tsuji, H., 451. Tsukerman, E. A., 15. Tsung Lin, 591. Tsvetkov, D., 347. Tsyplenkov, E. P., 13. Tubeuf, C. von, 251. Tuft, T. O., 483. Turley, D. L., 473. Turnbull, A. L., 586. Turner, A. J., 232. Turnock, W. J., 125. Tuttle, D. M., 396. Tyler, P. S., 582. Tyrrell, C., 526.

Uberoi, N. K., 626. Ueno, H., 450. Ural, I., 82. Ushatinskaya, R. S., 543.

Vaclav, V., 54, 369. Vago, C., 606. Valcarce, A. C., 514. Vallo, V., 305. Van, T. K., 78. van Bruggen, A. C., 531. van den Bosch, R., 90, 151. Van den Brande, J., 607. van den Bruel, W. E., 374, 607, 683. VanDenburgh, R. S., 509, 511, 630. van der Laan, P. A., 94. van der Merwe, J. H., 694. Vanderplank, F. L., 28. Vanderveken, J., 610. Vanderzant, E. S., 404, 517. van de Vrie, M., 444. Van de Woestijne, N., 607. Van Dijck, W., 374. van Dinther, J. B. M., 93. van Emden, H. F., 319, 320, 495. van Eyndhoven, G. L., 444. Van Middelem, C. H., 64. van oud Loosdrecht, W. E. de Mol, 42. Van Steyvoort, L., 249. van't Sant, L. E., 299, 608, 673. Vanwetswinkel, G., 302. Vappula, N. A., 103. Varis, A. L., 103, 674. Varma, B. K., 495. Varma, P. M., 80. Vasakas, D. T., 237, 238. Vasantharaj, B., 79. Vasantharaj David, B., 424. Vasev, A., 540. Vashkov, V. I., 541. Vasić, K., 54, 55, 367, 368. Vasil'ev (Vassilev), V. P., 609. Vasiljević, L., 55, 369. Vasilyan, V. V., 113. Vasquez, L. E. R. Lajo, 357. Vassilev, V. P. (see Vasil'ev, V.P.). Vasudeva, R. S., 80. Vasudeva Menon, P. P., 424. Vedeneeva, N. A., 495. Velez, R. Muñiz, 356. Venables, D. G., 197. Ventura, E., 669. Venugopal, S., 316.

Verhoyen, M., 609. Vernon, J. D. R., 432. Veselá, O., 247. Vessia, R., 48. Vidano, C., 660. Vijzelman, H. E., 100, 673. Vilkova, N. A., 361. Vincent, L. E., 619. Vinogradova, N. M., 446. Vité, J. P., 339, 340. Voegelé, J., 355, 382, 653, 654, 655. Vogel, J., 9. Vogel, R., 383. Vogel, W., 664. Vojdani, S., 520. Vojnović, V., 370. Völk, J., 303. von Bernegg, A. Sprecher, 693. von Tubeuf, C., 251. Vorontsov, A. I., 354. Voroshilov, N. V., 62. Voss, G., 443. Vrie, M. van de, **444**. Vries, J. E. de, 91.

Waddle, B. A., 220. Wade, W. H., 482. Wadhi, S. R., 170, 499. Wagner, R. E., 496. Wagner, W., 81. Wai (Hung-chuen), 593. Waites, R. E., 64. Walcher, H. Pschorn-, 184, 185. Waldbauer, G. P., 552. Wali, M., 176. Walkden, H. H., 554. Walker jr., J. K., 300, 576. Walker, P. T., 179. Walker, R. L., 322. Walker, T. J., 567. Wall, H. C., 21, 65. Waloff, N., 426. Walter, E. V., 327, 476. Walther, B., 91. Walton, R. R., 338. Wang, L. Y., 625. Wang (P'ing-yuan), 624. Wang (Yun-pei), 591. Wang, Y. S., 624. Ware, G. W., 323, 326. Warner, R. M., 265. Warren, L. O., 421. Warry, J. P., 40.

Watson, M. A., 358.

Watson, W. Y., 124. Watt, K. E. F., 90. Watters, F. L., 225. Watts, W. S., 164. Wave, H. E., 460, 640. Way, M. J., 550. Wearne, G. R., 496. Weaver, H., 299. Webb, D. v. V., 425. Webb, F. E., 121. Webley, D. P., 430. Wegorek, W., 366, 683. Weiser, J., 89, 105, 186, 283. Weismann, L'., 235, 305. Welch, H. E., 162, 207. Wells, A., 16. Wene, G. P., 638. Wessel, R. D., 573. West, A. S., 326. Westdal, P. H., 524, 551. Westigard, P. H., 402. Wheatley, G. A., 432. Wheatley, P. E., 179. Whellan, J. A., 81. Whitcomb, W. H., 21, 65. White, E. B., 356. White, J. H., 430. Whitfield, F. G. S., 496, 553. Whitlaw jr., J. T., 222. Whitney, W. K., 554. Wiąckowski, S., 189, 545. Wickramasinghe, Nallini, 484. Wiesmann, R., 100. Wilbert, H., 380, 683. Wilcox, J., 217. Wilcoxson, R. D., 398. Wilde, J. de, 6, 90, 204. Wilkes, L. H., 300, 417. Wilkinson, W., 485. Wille T., J. E., 559. Williams, C. B., 30. Williams, D. J., 234. Williams, G. C., 694. Williams, K., 388, 394, 525. Willis, E. R., 274, 565. Wilson, D. M., 620. Wilson, E. L., 221. Wilson, F., 589. Wilson, H. L., 563. Wilson, J. A., 575. Wilson, J. W., 327. Wilson, M. C., 572. Wilson, W. T., 582. Winfield, A. L., 318, 373. Winton, M. Y., 407, 634.

Witkowski, W., 61. Woehst, J. E., 561. Woestijne, N. Van de, 607. Wojnarowska, P., 365. Wolf, 489. Wolfenbarger, D., 328, 397, 413, 463. Wong Kong, 501. Woo (Wei-chün), 625. Wood, D. L., 300, 339, 340. Wood jr., E. A., 219, 470. Woodard, G. T., 384. Woodham, D. W., 479. Woodroffe, G. E., 657. Woodruff, R. E., 495. Woodville, H. C., 288. Wootten, N. W., 291. Wray, D. L., 517. Wressell, H. B., 163. Wright, D. W., 432. Wylie, H. G., 257, 378.

Yakovlev, B. P., 244. Yaman, I. K. Abu, 39. Yaruss, F. R., 271. Yates, H., 227. Yates III, H. O., 495. Yathom, S., 351. Yeh (Jen-chu), **501.** Yeomans, A. H., **226.** Yerington, A. P., **265.** York, G. T., **509.** Young, V. L., **593, 625.**

Zabirov, Sh. M., 542. Zacha, V., 444. Zaher, M. A., 149. Zaitzov, A., 613. Zaki, M., 277. Zangheri, S., 136, 661. Zawirska, I., 366. Zečević, D., 54, 55, 370, 371. Zelený, J., 678. Zhu (Zhao-qi), 501. Zhukovskiĭ, A. V., 13. Zhukovskii, S. G., 446. Ziady, S. A. E. El-, 30. Ziarkiewicz, T., 545. Zimmermann-Gris, S., 614. Živanović, V., 240. Živojinović, S., 54. Zocchi, R., 40, 136, 137, 311, 535. Zweig, G., 92, 384. Zwölfer, H., 168, 184, 185.

GENERAL INDEX

In the case of scientific names of arthropods, the page references are cited only under the generic headings. Chemical compounds having numerical prefixes (bis-, di-, etc.) are indexed under these prefixes, but the prefixes o-, m- and p- (ortho, meta and para), O-, N- and S- to indicate position of radicals, and normal, secondary and tertiary are ignored for alphabetisation.

A

Abacarus hystrix, transmitting mosaic virus of rye-grass in Europe, 71; and virus diseases of wheat and Agropyrum repens in Ontario, 71, 523.

Abbella subflava, parasitising leafhoppers in California, 630.

abbreviatus, Hypolithus,

Abelmoschus esculentus (see Hibiscus).

aberrans, Typhlodromus.

Abies, Lepidoptera on, in Canada, 208, 529; pests of, in U.S.S.R., 13, 112; Choristoneura fumiferana on, in U.S.A., 208; Argyresthia fundella causing die-

back of, in Yugoslavia, 535.

Abies alba, Lepidoptera on, in Czechoslovakia, 602, 603; Chermes nordmannianae on, in Denmark, 674; Choristoneura murinana on, in France, 168; pests of, in Germany, 168, 185, 186, 251, 375; Formica spp. in forests of, in Italy, 615; C. merkeri on, in Sweden, 185; C. piceae on, in Washington, 209; physiological effects of Chermes on, 45; injury to, by insecticidal dips, 674. Abies amabilis, Chermes piceae on, in

Abies balsamea, pests of, in Canada, 25, 26, 121, 185, 394, 455, 456, 522; pests of, in U.S.A., 121, 122, 149, 337, 411, 419; Chermes piceae on, in Holland, 185; silvicultural methods for control of C. piceae on, 457; annual growth of, as indicator for timing spray programmes, 411.

Abies fraseri, Chermes piceae on, in Holland, 185; C. piceae on, in N.

Carolina, 513.

U.S.A., 209.

Abies grandis, Chermes piceae on, in Oregon, 209.

Abies lasiocarpa, Chermes piceae on, in U.S.A., 209.

Abies nobilis, Chermes piceae on, in Holland, 185.

Abies pectinata (see A. alba).

abietella, Dioryctria.

abieticola, Gilpinia.

abietina, Pristiphora.

abietinum, Rhopalosiphum (Liosomaphis). abietis, Agevillea (Diplosis); Baeacis; Cephalcia; Chermes; Ernobius; Hylobius; Neodiprion.

abietivorella, Dioryctria.

ablutellus, Raphimetopus.

abnormis, Leptomastidea.

Abraxas miranda, bionomics of, on Euonymus in China, 503; measures against, 503.

abruptalis, Syngamia.

Absidia caerulea, infesting termites, 428.
Abutilon indicum, Earias spp. on, in India, 37.

abutilonea, Trialeurodes.

AC-5727 (see m-Isopropylphenyl Methylcarbamate).

Acacia, White (see *Robinia pseudacacia*).

Acantholyda erythrocephala (on pines), in Germany, 10; in U.S.S.R., 598; bionomics and parasites of, 10; insecticides against, 10.

acanthopus, Hoplopleura.

Acanthoscelides obtectus (infesting stored beans), in Angola, 99; in Colombia, 560; insecticides against, 560, 561, 603. Acarine Disease (of bees) (see Acarapis

woodi).

Acarapis (infesting honey bees), in Australia, Italy, New Zealand and U.S.A., 295, 342, 504, 505.

Acarapis dorsalis, 295; (external parasite of honey bees), in Canada and U.S.A., 342; species allied to, 505.

Acarapis externus (infesting bees), in Canada, 295, 342; in U.S.A., 342.

Acarapis woodi, 505; (infesting honey bees), question of occurrence of, in N. America, 342; in Britain, 617;

world distribution of, 295, 296, 507; pathogenicity of, 296; not considered as cause of Isle of Wight disease, 342; annual fluctuations of, 617; survey of data on, 507.

Acaricalus hederae, on Ilex opaca in British Columbia, 529.

acarorum, Gelis.

Acarus farris, in Britain, 549; hypopus of, misidentified as that of A. siro, 549.

Acarus siro (infesting stored products), in Britain, 91, 549; in Canada, 424; in Poland, 364; in U.S.A., 518; food preferences of, 518; ecology of, 549; predation of Cheyletus eruditus on, 91, 549; hypopus of A. farris mistaken for that of, 549; enzyme activity of homogenates of, on phosphorus compounds, 213.

Acer platanoides, Japananus hyalinus on,

in Ontario, 529.

Acer pseudoplatanus, Xyleborus germanus

infesting, in Germany, 11.

Aceratagallia, species of, on beet in California, 630; parasites of, 630. Aceria, on litchi in Pakistan, 88.

Aceria ficus, transmitting fig mosaic, 70. Aceria litchii, on litchi in Hawaii, 463.

Aceria merwei, on sugar-cane in Natal, 83.

Aceria neocynodonis, injuring Bermuda grass in Arizona, 396; acaricides against, 396.

Aceria phloeocoptes, bionomics of, on plum and other stone fruits in Yugoslavia, 241; tests of sprays against, 241.

Aceria sheldoni (on Citrus), in Aden, 80;

in S. Africa, 83.

Aceria tulipae, in Canada, 71, 156, 524; in U.S.A., 71; on Agropyrum repens, 71, 523; on wheat, 71, 156, 524; transmission of viruses by, 71, 523, 524; relation of, to silver top of grasses, 156. aceris, Phenacoccus.

Acetal, in bait for Drosophila, 327.

Acetic Acid, as constituent of Celcure

(q.v.), 485.

Acetone, use of, for removal of eggs of Pieris brassicae, 291; inactivation of insecticide-induced neuroactive sub-

stance by, 63.

Acetylcholine, in Periplaneta americana, 62; in Tetranychus telarius, 40; effects of insecticides on synthesis of, 62; effects of insecticide-induced neuroactive substance on activity of, 63.

Acetyl Glycine, effects of, in diet of Drosophila melanogaster, 638.

Achaetoneura, parasitising Oiketicus kirbyi in Costa Rica, 638; parasitising Caligo eurilochus in Ecuador, 274.

Acheta, scope of, 567.

Acheta assimilis (see Gryllus).

Acheta commodus, urea preventing diapause in eggs of, 296.

Acheta domesticus, relation of congeners to, in America, 567; toxicity of ethion to, 603.

Achrysocharella orientalis, hyperparasite of Promecotheca papuana in New Britain, 166.

Achrysocharella ritchiei, parasitising Leucoptera spp. in Tanganyika, 653; effects of sprays on populations of,

Achrysocharoides, parasitising Evagora starki in Canada, 586.

Achrysopophagus, species of, as hyperparasites of *Planococcus citri* in Israel, 614.

Achrysopophagus insolitus, sp.n., reared from Centrococcus insolitus in India,

Acidia (see Euleia).

Acorns (see Oak).

Acorus calamus, toxicity of extracts of, to insects, 171, 173.

acraea, Estigmene.

Acrobasis angusella, hickory on Ontario, 529.

Acrobasis feltella, on hickory in Ontario,

Acrocercops amurensis, sp.n., bionomics of, on Quercus mongolica in U.S.S.R., 242.

Acrolepia assectella, characteristics and world distribution of, 507.

Acroleucus coxalis, on pimento in Brazil, 356.

Acrylonitrile (as fumigant), against pests of stored products, 267, 499; factors affecting toxicity of, to Tribolium confusum, 274; determination of, in air, 496; and carbon tetrachloride, 499.

Actia, key to species of, 293.

Actia crassicornis, parasitising Tortricids in Czechoslovakia, 683.

Actia painei, sp.n., parasitising Agonoxena pyrogramma in New Britain and Solomon Islands, 293.

aculeatus, Haplothrips.

Aculus cornutus, on peach in S. Africa, 83; Typhlodromus caudiglans feeding on, 522.

Aculus lycopersici, on tomato in Africa, 83.

acuminata, Aelia.

acutangulus, Agrilus.

acutus, Agrilus; Chironomus; Scaphytopius.

Acyrthosiphon (see Macrosiphum).

Adalia bipunctata, predacious on Aphis spp. in Bulgaria, 59, 537; predacious on Psylla pyri in Poland, 365.

adalia, Laelia.

Adelencyrtus, key to females of species of, 490; new species of, 491.

Adelencyrtus aulacaspidis, parasitising Aulacaspis rosae in Argentina and California, 491.

Adelencyrtus femoralis, sp.n., parasitising Lepidosaphes sp. in India, 491; L. beckii not attacked by, 491.

Adelencyrtus tibialis, sp.n., parasitising Ceroplastes destructor in Nepal, 491.

Adelges (see Chermes).

Adelphocoris lineolatus (on lucerne), in Canada, 72; in Bulgaria, 60; sprays against, 72.

Adelphocoris rapidus, on lucerne in Canada, 72; sprays against, 72.

Aden Protectorate, pests and tristeza disease of Citrus in, 80; Diparopsis watersi on cotton in, 486; first record of Contarinia sorghicola in, 684; parasite of Phyllocnistis in, 80.

Adhesive Bands, for trapping insects, 336, 482.

Adhesive Traps, for aphids, 131, 320; for bark-beetles, 459; use of, in determination of emergence date of *Ephestia elutella*, 374; for *Frankliniella tritici*, 212; for *Hypera variabilis*, 576; use of, for collection of insects at low levels, 29, 30; evaluation of adhesives for, 424.

Adhesives, influence of, on effectiveness of seed treatments, 674; toxicity of, in seed treatments of beet, 670; with

Bacillus thuringiensis, 107.

Adoretus sinicus, on litchi in Hawaii, 463; characteristics and world distribution of, 507.

Adscita pruni, on oak in Ukraine, 620.

advena, Ahasverus.

Aedes aegypti, survival of, in nitrogen atmospheres, 416; use of, in detection

of BHC residues, 324.

Aegeria exitiosa (on peach), in Ontario, 154; in U.S.A., 226, 326, 631, 641, 642; insecticides against, 155, 226, 326, 631, 641, 642; Bacillus thuringiensis against, 642.

Aegeria pictipes (on peach), in Ontario, 154; in U.S.A., 335, 336, 642; sprays against, 155, 335, 336; soil treatment with phorate ineffective against, 642.

Aegeria tipuliformis, bionomics of, on black current in Czechoslovakia, 445.

Aegeria vespiformis, bionomics of, damaging vines in Bulgaria, 539; development of larvae of, in apples, 539; on oak, 539.

Aeglopsis chevalieri, use of, in tests of transmission of tristeza disease of Citrus by aphids, 80.

aegypti, Aedes.

aegyptiaca, Icerya.

aegyptium, Anacridium.

Aelia, egg-parasites of species of, in Morocco and Middle East, 654, 655; FAO report on control of, on cereals, 300.

Aelia acuminata, in Morocco, 355, 382, 653, 654; in U.S.S.R., 363; on cereals, 363, 382, 653, 654; feeding on Cynodon dactylon, 653; diapause in, 653; parasites of, 654, 655; method of rearing, 654.

Aelia cognata, in Morocco, 355, 382, 653, 654; on cereals, 382, 653, 654; diapause in, 653; new parasite reared from eggs of, 654.

Aelia germari, in Morocco, 355, 382, 653, 654; on cereals, 382, 653, 654; diapause in, 653; new parasite reared from eggs of, 654.

Aelia rostrata, on wheat in Middle East, 485; natural enemies of, 485.

Aelia sibirica, on cereals and grasses in U.S.S.R., 363.

Aeneolamia varia saccharina, on sugarcane in Trinidad, 332; number of nymphal instars of, 248; sampling method for, 332.

aeneus, Meligethes.

aequatus, Rhynchites (Coenorhinus).

aereus, Monodontomerus. aeripennis, Ctenicera.

Aerosols, acaricides applied in, against mites in houses or greenhouses, 419, 517; insecticides applied in: against adults of *Melolontha*, 682; against Trypetids, 676, 680; other outdoor applications of insecticides in, 10, 27, 55, 85, 247, 348, 536, 537, 539, 550, 600, 623; applied from aircraft, 55, 680; tests of, for use against insects in aircraft, 418.

aestivum, Apion (see A. trifolii).

Aethecerus pinifolii, parasitising Evagora milleri in California, 222.

aethiops, Microctonus.

affaber, Alcidodes.

affinis, Agria; Dasyneura; Drosophila; Lounsburyia; Pempherulus; Perga.

Africa, survey of data on Spodoptera exempta in, 530; natural enemies of insects in tropical areas of, 653.

Africa, French West (see Ivory Coast and

Guinea).

Africa, South, insects reared from pomegranates in, 531; first record of Contarinia sorghicola in, 684; vine pests in, 83, 298, 650; Nudaurelia cytherea capensis on pines in, 425; Lasioderma serricorne in stored tobacco in. 650; Dermestes infesting hides and skins in, 49; Pheidole megacephala punctulata in, 87; question of Cryptolestes capensis in, 656; Eriophyids of economic importance in, 83; natural enemies of insects in, 491, 653.

Africa, West, termites in, 485, 495. afrum, Zagrammosoma variegatum.

Agar, in diets for insects, 155, 339, 342, 404; use of, in rearing insects, 281; use of, in media for transporting and rearing mites, 640, 641.

agilis, parasitising Agathis

nubilalis in Delaware, 509.

Ageniaspis, parasitising Leucoptera spp. in Tanganyika, 652; effects of sprays

on populations of, 653.

Ageniaspis fuscicollis (parasitising forms of Hyponomeuta padellus), in Bulgaria, 111; in Finland, 195; bionomics of, 195.

Agevillea abietis, on silver fir in Germany, 251, 375; parasites of, 252, 375; biological control of, by *Platygaster* manto, 375; Dasyneura balsamicola considered distinct from, 252; synonymy of, 251.

agilis, Agathis; Anystis; Phorocera.

Agonoderus lecontei, on maize in Illinois, 420; insecticides against, 420.

Agonoxena, new Chalcidoid reared from species of, in Queensland, 52.

Agonoxena argaula, on coconut in Fiji, 52; Elachertus agonoxenae introduced into Fiji against, 52.

Agonoxena pyrogramma, on coconut in New Britain, 293; in New Guinea, 52; in Solomon Islands, 293; new parasites of, 52, 293.

agonoxenae, Elachertus,

agonus, Limonius.

Agria affinis, attacking pupae of Stilpnotia salicis in British Columbia, 126; effects of diet on parasitism of, by Aphaereta pallipes, 389.

Agria mamillata, parasitising Hyponomeuta padellus malinellus in Finland,

195.

Agrilus acutangulus, bionomics of, on basket willows in Yugoslavia, 138; destroyed by sap flow, 138; sprays against, 138.

Agrilus acutus, factors affecting incidence

of, on Hibiscus spp. in India, 594; larval habits of, 594.

Agrilus sinuatus, bionomics of, on pear in Sicily, 299.

Agriophara coricopa, on Olearia in New Zealand, 75; tests of Bacillus thurin-

giensis against, 75.

Agriotes, population fluctuations of, attacking maize in old grassland in France, 371; question of occurrence of European species of, in Nova Scotia, 73; effects of cultural practices on populations of, 371; soil treatments against, 371, 372.

Agriotes fuscicollis, on wheat in China,

625.

Agriotes litigiosus tauricus, bionomics of, on maize and other crops in U.S.S.R., 14; characters of stages of, 14.

Agriotes mancus, on potato and grasses in Nova Scotia, 73; destroyed by bac-

terial toxins, 582.

Agriotes sputator, in grassland in Nova Scotia, 73; bionomics of, on maize and other crops in U.S.S.R., 14; characters of stages of, 14.

Agromyza obtusa (see Melanagromyza). Agromyza oryzae, characteristics and

world distribution of, 507.

Agromyza phaseoli (see Melanagromyza). agromyzae, Euderus.

agropyrella, Sipha (see Rungsia kurdjumovi).

Agropyrum, aphids on, in New Brunswick, 585, 586; Ochsenheimeria vacculella on, in U.S.S.R., 596.

Agropyrum cristatum, relation of mites and thrips to silver top of, in Alberta, 156.

Agropyrum repens, pests of, in Canada, 71, 162, 523, 585; Meromyza saltatrix on, in Japan, 96, 97; Haplodiplosis equestris on, in Sweden, 101; oviposition of Hadena sordida on, in U.S.S.R., 447; mites in relation to viruses of, 71, 523.

Agrotis munda (see Euxoa radians).

Agrotis orthogonia, biochemical study of varieties of wheat in relation to feeding by, 527; rôle of digestive enzymes in larval nutrition of, 69.

Agrotis segetum, experiments with Thelohania weiseri and, 105; characteristics and world distribution of, 506.

Agrotis ypsilon, on potato, tomato and other vegetables in Brazil, 556, 557; bionomics of, on vegetables and field crops in Bulgaria, 346; on maize in Illinois, 420; in Ontario, 210; effects of food-plants on mortality of, 346; INDEX 721

parasites of, 346; insecticides against, 210, 347, 420, 557.

Agrypon flaveolatum (parasitising Operophtera brumata), in Europe, 258, 378; introduced into Canada, 378; in U.S.S.R., 190; effects of temperature on survival of, 258.

aguacatae, Copturus.

Ahasverus advena, dietary requirements of, in relation to infestation of stored products, 657.

ainsliella, Bucculatrix.

Aircraft, dispersal of *Popillia japonica* by, 418; tests of aerosols for use in, 418; container for shipment of insects in, 25; used for applying insecticides, 7, 55, 121, 122, 211, 220, 222, 276, 280, 383, 395, 411, 413, 425, 455, 462, 530, 540, 541, 569, 574, 659, 680; application of virus suspensions from, 336, 531; effects on general insect fauna of insecticides applied from, 55; survey of forest infestations by, 456; collection of arthropods by, 578.

Akar 338 (see Chlorobenzilate).

Alabama, Thysanoptera on groundnuts in, 151; Heliothis zea on maize in, 227; Scapsipedus micado in, 567.

Alabama argillacea (in Texas), on cotton, 16, 413; development of, on Cienfuegosia sulphurea, 518; insecticides against, 16, 413.

Alanine, in saliva of Aphis pomi, 95.

l-Alanine, in relation to feeding response of wireworms, **528**.

alaskensis, Microplitis; Pachynematus (Pikonema).

Alaxon, 678. alba, Polyphylla.

albatorius, Barichneumon.

Alberta, pests of beet in, 402, 468; Macrosiphum pisum on lucerne in, 210; Chorizagrotis auxiliaris on mustard in, 640; Cephus cinctus on wheat in, 72, 161, 527; Aceria tulipae transmitting wheat mosaic virus in, 524; mites and thrips associated with silver top of grasses in, 156; forest pests in, 208; Acarapis sp. infesting honey bees in, 295; beneficial insects in, 161, 469; Coleoptera established in, for control of Linaria vulgaris, 386.

albicans, Monochaeta (Cyzenis).

albicostalis, Assara.

albifrons, Spilochalcis. albimanus, Anopheles.

albinella, Rupela.

Albizia julibrissin, damaged by Homadaula albizziae in Kentucky, 273.

albizziae, Homadaula. alboannulatus, Dirhicnus. albocingulata, Staurochaeta.
albofasciatus, Spanogonicus (Leuco-

poecila).

Albumen, in baits for *Gryllotalpa*, **442**. *Alcidodes affaber*, bionomics of, on cotton and *Hibiscus* in India, **314**.

Alder (see Alnus). aldrichi, Sarcophaga.

Aldrin, against ants, 120, 420, 421, 589; against aphids, 420, 421, 468, 498; against other Hemiptera, 213, 316, 557; against Curculionids, 16, 109, 114, 152, 255, 275, 280, 329, 360, 417, 421, 500, 509, 673; toxicity of, to Hypera variabilis, 317; against Elaterids, 371, 372, 420, 421, 433; against other Coleoptera, 171, 173, 316, 323, 324, 366, 396, 417, 420, 421, 471, 493, 226, 288, 330, 484, 608, 623; against Lepidoptera, 16, 211, 277, 297, 309, 383, 420, 421, 534, 553, 557, 661; toxicity of, to locusts and grasshoppers, 172, 498, 521; effects of humidity on susceptibility of Schistocerca gregaria to, 170; excretion of, by S. gregaria, 321, 322; against termites, 77, 119, 293, 553, 577, 627; against Thysanoptera, 71; toxicity of, to Schizotetranychus andropogoni, 628; against cotton pests, 352; Scutigerella immaculata favoured by, 433; in dusts, 71, 109, 114, 120, 170, 288, 293, 309, 316, 323, 366, 417, 534, 553, 555, 557, 568, 627, 661, 673; humidity in relation to fumigant action of, 471; in granules, 152, 323, 325, 330, 396, 417, 421, 568; in sprays, 16, 71, 101, 170, 173, 175, 213, 330, 360, 383, 484, 509, 557, 572, 608; as deposit, 316, 471; seed treatment with, 50, 51, 65, 66, 101, 226, 329, 421, 433, 484, 568, 589, 608, 627, 674; soil treatment with, 40, 50. 51, 71, 101, 152, 171, 226, 254, 255, 275, 280, 288, 293, 309, 324, 325, 330, 352, 366, 371, 372, 388, 396, 417, 421, 433, 468, 493, 526, 534, 553, 555, 568, 570, 661; persistence and metabolism of, in soils, 16, 77, 119, 146, 255, 272, 416, 417, 527, 572, 573; treatment of planting material with, 77, 119, 288, 297, 557; pre-flooding treatment of ricefields with, 329; treatment of planting water with, 388; treatment of selfing bags with, 277; treatment of nests with,

627; uses of, against pests of stored products, 500; systemic activity of, in peas, 146; toxicity of, to plants, 65, 101, 288, 297, 360, 557, 675; wheat stimulated by, 627; effects of, on beneficial insects, 40, 469, 627; restriction in use of, in relation to hazards to wild life, 433; residues of, on plants, 330, 417; residues of, in milk or tissues of domestic animals, 384, 521; question of residues of, 570; analysis of residues of, 91; metabolism of, 63, 64, 146, 321; dieldrin as metabolite of, 272, 322, 384, 417; bioassay of, 146, 397; labelled with 36Cl, 321; and endrin, 509; and fungicides, 50; antagonistic action of sesamex with, 63, 64.

Aleochara, bionomics of, attacking Hylemya brassicae in U.S.S.R., 362.

Aleochara bilineata, parasitising Hylemya brassicae in Holland, 39.

Aleurites fordii, review of pests and diseases of, 693.

Aleurites montana, review of pests and diseases of, 693.

Aleurocanthus spiniferus, characteristics and world distribution of, 507.

Aleurocanthus woglumi, on Citrus in Aden, 80; sprays against, 80.

Aleurolobus barodensis, on sugar-cane in India, 497, 498; effects of, on sugar content of cane, 498; sprays against, 498.

alexanderi, Apanteles.

Alfalfa (see Lucerne).

Alfalfa Aphid, Spotted (see Therioaphis maculata).

Algeria, Bruchids on flax in, 249.

algiricus, Bruchidius.

algonquin, Drosophila.

Alhagi camelorum, as water source for Eurygaster integriceps, 429.

alhambra, Myzocallis.

Alkali Bee (see Nomia melanderi).

Alkaloids, in experiments on resistance of Solanum to Leptinotarsa decemlineata, 318.

alkani, Idiocerus.

Allatectomy, pseudo-diapause induced in Leptinotarsa decemlineata by, 7.

Allethrin, experiments with Alphitobius laevigatus and, 201, 687; resistance to, in strains of Sitophilus granarius, 82, 128; question of metabolism of, 64; and piperonyl butoxide, 82, 128, 687; and other synergists, 687.

Alloglycaspis (see Glycaspis).

Allolobophora caliginosa, insecticide residues in, 641.

Allotropa burrelli, destroying Pseudococcus comstocki, 164; introduced into Ontario from Virginia, 164.

Allotropa convexifrons, destroying Pseudococcus comstocki, 164; introduced into Ontario from Virginia, 164.

Allvl Isothiocvanate (Mustard glucosides associated with, in relation to oviposition by Pieris brassicae, 290; tests of fumigant action of, on Sitophilus granarius, 9.

Almond, pests of, in California, 405, 482;

pests of, in Italy, 38, 181.

Alnus, Cryptorhynchus lapathi on, in Hungary, 307.

Alphitobius diaperinus, intercepted

oil-seed cake in Germany, 11.

Alphitobius laevigatus, infesting dry fruits of tamarind in India, 499; factors affecting susceptibility of, to insecticides, 4, 201, 687.

Alsike Clover (see Trifolium hybridum). Alsophila pometaria, sprays against, on

apple in Nova Scotia, 159.

alternans, Itoplectis.

Althaea rosea, Earias spp. on, in India,

Aluminium Phosphide, fumigation of stored products with, against pests, 483, 558; tablets containing, 483, 558.

Aluminium Silicate, as carrier for insecticide dusts, 50.

alveofrons, Dasiops. amalthea, Trigona.

Amarantus, Hymenia recurvalis on species of, in India, 498.

Amarantus gangeticus, 498.

Amarantus retroflexus, infected by beetyellows virus, 17.

Amarantus viridis, Hymenia recurvalis

on, in Fiji, 88.

Ambeodontus tristis, effects of boron compounds on larvae of, in timber, 648.

ambiguella, Clysiana (Clysia).

ambiguus, Psallus.

Amblymerus (see Mesopolobus).

Amblypelta cocophaga, 235.

Amblypelta lutescens papuensis, on coconut, rubber, cassava and Ipomoea spp. in Papua, 235; ants associated with, 235; parasites of, 235; dieldrin spray against, 235.

Ambrosia, Micrutalis malleifera transmitting curly-top virus to, in Florida,

Ambrostoma quadriimpressum, bionomics and natural enemies of, on elms in China, 593; measures against, 593.

America, nomenclature of crickets in, 567.

723 INDEX

Am. Cyanamid 12008 (see O. O-Diethyl S-Isopropylthiomethyl Phosphorodithioate).

Am. Cyanamid 12009 (see O, O-Diethyl S-Propylthiomethyl Phosphorodithi-

Am. Cyanamid 12503 (see O. O-Diethyl O-(2-Phenyl-6-(3(2H)-pyridazinonyl)) Phosphorothioate).

Am. Cyanamid 18133 (see O, O-Diethyl O-2-Pyrazinyl Phosphorothioate).

Am. Cyanamid 18706 (see O, O-Dimethyl S-Ethylcarbamoylmethyl Phosphorodithioate).

Am. Cyanamid 24055 (see 1, 1-Dimethyl-3-(p-acetamidophenyl)triazene).

Am. Cyanamid Anti-feeding Compound 24055 (see 4'-(Dimethyltriazeno) Acetanilide).

americana, Leucopis; Periplaneta; Platy-

americanum, Malacosoma; Zagrammo-

amerinda, Eurygaster,

Ameroseius, in stored grain in Canada,

Ametastegia glabrata, on apple Finland, 103.

amethystinus, Tetrastichus (Geniocerus). dl-y-Aminobutyric Acid, relation of, to feeding response of wireworms, 528.

3-Amino-1, 2, 4-triazole (see Amitrole). Amiton Oxalate, against *Planococcus citri*, 612; mode of action of sesamex as synergist with, 63.

Amiton Toluene-p-sulphonate, toxicity of, to Tetranychus telarius, 478.

Amitrole, effects on Macrosiphum pisum of broad beans treated with, 525; not toxic in sprays to M. pisum, 525.

Ammonium Carbamate, fumigation tablets of aluminium phosphide (q.v.)

and, 483, 558.

Ammonium Nitrate, effects on mite populations of treatment of apple trees with, 259.

Amnemus quadrituberculatus, characteristics and world distribution of, 507. amoena, Sciara (Neosciara).

Amorphoidea lata, characteristics and world distribution of, 507.

Amphimallon solstitiale, injuring pines in U.S.S.R., 111; characteristics and world distribution of, 507.

Amphorophora forbesi, on Rubus spectabilis in British Columbia, 529.

Amphorophora parviflori, on Rubus parviflorus in British Columbia, 529; as vector of virus diseases of Rubus, 529.

Amphorophora rubi, on strawberry in British Columbia, 692; on Rubus spp. in U.S.A., 132; transmitting virus diseases of plants, 132, 692.

Amsinckia, Eurygaster spp. on, in California, 520.

amurensis, Acrocercops.

amygdali, Eurytoma; Hyalopterus.

Amylases, rôle of, in nutrition of Agrotis orthogonia, 69; in relation to transmission of viruses by Tettigonia viridissima, 443.

Anabrolepis extranea, in China, 491; introduced into California for control

of Aonidiella aurantii, 491.

Anabrolevis zetterstedtii, parasitising Quadraspidiotus perniciosus in Switzerland, 664.

Anacanthotermes ochraceus, attacking straw in bricks in Egypt, 180; measures

against, 181.

Anacentrinus deplanatus, bionomics of, in U.S.A., 219; on sugar-cane, sorghum and grasses, 219; insecticides against,

Anacridium aegyptium, on Citrus in Turkey, 82; parasitised by *Ceracia* mucronifera in France, Spain and Tunisia, 313.

Anaesthetisation, of insects, 356.

Anagasta kuehniella, study on multiparasitism of larvae of, 618, 619; use of eggs of, in study of parasitism by Trichogramma cacoeciae, 681; susceptibility of sexes and stages of, to pyrethrins, 3, 4; effects of y-radiation on stages of, 607.

Anagrus giraulti, parasitising leafhoppers

in California, 630.

Anagyrus kivuensis, parasitising Planococcus citri in Israel, 614; parasites of,

Ananas comosus (see Pineapple).

Anaphes (see Mymar).

Anarsia lineatella, on almond in California, 405; fluctuations in populations of, 405.

Anastatus, egg-parasite of Amblypelta lutescens papuensis in Papua, 235.

Anastatus disparis, egg-parasite Lymantria dispar in Yugoslavia, 54, 55, 367.

anastomosalis, Omphisa.

Anastrepha ludens, in California, 279; in Mexico, 279, 280, 325; on Citrus, 280; measures against, 280, 325.

Anastrepha mombinpraeoptans, on coffee in Brazil, 643; baits and traps for, 643.

anceps, Apamea.

Ancylis comptana (fragariae), on strawberry in Kentucky, 570; insecticides against, 570.

Ancylis mitterbacheriana, parasites reared from, in Czechoslovakia, 683.

ancylivorus, Macrocentrus. andersoni, Typhlodromus.

andropogoni, Schizotetranychus. Anemone, Hepialus lupulinus on, in Britain, 430.

angaleti, Apanteles.

Angitia (see Nythobia).

Angola, Coleoptera attacking timber trees in, 608; Coleoptera in stored beans in. 99.

Angophora intermedia, pests of, in New

South Wales, 232.

angulatus, Blepharidopterus.

angusella, Acrobasis.

angusticollis, Zootermopsis.

Anisandrus (see Xyleborus).

Anisota senatoria, on oak in Connecticut, 271, 407; bionomics of, 407; parasites and diseases of, 271, 407.

anitus, Asolcus (Microphanurus) (see A. rufiventris).

annulipes, Chelonus.

Anobium punctatum (infesting timber), in Britain, 489: natural enemies of. 489; surveys of data on, 489, 646, 648; artificial egg-laying sites for, 231; tests and uses of timber preservatives against, 646, 647, 648.

Anodontonyx noxius, damaging wheat and oats in New South Wales, 493;

soil treatments against, 493.

Anolis, destroying Oiketicus kirbyi, 638; destroying beneficial insects, 229; bird introduced into Bermuda against, 229. Anomala, on cereals and other crops in

China, 593.

Anomala corpulenta (in China), on cereals and other crops, 593; on crab apple and other trees, 501, 502; bionomics of, 501; DDT sprays against, 502.

Anomala horticola, characteristics and

world distribution of, 506.

Anomis editrix, insecticides against, on jute in British Guiana, 27.

Anomis texana, on cotton in Brazil, 559, 560; natural enemies of, 560; sprays against, 559.

Anopheles albimanus, used in tests of carbamate insecticides, 516.

Anopheles quadrimaculatus, used in tests of dichlorvos, 562.

Anoplolepis custodiens, established in Zanzibar, 29; predacious on Oecophylla

Anoplolepis longipes, in Papua, 235; established in Zanzibar, 29; predacious on Oecophylla sp., 29; occurrence of Amblypelta lutescens papuensis absence of, 235.

Antestiopsis, species of, on coffee in Uganda, 490; sampling methods for, 490.

Antestiopsis lineaticollis, sprays against, on coffee in Tanganyika, 653.

Antheraea pernyi, feeding on dried oak leaves in U.S.S.R., 192.

Anthocoris, predacious on Aphis pomi in Wisconsin, 415.

Anthocoris confusus, in Canada, 429; development and prey of, 429.

Anthocoris gallarumulmi, in Canada, 429;

development and prev of, 429.

Anthocoris melanocerus, bionomics of, predacious on Psylla pyricola in British Columbia, 158; effects of sprays on populations of, 158.

Anthocoris minki, in Canada, 429; devel-

opment and prey of, 430.

Anthocoris musculus, predacious on Psylla mali in Nova Scotia, 523; effects of sprays on populations of, 523.

Anthocoris nemoralis, in Canada, 429; development and prey of, 429.

Anthocoris nemorum, in Canada, 429; in Poland, 365; development and prey of, 429; predacious on Psylla pyri, 365. Anthocoris sarothamni, in Canada, 429;

Anthonomus eugenii, on Capsicum in California, 637; insecticides against,

development and prey of, 429.

Anthonomus grandis (on cotton), in Mexico, 407; in U.S.A., 16, 147, 264, 322, 330, 407, 408, 410, 412, 413, 464, 467, 473, 515, 517, 568, 573, 574, 576; selection of morphological varieties of cotton by, 464; effects of, on yields, 574; development of, on Cienfuegosia, 518; adult habits of, 410, 473, 576; factors affecting life-span in strains of. 407, 408; diets for, 404, 407; collection of eggs of, 225, 404; methods of labelling, with ³²P, 471; effects of ³²P on stages of, 472; effects of γ-radiation on stages of, 568; insecticides against, 16, 322, 330, 331, 338, 408, 412, 413, 467, 515; distribution of dimethoate in plants in relation to, 472; resistance to insecticides in strains of, 147, 218, 219; genetic basis of resistance to endrin in strains of, 220; biological differences between susceptible and endrin-resistant strains of, 264; metabolism of insecticides in, 218, 219, 476.

Anthonomus pomorum, characteristics and

world distribution of, 507.

Anthonomus rubi, on strawberry, raspberry and other plants in U.S.S.R., 360; bionomics of, 360; insecticides against, 360.

725 INDEX

Anthonomus vestitus, characteristics and world distribution of, 507.

Anthophila pariana, on apple in Austria, 302.

Anthoxanthum odoratum, stem-boring Diptera on, in England, 487.

Anthracene Oil (see Tar Distillates).

Anthrenocerus australis, treatments

flannel against, 587.

Anthrenus flavipes, basis of attraction of, to textile stains, 615; not attracted by Kepone bait, 339; insects infesting cultures of, 339.

Anthrenus verbasci, in Britain, 249; damaging textiles in Germany, 377; effects of photoperiod on pupation of, 249.

Anthribus

nebulosus, parasitising lecanium corni in France, 683.

Antigastra catalaunalis, bionomics and parasite of, on sesame in India, 316.

antiqua, Hylemya; Orgyia. antirrhini, Gymnetron.

antonii, Helopeltis.

Antonina graminis, in India and Texas, 406; on grasses, 406; parasite of, 406. Antrocephalus, species of, parasitising Corcyra cephalonica in India, 314; sex of, in relation to that of host, 314;

diet for, 314.

Ants, in Brazil, 120; relation of, to fungus disease of aphids in Congo, 610; prevention of establishment of pastures by, in Queensland, 589; in U.S.A., 420, 517; on maize, 420; nesting on railway embankment, 517; associated aphids, 144, 419; associated with Coccids, 229, 519, 531; effects of, on natural enemies of Coccids, relation of, to Pseudotheraptus wayi on coconut, 28, 29; occurrence of Amblypelta lutescens papuensis in absence of, 235; destroying other insects, 24, 104, 138, 141, 157, 166, 167, 251, 352, 363, 393, 596, 615, 616, 638, 682; woodpeckers predacious on, 104, 141; measures against, 120, 145, 420, 464, 589, 639; transplantation of colonies of, 615, 616; classification of, 104.

Anuraphis bakeri, transmitting mosaic viruses of bean in Washington, 264.

Anuraphis helichrysi, transmitting virus disease of beet, lettuce and radish in California, 131.

Anuraphis maidiradicis, on maize in Illinois, 420; insecticides against, 420. Anuraphis plantaginea (on apple), bionomics of, in Holland, 673; in Nova

Scotia, 388; natural enemies of, 388, 673; synonymy of, 388.

Anuraphis roseus (see A. plantaginea).

Anuraphis tulipae, in Canada, 529; on tulip, carrot and iris, 529; transmitting

tulip-breaking virus, 529.

Anystis agilis, predacious on Phytodecta olivaceus in Britain, 30; predacious on Psylla mali in Nova Scotia, 523; effects of sprays on populations of, 523.

Aonidiella, parasite of species of, on

Citrus in Burma, 491.

Aonidiella aurantii, in Aden, 80; in Burma, 266, 491; in India, 626; in Israel, 22, 97; in Mexico, 22; in Morocco, 382; in Siam, 491; in U.S.A., 22, 221, 266, 491, 561, 688; on Citrus, 22, 80, 97, 221, 382, 491, 561, 688; on mulberry, 626; natural enemies of, 21, 22, 266, 267, 491; large-scale rearing of natural enemies of Coccids on, 229: survey of data on biological control of, 688; other measures against, 221, 383, 561; congeners misidentified as, 688.

Aonidiella citrina, on Citrus in Aden, 80; misidentified as A. aurantii, 688.

Aonidiella orientalis, in China, India and Pakistan, 491; on Carissa grandiflora, Euonymus and papaya, 491; parasites of, 491.

Aonidiella taxus, A. aurantii misidentified

as, 688.

aonidum, Chrysomphalus (see C. ficus).

Apamea anceps, bionomics of, on cereals in U.S.S.R., 362, 363; natural enemies and virus disease of, 363; measures

against, 363.

Apanteles, parasitising Evagora starki in Canada, 586; parasitising Ceramidia viridis in Ecuador, 353; parasitising Choristoneura spp. in Europe and N. America, 169; parasitising Callimorpha quadripunctaria, 87; experiments with Galleria mellonella and larvae of, 87. Apanteles alexanderi, parasitising larvae

of Eulia loxonephes in Argentina, 144. Apanteles angaleti, parasitising Pectinophora gossypiella in India, 317; introduced into U.S.A., 317; reared on Corcyra cephalonica, 317; factors affecting reproductive potential of, 317.

Apanteles bordagei, parasitising Leucop-

tera in Tanganyika, 652.

Apanteles californicus, parasitising Eva-gora milleri in California, 124, 221; parasitising E. starki in Canada, 586; parasites of, 124.

Apanteles fumiferanae, parasitising Choristoneura fumiferana in U.S.A., 24, 337; vertical distribution of, 24.

Apanteles glomeratus, in Germany, 380, 683; in Japan, 78; in New York, 401; in U.S.S.R., 114, 115; parasitising Aporia crataegi, 380, 683; parasitising Pieris spp., 78, 87, 114, 115, 401, 678; experiments with Galleria mellonella and larvae of, 87; range of parasitism by, of other Lepidoptera, 115; effects of, on development of Apanteles pieridis, 381; factors affecting induction of diapause in, 678.

Apanteles jucundus, parasitising larvae of Operophtera brumata in Europe, 378; introduced into Canada, 378.

Apanteles lacteicolor, parasitising Cydia funebrana in U.S.S.R., 112.

Apanteles liparidis, parasitising Dendrolimus superans in Mongolia, 116.

Apanteles longicauda, parasitising Eucosma nigricana in Czechoslovakia, 602; effects of sprays on populations of, 603.

melanoscelus Apanteles (parasitising Lymantria dispar), in Massachusetts, 400; in Yugoslavia, 54; persistence of, in sprayed areas, 400.

Apanteles murinanae, parasitising Eucosma nigricana in Czechoslovakia, 602; survey of data on, parasitising Choristoneura murinana in France and Germany, 168.

Apanteles opsiphanis, parasitising larvae of Caligo eurilochus in Ecuador,

274.

Apanteles ordinarius, parasitising larvae of Dendrolimus superans in Mongolia, 116.

Apanteles ornigis, parasitising stages of Bucculatrix ainsliella in Michigan, 327. Apanteles parasitellae, parasitising

Spilonota prognathana in U.S.S.R., 112. Apanteles pastranai, sp.n., parasitising Rhyaciona buoliana in Argentina, 144.

Apanteles pieridis, parasitising Aporia crataegi in Germany, 380, 683; parasitising Pieris brassicae, 381; bionomics of, 380; effects of Apanteles glomeratus on development of, 381; parasite of, 381.

Apanteles sarcitorius, parasitising Cydia funebrana in U.S.S.R., 112.

Apanteles sericeus, parasitising Operophtera brumata in U.S.S.R., 190.

Apanteles sesamiae, parasitic on Busseola

fusca in Nigeria, 292.

Apanteles solitarius, established in British Columbia, 126; parasitising Lymantria dispar and Apatele psi in Czechoslovakia, 683; larvae of L. dispar parasitised by, in Yugoslavia, 54; parasitising Stilpnotia salicis, 126.

Apanteles starki, parasitising Evagora starki in Canada, 586.

Apanteles xanthostigma, parasitising Lepidoptera in U.S.S.R., 112.

Apate monachus rufiventris, on almond, lemon, orange and peach in Sardinia, 181; question of other food-plants of, 181; mining habits of, 181; measures

against, 181. Apatele psi, parasitised by Apanteles solitarius in Czechoslovakia, 683.

Apechthis resinator, parasitising Tortricids in Czechoslovakia, 683; survey of data on, parasitising Choristoneura murinana in France and Germany, 168.

Apechthis rufata (parasitising Tortricids), in Czechoslovakia, 683; survey of data on, in France and Germany, 168; in

U.S.S.R., 598.

Aphaereta pallipes, effects of host diet on parasitism of Agria affinis by, 389.

Aphelenchylus elongatus, bionomics of, parasitising Ips confusus, 496.

Aphelia viburniana, bionomics of, on conifers in Europe, 375; parasites and fungus disease of, 375, 376.

Aphelinoidea plutella, parasitising leafhoppers in California, 630.

Aphelinus, parasitising Aphis fabae in Bulgaria, 537.

Aphelinus howardii, parasitising Macro-siphum pisum in California, 66.

Aphelinus mali (parasitising Eriosoma lanigerum), in China, 591; in Czechoslovakia, 678; in France, 312, 313; in U.S.S.R., 592; bionomics and factors affecting populations of, 591, 592; toxicity of insecticides to, 312, 313, 678.

Aphelinus perpallidus, parasitising Chromaphis juglandicola in California, 151. Aphid, Black-margined (see Monellia

costalis).

Aphid, Cowpea (see Aphis craccivora). Aphid, Spotted Alfalfa (see Therioaphis maculata).

Aphidius, species of, parasitising Anuraphis plantaginea in Holland, 673.

Aphidius avenae (picipes), parasitising Anuraphis plantaginea in Holland, 673. Aphidius pisivorus, parasitising Macrosiphum pisum in California, 66.

Aphidius rapae (see Diaeretus).

Aphidius smithi, parasitising Macrosiphum pisum, 66; established in California from India, 66; characters distinguishing, from other parasites of aphids on lucerne, 66.

Aphidoletes thompsoni, predacious on Chermes piceae, 25; container for

shipment of, in aircraft, 25.

Aphids, sooty mould favoured by, 415; and virus diseases of plants, 16, 17, 23, 34, 35, 71, 80, 81, 95, 97, 101, 106, 107 INDEX 727

547, 549, 605, 610, 611, 614, 636, 669, 680, 684, 688, 689, 690, 691, 692; injection of plant viruses into, 133; routes of virus transport in, 358; classification of transmission of viruses by, 691; inhibition of non-persistent viruses by body fluids of, 250; preference of, for excised or virus-infected leaves, 584; ants associated with, 144, 419; natural enemies of, 151, 240, 243, 279, 281, 319, 387, 388, 415, 429, 456, 459, 469, 605, 673, 683; fungus diseases of, 459, 462, 570; mechanism, habits and physiology of feeding of, 44, 45, 92, 105, 423, 482, 493; composition of saliva of, 45; methods for collecting saliva from, 95; life-cycles of, in relation to biochemistry of food-plants, 423; rôle of wavelength in food-plant selection by, 200, 201; varietal susceptibility of plants to, 167, 219, 405, 406, 470, 551, 575; plant-growth regulators in relation to, 525, 551, 629; relation of populations of, to planting patterns in crucifers, 68, 69; factors affecting populations of, 320, 377; influence of electromagnetic fields on moulting of, 444; factors affecting dispersal and flight of, 305, 320, 481, 482, 596, 605; physiology and duration of flight in, 203, 204; respiration and water losses of, 322, 443; sampling methods for, 402, 463, 551, 620; traps for, 131, 160, 320, 482, 639; high-altitude catches of, 199; forecasting of outbreaks of, ecology of, on apple, 673; nature of injury to lucerne by, 322; identity of species of, on strawberry in N. America, 23; in Finland, 101; on strawberry in Germany, 106; on fruit trees in Poland, 188; in U.S.A., 17, 19, 131, 160, 278, 339; rearing methods for, 496; labelled with radioactive isotopes, 93; tests of systemic insecticides against, 249; resistance to phosphorus compounds in strains of, 284; classification and new species of, 144, 262, 358, 494, 584, 585. (See also Honeydew.)

Aphis, species of, on Citrus in Aden, 80.

Aphis citricidus (see Toxoptera).

Aphis craccivora, in Aden, 80; bionomics of, on lucerne, beans, clover and other plants in Bulgaria, 59, 110; in Congo, 610; in Gaza Strip, 97; on cowpeas in India, 498; in Senegal, 298; in U.S.A., 262; on broad bean in Victoria, 34; on Citrus, 80, 97; on groundnuts, 298,

610; on vines, 110; as vector of plant viruses, 34, 35, 298, 610; natural enemies of, 59; relation of ants to fungus infesting, 610; insecticides against, 59, 498; misidentified as A. medicaginis, 262.

Aphis cuscutae, transmitting virus yellows

of beet, 17.

Aphis fabae, in Britain, 199, 201, 320, 549, 550; in Bulgaria, 59, 536; in California, 419; in Canada, 429; in Czechoslovakia, 305; in France, 605, 669, 670; in Germany, 377, 545, 546, 547; in U.S.S.R., 191; on beet, 59, 191, 377, 536, 545, 546, 549, 605, 669, 670; on broad beans, 536, 550, 605; on Euonymus europaeus, 305; on potato, 320, 546, 547; on Tropaeolum, 603; on vines, 303; primary food-plants of, 536, 537; transmitting virus diseases of plants, 17, 303, 545, 546, 547, 549, 605, 669; stomach size of, in relation to food-plants, 423; bionomics of, 537; respiration of, 443; flight habits of, 199, 320, 537, 547, 605; physiology and duration of flight in, 203, 204; effects of flight on length of life and fecundity of, 204; colour preferences of, 201; traps for, 320, 547; ant associated with, 419; natural enemies of, 192, 429, 537; factors affecting populations of, 192, 305, 320, 377; measures against, 59, 537, 549, 550, 603, 605, 670.

Aphis forbesi (see Cerosipha).

Aphis frangulae, auct. (see A. gossypii). Aphis gossypii, in Aden, 80; in Brazil, 357, 559, 643, 645; in Gaza Strip, 97; in Greece, 352; in Israel, 615; in Poland, 544; in U.S.A., 16, 131, 263, 278, 279, 338, 480; transmitting virus yellows of beet, 17; on Citrus, 80, 97; as vector of tristeza disease, 80, 97; on cotton, 16, 263, 352, 467, 473, 479, 559, 643, 645; on Hibiscus esculentus, 357; on Easter lily, 278; on melon, 279; and virus diseases of potato, 544; transmitting virus disease of spinach, 480; question of transmission of sweet-potato virus by, 131; transmitting virus disease of lettuce to zinnia, 615; on greenhouse plants, 338; parasite of, 279; insecticides against, 263, 278, 279, 352, 357, 473, 479, 559, 643, 645; use of, in tests of residues of phorate, 395.

Aphis helichrysi (see Anuraphis).

Aphis idaei, in British Columbia, 692;
transmission of strawberry viruses by,
692; misidentified as A. rubifolii, 692.

Aphis maidis, in Japan, 452; in Kenya,
179; in New Brunswick, 585; in U.S.A.,

160, 639; in Yugoslavia, 80; bionomics

of, on barley, 452, 585; as vector of barley yellow-dwarf virus, 160; on grasses, 160, 585; on maize, 179; on sorghum, 80, 179; question of transmission of sorghum red-stripe disease by, 80; trap for, 639.

Aphis medicaginis, auct. (see A. craccivora). Aphis medicaginis Koch, on leguminous plants in Europe, 262; A. craccivora

misidentified as, 262.

Aphis nasturtii, distribution of Rhamnus cathartica in relation to, in Germany, 255; in New Brunswick, 633; in Poland, 544; on potato, 544, 633; transmitting potato viruses, 255, 544; insecticides against, 633; honeydew of, in diets for Ephialtes buolianae, 391.

Aphis pisi (see Macrosiphum pisum).

Aphis pomi (on apple), in Canada, 388, 394, 395, 455, 524; in France, 604; in Holland, 673; in Italy, 616; in Poland, 188, 677; in U.S.S.R., 359; in U.S.A., 402, 415; development of sooty mould favoured by, 415; contamination of apples by honeydew of, 402; methods for collecting saliva from, 95; amino acids in saliva and honeydew of, 95; bionomics of, 673; effects of climatic factors on mortality of, 455; fluctua-tions in populations of, 415; natural enemies of, 388, 415; question of migration of parasites of, 673; insecticides against, 188, 388, 389, 395, 402, 415, 524, 604, 616, 677; effects of sprays against, on populations of Tetranychus mcdanieli, 394; sampling methods for, 402, 551.

Aphis rubifolii, A. idaei misidentified as,

692.

Aphis rumicis, auct. (see A. fabae).

Aphis sacchari, bionomics of, on sorghum in China, 624; overwintering on Miscanthus, 624; measures against, 624. Aphis spiraecola, on orange, 516; toxicity

of malathion and methylcarbamates to,

516.

Aphodius granarius, on Mentha spp. in Washington, 324; soil treatments

against larvae of, 324.

Aphodius tasmaniae, infesting pastures in S. Australia, 165, 166, 504; effects of grassland management on populations of, 504; effects of moisture on immature stages of, 165, 166; fungus infesting, 166.

Aphomia gularis, infesting dry fruits of tamarind in India, 499.

Aphrastasia (see Chermes).

Aphrodes bicinctus, in Czechoslovakia, 304; in Germany, 441; in Hungary, 307; in Ontario, 686; on clover, 304,

307, 686; on lucerne, 307; on potato, 307; on strawberry, 441, 686; as vector of plant viruses, 304, 307, 441, 686.

Aphthona euphorbiae, bionomics of, on flax in Rumania, 608; insecticides

against, 608.

Aphycus brasiliensis, sp.n., parasitising Chaetococcus bambusae in Brazil, 491.

Aphycus californicus, parasitising Eulecanium pruinosum, 491; in California, 491; A. insidiosus indistinguishable from, 491.

Aphycus helvolus, parasitising Coccids in California, 519; influence of ants on

populations of, 519.

Aphycus insidiosus, parasitising Eulecanium corni in France, 683; imported into California, 491; A. californicus indistinguishable from, 491.

Aphycus lichtensiae, parasitising Saissetia oleae in Pakistan, 491; introduced into

California, 491.

Aphycus luteolus (parasitising Coccids), in California, 519; introduced into U.S.S.R., 597; influence of ants on populations of, 519.

Aphycus punctipes, parasitising Eulecan-

ium coryli in France, 683.

Aphycus stanleyi, parasitising Coccids, 491; introduced into California, 491; question of identity of, 491.

Aphytis coheni, sp.n., parasitising Aonidiella aurantii in Israel, 22; introduced into California, 22; misidentified as A. lingnanensis, 21, 22.

Aphytis diaspidis (parasitising Quadraspidiotus perniciosus), introduced into Pakistan, 170; in Switzerland, 664.

Aphytis holoxanthus, sp.n., parasitic on Chrysomphalus ficus, 22; in Hong Kong, 22; introduced into California from Israel, 22; misidentified as A. lingnanensis, 21, 22.

Aphytis lepidosaphes, parasitising Lepidosaphes beckii, 564; established in

U.S.A., 564.

Aphytis lingnanensis, parasitising Aonidiella aurantii, 22; in China, 21; established in Mexico and U.S.A., 22; not occurring in Israel, 22; new species misidentified as, 21, 22.

Aphytis maculicornis, parasitising Parlatoria oleae, 227; in Iraq and Persia, 227; introduced into California, 227; effects of sprays on populations of, 227.

Aphytis mytilaspidis, parasitising Quadraspidiotus perniciosus in Switzerland, 664.

Aphytis proclia, parasitising Quadraspidiotus perniciosus in France, 380;

competition of, with Prospaltella perniciosi, 380.

Aphytis spiniferus, parasitising Chrysomphalus in Brazil, 491.

apiella, Depressaria.

Apion, species of, in Bulgaria, 540; in U.S.S.R., 114, 355; in Yugoslavia, 32; on clover, 114, 540; effects of infestation by, on seed yields of red clover, 32; on trees and shrubs, 355; bionomics and natural enemies of, 540; measures against, 540.

Apion aestivum (see A. trifolii).

Apion apricans (on clover), in Bulgaria, 540; in Czechoslovakia, 445; bionomics of, in Rumania, 56; parasites of, 56; DDT against, 56.

Apion assimile, on red clover in Czecho-

slovakia, 445.

Apion dichroum, on red clover in Czechoslovakia, 445.

Apion flavipes (see A. dichroum).

Apion seniculum (on clover), in Czechoslovakia, 445, 446; in U.S.S.R., 114; bionomics of, 114; dusts against, 114. Apion trifolii (on clover), in Bulgaria, 540; in Czechoslovakia, 445, 446; bionomics of in Rumania 56; paragites

bionomics of, in Rumania, 56; parasites of, 56; DDT against, 56.

Apion varipes, on red clover in Czechoslovakia, 445.

Apion varium, infestation of cowpea by, in Nigeria, 49, 50; measures against, 50.

Apion virens (on clover), in Czechoslovakia, 445, 446; in U.S.S.R., 114; bionomics of, 114; dusts against, 114. Apleurotropis lalori, parasitising Promecotheca papuana in New Britain, 166;

misidentified as Derostenus sp., 166. Apomecyna binubila, characteristics and

world distribution of, 507.

Aporia crataegi, in Germany, 380, 683; in Poland, 364; in U.S.S.R., 115; on fruit and other trees, 364, 380; bionomics of, 364; parasites of, 115, 380, 683;

virus disease of, 683.

Apple, Eriophyes pyri on, in S. Africa, 83; Eulia loxonephes on, in Argentina, 144; Cydia pomonella on, in Australia, 165; pests of, in Austria, 302; pests of, in Britain, 379, 670, 671, 672; pests of, in Bulgaria, 59, 60, 110, 111; pests of, in Canada, 26, 159, 163, 259, 262, 273, 388, 394, 395, 454, 455, 523, 524, 529; pests of, in China, 591, 592, 625; Eriosoma lanigerum on, in Czechoslovakia, 678; pests of, in Finland, 103, 194; pests of, in France, 8, 84, 312, 313, 379, 434, 435, 550, 604, 665, 666, 669; pests of, in Germany, 9, 107, 319, 381,

439, 440, 442, 681, 682; pests of, in Greece, 239; pests of, in Holland, 444, 673; Tetranychids on, in Hungary, 307; pests of, in Italy, 136, 309, 533, 534, 616; pests of, in Japan, 35, 36; Quadraspidiotus perniciosus on, in New Zealand, 167; Q. perniciosus on, in Pakistan, 170; pests of, in Poland, 188. 189, 676, 677, 678; pests of, in Switzerland, 46, 86, 100, 663, 664; pests of, in Turkey, 177, 237; pests of, in U.S.S.R., 15, 112, 242, 343, 359, 360, 361, 543, 597; pests of, in U.S.A., 18, 20, 23, 266, 268, 400, 402, 403, 415, 417, 466, 470, 481, 694; sampling method for Aphis pomi on, 551; A. pomi in relation to sooty mould on, 415; physiology of necrosis caused by Q. perniciosus in, 186; Typhlodromus pyri feeding on pollen of, 319; spatial distribution of parasite of C. pomonella on, 681; factors affecting arthropod fauna of, 159, 259, 262, 376, 410, 439, 682, 694; hail damage to, 159; assessment of damage to fruit of, 159, 160; forecasting infestation of, 84: systemic activity of insecticides in. 188, 395; toxicity of insecticides to, 394, 466, 669; insecticide residues on, 525; (fruit), development of larvae Aegeria vespiformis in, 539; speed of entry of larvae of C. pomonella into, 637; E. lanigerum in, 481, 482; larvae of pests destroyed by controlled-atmosphere storage of, 403; contamination of, by honeydew, 402; factors affecting adherence and toxicity of spray deposits on, 21; use of, in rearing of wireworms, 575; in diets for insects, 216, 418.

Apple, Crab (see Pyrus prunifolia).

Apple Maggot (see *Rhagoletis pomonella*). Apple Scab (see *Venturia inaequalis*).

Apple Sucker (see Psylla mali).

apricans, Apion.

Apricot, pests of, in California, 128, 271; Cydia molesta on, in Ontario, 163; Ceratitis capitata on, in Spain, 98; Lepidoptera on, in Switzerland, 46, 100; Perotis malachitica on, in Turkey, 495; pests of, in U.S.S.R., 112, 242.

Aprostocetus, new species of, in England and Sweden, 493; scope of, 493.

Apsylla cistellata, bionomics of, on mango in India, 315.

apterus, Lethrus.

Aptesis, parasitising Pristiphora erichsonii in Michigan, 210.

Aptesis basizona, 210; (parasitising sawflies), introduced into Ontario from Europe, 155, 209, 387; bionomics of 387; factors affecting reproduction of, 155, 209.

Aptesis galactina, parasitising Pristiphora abietina in Germany, 682; effects of BHC applied against Melolontha on populations of, 682.

aquilonia, Formica.

arachidis, Cicadulina.

Arachis hypogaea (see Groundnuts).

Aradus cinnamomeus, on pines in U.S.S.R., 116.

Araecerus fasciculatus, in stored coffee beans in Brazil, 558; phosphine fumigation against, 558.

Araecerus suturalis, infesting dry fruits of

tamarind in India, 499.

Aramite (see 2-Chloroethyl 2-(p-tert.-Butylphenoxy)-1-methylethyl Sulphite).

Arbor-vitae (see Thuja).

arborea, Bryobia (see B. rubrioculus).

Arborol, containing DNC and tar distillate (q.v.), 600, 678.

Archips argyrospilus, on apple in Quebec, 26, 159; bionomics of, 26; sprays against, 26.

Archips cerasivoranus, in Canada, 391; Ephialtes buolianae reared on, 391.

Archips crataeganus, bionomics and natural enemies of, on fruit trees in U.S.S.R., 15; sprays against, 15.

Archips rosanus, in Czechoslovakia, 683; on fruit trees in U.S.S.R., 360; parasites of, 683; DDT sprays against, 360.

Archips xylosteanus, parasites of, in Czechoslovakia, 683.

arenacearia, Tephrina.

arenaria, Vespula.

argaula, Agonoxena. argentifera, Plusia.

Argentina, pests of cotton in, 694; pests of fruit trees in, 144, 299, 558; Gryllus (Acheta) assimilis on green forage crops in, 356; Ptericoptus sp. on potato in, 552; Rhyacionia buoliana on pine in, 144, 230; new Tortricid on Solanum bonariense in, 230; new Pyralid on sunflower in, 230; Aulacaspis rosae in, 491; Acarapis woodi infesting honey bees in, 296; beneficial insects in, 144, 230, 356, 491, 552.

argentinensis, Tersilochus (Porizon).

argentosus, Poliaspis. argillacea, Alabama.

Argon Ionisation Detector, use of, in gas chromatography of insecticides, 577. Argon-glow Lamps, in light-traps for

Lepidoptera, 476, 477.

Argyresthia conjugella, in Germany, 9, 10, 441, 442; in U.S.S.R., 112; on apple, 9, 10, 112, 442; on Sorbus aucuparia,

9, 10, 441; bionomics of, 9, 10; parasites of, 10.

Argyresthia fundella, bionomics of, causing die-back of fir in Yugoslavia, 535, 536; tests and uses of insecticides against, 536.

argyrolepia, Chilotraea.

Argyroploce leucotreta (see Cryptophlebia).

Argyroploce variegana, bionomics of, on fruit trees in U.S.S.R., 360.

argyrospilus, Archips.

Argyrotaenia pulchellana, bionomics of, on apple, apricot and vine in Switzerland, 100.

Argyrotaenia velutinana, photoperiod in relation to diapause in, 634; continuous rearing of, 633.

aridula, Chaetocnema.

Arizona, pests of cotton in, 219, 478, 480, 638, 639; Lepidoptera on lettuce in, 514; Myzus persicae and Circulifer tenellus transmitting virus diseases of melon in, 408; pests of potato in, 473, 474; new Lonchaeid on walnut, sunflower and Juglans major in, 128; pests of Pinus ponderosa in, 460; Therioaphis maculata on lucerne in, 462; Aceria neocynodonis on Bermuda grass in, 396; grasshoppers on rangelands in, 224; notes on miscellaneous insects of, 221; beneficial insects in, 460, 638, 639; fungus disease of T. maculata in, 462; virus disease of Trichoplusia ni in, 639.

Arkansas, Zeadiatraea grandiosella on maize in, 21, 65; pests of rice in, 421; new sawfly on Pinus echinata in, 462; Thyridopteryx ephemeraeformis on ornamental evergreens in, 421.

armatus, Myochrous. armicollis, Magdalis.

armigera, Dicladispa; Heliothis.

Armillaria matsutake (see Tricholoma).

armillata, Nythobia (Angitia).

Arnica, Tyria jacobaeae feeding on, 260. Aroclor 5460, stabilisation of heptachlor with, 145.

Arrhenatherum elatius, of little effect on populations of Apion in mixture with clover, 445.

Arsenic Trioxide, against termite, 77.

Arsenious Oxide, as fumigant against termite, 181; and sulphur, 181.

Artemisia vulgaris, Ostrinia nubilalis on, in U.S.S.R., 14; experiments with strains of O. nubilalis and, 14.

Artichoke, Globe (see *Cynara scolymus*). Arum Lily (see *Zantedeschia aethiopica*). arvensis, Cephalcia.

ascalonicus, Myzus (Rhopalomyzus).

Ascorbic Acid, in diets for insects, 5, 404, 517.

Ash, Operophtera brumata on, in Belgium, 85; Hylesinus fraxini on, in Czechoslovakia, 187.

Ash, Mountain (see Sorbus aucuparia).

Asolcus, species of, parasitising eggs of Pentatomids in Middle East, 485, 654; in Morocco, 654; key to, 654.

Asolcus anitus (see A. rufiventris).

Asolcus basalis (egg-parasite of Pentatomids), in Morocco, 654, 655; in West Indies, 589; reproductive behaviour of, 589; A. perrisi probably identical with, 655.

Asolcus choaspes (see A. vassilievi).

Asolcus ghorfii, sp.n., egg-parasite of Pentatomids in Morocco, 654, 655; misidentified as A. scutellaris, 654.

Asolcus grandis, egg-parasite of Pentatomids in Morocco, 654, 655; misidentified as A. semistriatus, 655.

Asolcus perrisi, probably identical with

A. basalis, 655.

Asolcus rufiventris (egg-parasite of Pentatomids), in Morocco, 654, 655; in Turkey, 485; synonymy of, 655.

Asolcus scutellaris, distribution of, 654; new species misidentified as, 654.

Asolcus semistriatus (egg-parasite of Pentatomids), in Morocco, 654, 655; in Persia, 485; in Turkey, 485; factors affecting development of, 601; A. grandis misidentified as, 655.

Asolcus simoni reticulatus, subsp.n., eggparasite of Eurygaster integriceps in

Middle East, 655.

Asolcus tumidus, hibernation of, in Persia, 654.

Asolcus vassilievi, egg-parasite of Eurygaster integriceps in Persia, 485; synonymy of, 654.

Asparagin, in honeydew of Aphis pomi, 95; relation of, to oviposition by Tetranychus telarius, 547; differences in content of, in wheat varieties, 527.

1-Asparagine, in relation to feeding response of wireworms, 528.

Aspartic Acid, in saliva of *Aphis pomi*, **95**; relation of, to oviposition by *Tetranychus telarius*, **547**.

l-Aspartic Acid, in relation to feeding response of wireworms, 528.

Aspen, Trembling (see Populus tremuloides).

Aspergillus, relation of Sitotroga cerealella to infestation of stored grain by species of, 415; Tyrophagus putrescentiae reared on, 583.

Aspergillus amstelodami, 415.

Aspergillus flavus, experiments with Eurygaster integriceps and, 115.

Aspergillus glaucus, repens and ruber, 415. aspidioti, Habrolepis.

Aspidiotiphagus citrinus, parasitising Quadraspidiotus perniciosus in Switzerland. 664.

Aspidiotiphagus lounsburyi, parasitic on Lepidosaphes beckii in Morocco, 382.

Aspidiotus destructor, distribution of, in Pacific area, 81; on coconut, 81.

Aspidiotus hederae, large-scale rearing of natural enemies of Coccids on, 229.

Aspidiotus juglansregiae (see Quadraspidiotus).

Aspidiotus perniciosus (see Quadraspidio-

tus).

Aspirators, for collection and handling of insects, 372, 694.

Assara albicostalis, infesting dry fruits of tamarind in India, 499.

assectella, Acrolepia.

assimile, Apion.

assimilis, Ceutorhynchus; Gryllus (Acheta). assuanensis, Psammotermes.

assulta, Heliothis.

assumthi, Odontotermes.

Aster, China (see Callistephus chinensis). atalantae, Theronia.

ater, Cryptolestes; Dermestes.

Athalia rosae (colibri), bionomics of, on rape in Hungary, 308; characteristics and world distribution of, 507.

Atherigona indica, on sorghum in Kenya, 179; recovery of sorghum from attack

by, 179, dusts against, 179.

Athous subfuscus, cocoons of Pristiphora abietina destroyed by, in Germany, 141.

atlanticus, Tetranychus.

Atomaria linearis, on beet in Germany, 376, 377; infected by Cephalosporium, 377; effects of cultural measures on populations of, 377.

Atoposoma (see Zagrammosoma).

atra, Phyllotreta.

Atractotomus mali, destroying stages of Hyponomeuta padellus malinellus in Finland, 195; predacious on apple pests in Nova Scotia, 159; effects of sprays on populations of, 159; misidentified as Criocoris saliens, 159.

atramentaria, Campoplex (Omorga).

atriclavus, Tetrastichus.

atricornis, Phytomyza.

Atriplex, development of Agrotis ypsilon on, in Bulgaria, 346.

Atriplex semibaccata, infected by yellows virus of beet, 17.

atriplicella, Scrobipalpa.

atrolineatus, Bruchidius.

atropicta, Tegea.

Atropine, effect of, in insects, 62.

atropurpureus, Eupelmus.

atrum, Colaspidema.

Atta laevigata, dusts against, in Brazil, 120.

Atta sexdens, characteristics and world distribution of, 506.

Atta sexdens rubropilosa, dusts against, in Brazil, 120.

Attaclay, as carrier for insecticides, 277. Attacus cynthia ricini, Trichogramma evanescens reared in, in China, 623.

Attagenus megatoma (piceus), basis of attraction of, to textile stains, 635; enzyme activity of homogenates of, on phosphorus compounds, 213.

Attagenus pellio, damaging textiles in

Germany, 377.

Attapulgite, granules of, as carrier for heptachlor, 396.

attenuatus, Psylliodes.

aulacaspidis, Adelencyrtus.

Aulacaspis rosae, parasitised by Adelencyrtus aulacaspidis in Argentina and California, 491.

Aulacophora, characteristics and world distribution of species of, 507.

Aulacophora foveicollis (on cucurbits), in India, 173; bionomics of, in Israel, 613; characteristics and world distribution of, 507; toxicity of insecticides to, 173.

Aulacophora hilaris, characteristics and world distribution of, 507.

Aulacorthum (see Macrosiphum).

aulicus, Mesoleius.

Aulocara elliotti, on rangelands in Arizona, 224.

aurantii, Áonidiella; Scirtothrips; Toxoptera.

aurata, Crepidodera (Chalcoides).

auratus, Carabus.

auricularia, Forficula.

Aurigena (see Perotis).

auropunctatum, Calosoma maderae.

Australia, Cydia pomonella on apple in, 165; aphids on broad beans in, 34, 35; pests of cereals and grasses in, 684; Eumargarodes laingi on sugar-cane in, 265; forest pests in, 32, 33, 231, 232, 504, 552; Acarapis sp. introduced into, on honey bees from U.S.A., 504; Trogoderma spp. infesting stored products in, 18; Actia spp. in, 293; Epiphyas spp. in, 232; Eriococcus orariensis introduced into New Zealand from, 587; parasites of Listroderes introduced into, 353; Tyria jacobaeae imported into, for control of Senecio jacobaea,

452; other beneficial insects in, 32; virus diseases of plants in, 34, 35, 70, 684. (See also under the individual States.)

Australia, South, pasture pests in, 165, 166, 504; lucerne pests in, 233, 234; bees and *Eucalyptus* in pollination of lucerne in, 234; *Acarapis* sp. on honey bees in, 505; fungus infesting *Aphodius tasmaniae* in, 166.

australiaca, Nomia.

australis, Anthrenocerus; Leptoglossus.

Austria, pests of crucifers in, 184; pests of orchard fruits in, 85, 247, 302, 312; pests of vines in, 85, 248; forest pests in, 100; beneficial arthropods in, 100, 302.

austriaca, Eurygaster.

Austrotortrix (see Epiphyas).

Autographa biloba, parasitised by Hyposoter exiguae, 66.

Autographa californica (see Plusia).

Autographa gamma, characteristics and world distribution of, 507.

Autographa nigrisigna, on cabbage and rape in Japan, 79; effects of temperature on development and pupal coloration of, 79.

Autographa precationis, on mint in

Indiana, 284.

auxiliaris, Chorizagrotis.

Auxins, in honeydew of aphids, 629; relation of content of, to resistance of plants to aphids, 551.

avellanae, Phytoptus (Eriophyes). Avena Coleoptile Test, 629.

avenae, Aphidius; Macrosiphum.
Avocado, pests of, in California, 200, 378; Copturus aguacatae on, in Mexico, 355; pests and diseases of, in Morocco, 383; biological control of pests of, 200; sorption and retention of ethylene dibromide by fruits of, 574.

Azalea, Paratetranychus ilicis on, in Japan, 452; P. ilicis on, in New York, 19; Tetranychus multisetis not feeding on, 67; systemic activity of insecti-

cides in, 19.

Azinphos-ethyl, in sprays against Lepi-

doptera, 466, 663.

Azinphos-methyl, against aphids, 16, 388, 468, 643, 645, 667; against Coccids, 120, 168; against other Hemiptera, 159, 395, 467, 515, 590; against Curculionids, 16, 322, 413, 436, 467, 515; against other Coleoptera, 211; against Hylemya brassicae, 510; tests and uses of, against Rhagoletis pomonella, 268, 273; against Tortricids, 9, 21, 239, 337, 398, 411, 434, 436, 466, 637; against other

Lepidoptera, 21, 161, 211, 326, 335, 336, 409, 469, 515, 559, 569, 590, 644, 663; Heliothis zea favoured by, 323; against Thysanoptera, 467, 477, 569; against Sminthurus viridis, 34; against Tetranychids, 16, 643, 645; cross-resistance to, in strain of Tetranychus pacificus, 512; effects of, on beneficial arthropods, 159, 412, 627; in dusts, 161, 409, 569, 644; in granules, 469; in sprays, 9, 16, 34, 120, 159, 168, 211, 213, 239, 268, 322, 326, 335, 336, 337, 388, 395, 396, 398, 411, 413, 434, 436, 466, 468, 477, 515, 525, 526, 559, 569, 590, 637, 643, 645, 663; as deposit, 273; lack of persistence of deposits of, 389; applied in drenches, 510; persistence of, in soil, 567; tests of, against pests of stored products, 213; effects of, on setting of potato tubers, 397; ovicidal action of, 326; toxicity of, to chicks, 475; question of residues of, in poultry foods, 475; residues of, on plants, 88, 398, 525, 526; residues of, in tobacco and cigarette smoke, 228; effects of weathering on residual toxicity of, 16; colorimetric determination of residues of, 88; degradation products of, 228; and DDT, 161, 409, 413, 467, 515; toxicity of DDT increased by, 21; and lead arsenate, 268; and malathion, 211; and Sevin, 268.

B

Bacillus, ingestion by forest insects of substances inhibiting growth of, 580. Bacillus cereus, pathogenicity of, to sawfly larvae, 282.

Bacillus fribourgensis, infecting Melolontha spp. in Czechoslovakia, 187.

Bacillus larvae, infecting honey bees, 332; toxicity of extracts and fractions of, to insects, 332.

Bacillus thuringiensis, infecting Lepidoptera, 75, 107, 184, 269, 292, 421, 465, 469, 514, 515, 525, 536, 642; ineffective against Lepidoptera, 9, 328; use of larvae of Trichoplusia ni in determination of activity of, 342; crystals not involved in killing of sawfly larvae by, 283; ineffective against aphids, 75; ineffective against Phyllotreta, 184; bees and other beneficial arthropods unharmed by, 75, 159, 582; laboratory experiments with, 184, 536; field use of, 9, 75, 107, 184, 421, 465, 469, 514, 515, 525, 536, 642; rain increasing effectiveness of, 107; commercial preparations of spores

of, 184, 269, 342, 421, 514, 525, 536, 582; tobacco injured by preparation of, 465; survey of data on, 140; and insecticides, 514, 525.

Bacillus thuringiensis sotto, pathogenicity

of, to sawfly larvae, 283.

Bacillus thuringiensis thuringiensis, infecting Lepidoptera, 569, 580; factors affecting susceptibility of Pectinophora gossypiella to, 427; infecting sawflies, 283, 341; not pathogenic to birds or mice, 282; survival of, in faeces of vertebrates, 282; field use of, 569, 580; commercial preparations of spores of, 580; formulation in relation to effectiveness of, 569.

Bacteria, occurrence of, in lamb pelts, 49; effect of insecticides on nitrifying, 553; ingestion by forest insects of substances inhibiting growth of, 580; (entomogenous), in Lepidoptera, 368, 389, 390, 531; infecting Melolontha, 186, 187; wireworms destroyed by toxins of, 582; ecology of use of, compared with chemical control, 262; surveys of data 88, 196. (See also Bacillus, Bacterium, Chromobacterium, Clostridium. Pseudomonas. Root Nodules. Serratia and Streptococcus.)

Bacterium juglandis, relation of Paramyelois transitella to, on walnut in

California, 276.

Bacto-agar, in diets for Diatraea saccharalis, 214.

Baeacis abietis, parasitic on Ernobius abietis in Italy, 136.

Baikiaea insignis minor, Platypodid on, in Uganda, 496.

baileyi, Glycaspis (Alloglycaspis).

Bait-sprays, against Trypetids, 98, 99, 279, 280, 308, 533; applied from aircraft, 280.

Bait-traps, use of, in sampling for Cydia

molesta, 665.

Baits, for ants, 464, 639; for Vespula germanica, 422; for Coleoptera, 319, 339, 577; sex attractant in, for Lyman*tria dispar*, **55**, **368**, **369**; for other Lepidoptera, **465**, **635**, **642**; for Trypetids, **383**, **384**, **439**, **533**, **590**, **643**; for other Diptera, 265, 326, 388; for cockroaches, 339; for Gryllotalpa gryllotalpa, 442; for insects damaging textiles, 635.

bajulus, Hylotrupes.

Meteorus: Typhlobakeri, Anuraphis; dromus.

Balaninus (see Curculio).

Balaustium, predacious species of, on soy bean in Delaware, 414. Balsam Fir (see Abies balsamea).

balsamicola, Dasyneura.

balteata, Diabrotica.

balteatus, Pamphilius; Syrphus.

Banana, Oiketicus kirbyi on, in Costa Rica, 637; pests of, in Ecuador, 268, 274, 353; Nacoleia octasema on, in Fiji, New Guinea and Indonesia, 353, 354; pests of, in Honduras, 462, 565, 566; pests of, in Queensland, 503; insects in relation to bacterial wilt of, 520; secretion from Scaptocoris divergens inhibiting soil fungi injurious to, 565; Chaetanaphothrips orchidii causing red rust of fruit of, 462; Cercospora leaf-spot disease of, 463; insecticides damaging fruit of, 503.

bandra, Chilotraea.

Bands (see Adhesive Bands).

banksi, Eutetranychus.

banksianae, Neodiprion pratti.

Barathra brassicae (see Mamestra).

Barbara colfaxiana, on Douglas fir in California, 355.

barberi, Dendroctonus.

barbita, Magdalis.

Barichneumon albatorius, parasitic on Pseudaletia unipuncta in Pakistan, 175. Baris, on cruciferous oil-seed crops in France, 373; natural enemies of, 373. Baris coerulescens, bionomics of, on rape

in Hungary, 308.
Barium Hydroxide, susceptibility of Galleria mellonella to virus disease

unaffected by, 427.

Barium Polysulphide, in sprays against *Leucoptera scitella*, **534**; and parathion, **534**.

Bark-beetles, associations of, 458; water supply of conifers in relation to infestation by, 253; factors influencing selection of logs by, 340; injurious fungi spread by, 124, 217; natural enemies of, 142, 283, 386, 387, 460; nematodes associated with, 619; X-ray detection of, in timber, 259; attractants and traps for, 44, 340, 341, 459, 460; measures against, 217, 218, 223, 448, 449, 460, 606, 641; methods of rearing, 390; temporary storage of larvae of, 281.

Barley, Macrosiphum avenae on, in Belgium, 375; pests of, in Bulgaria, 109, 537; pests of, in Canada, 70, 394, 529, 584, 585; Calligypona pellucida on, in Finland, 102, 103; Oscinella frit on, in Germany, 45; Meromyza saltatrix on, in Japan, 96, 97; Haplodiplosis equestris on, in Sweden, 101; Psylliodes elliptica on, in Turkey, 297; pests of, in U.S.S.R., 14, 243; pests of, in U.S.A., 95, 134, 214, 281, 401; Camarota

curvipennis on, in Yugoslavia, 249; survival of Chloriona orizicola on, 132; other insects on, 98, 113, 452, 629; Hemiptera in relation to virus diseases of, 95, 132, 133, 134, 375, 401; varietal susceptibility of, to insects, 22, 45, 629; seed treatment of, 401; toxicity of insecticides to, 401; detection of growth inhibitors in, 22; effects on vetch fauna of sprays against pests of, 211; protection of beet with shelter crop of, 606.

Barley (Stored), Coleoptera in, 154, 420, 554; other insects and mites in, 154; sorption of methyl bromide by, 554; effects of fumigants on germination

of, **579.**

Barley Groats, development of Sitophilus spp. on, 205, 206.

barodensis, Aleurolobus.

Barthrin (see 6-Chloropiperonyl Chrysanthemate).

basalis, Asolcus; Coccophagus; Orgyia.

basilinea, Manga. basizona, Aptesis.

Basudin (see Diazinon).

Bathyplectes curculionis, parasitising Hypera variabilis, 400; introduced into eastern U.S.A. from California, 400.

Bats, insects taken by species of, in

Germany, 684.

Bayer 19639 (see O, O-Diethyl S-2-(Ethylthio)ethyl Phosphorodithioate). Bayer 22408 (see O, O-Diethyl O-Naphthalimido Phosphorothioate).

Bayer 22684 (see 2-Chloroethyl 2.2-Dichlorovinyl Methyl Phosphate).

Bayer 23129 (see Thiometon).

Bayer 23453 (see O,O-Dimethyl S-2-(Ethylsulphinyl)ethyl Phosphorodithioate).

Bayer 24869 (see Ethyl 1-Chloro-2-propyl-2,2-dichlorovinyl Phosphate).

Bayer 25141 (see O,O-Diethyl O-p-Methylsulphinylphenyl Phosphorothioate).

Bayer 25198 (see O,O-Dimethyl O-p-Methylsulphinylphenyl Phosphorothioate).

Bayer 25820 (see Diethyl Naphthalimido Phosphate).

Bayer 29493 (see Fenthion).

Bayer 29709 (see O,O-Dipropyl O-p-Methylsulphinylphenyl Phosphorothioate).

Bayer 30911 (see O-Methyl O-2,4-Dichlorophenyl Methylphosphonothioate).

Bayer 37344 (see 4-Methylthio-3,5-xylyl Methylcarbamate).

Bayer 39007 (see o-Isopropoxyphenyl Methylcarbamate).

Baytex (see Fenthion).

Beans, Aphis craccivora on, in Bulgaria, 59; pests of, in Canada, 125, 454; Melanagromyza phaseoli on, in Ceylon, 484; pests of, in France, 604; Tetranychus telarius on, in Germany, 547; new Tetranychid on, in Japan, 452; Plusia chalcites on, in New Zealand, 75; instance pests and virus diseases of, in U.S.A., 16, 17, 264, 271, 279, 328, 397, 463; varietal susceptibility of, to insects, 328, 397, 414, 463; morphological study of injury to, by T. telarius, 107, 108; effects on T. telarius of treatment of, with gibberellin, 410; influence of, on resistance of T. telarius to acaricides, 548; use of T. telarius on, in tests of acaricides, 602, 603, 684; systemic activity of insecticides in, 16; effects of seed treatments on germination of, 484.

Beans (Stored), Coleoptera in, 99, 560; treatments of, 99, 560, 561.

Beans, Broad (Vicia faba), aphids on, in Australia, 34; Aphis fabae on, in Britain, 550; A. fabae on, in Bulgaria, aphids on, in France, 605; Taeniothrips lefroyi on, in India, 499; pests of, in U.S.A., 264, 413; Argyrovelutinana reared on, 634; Bruchidius spp. on, 249; resistance of, to Epilachna varivestis, 414; effects on Macrosiphum pisum of plant-growth regulators in, 525; M. pisum and varietal susceptibility of, 603, 629; oviposition of Pieris brassicae on, 290; aphids and virus diseases of, 34, 35, 264; transmission of clover virus to, by Aphrodes bicinctus, 304; effects of sprays applied to, on bees, 550. Beans, Field (see Beans, Broad).

Beans, Lima (see Lima Beans).

Beans, Mung (see Phaseolus aureus).

Beans, Wax, Tetranychus multisetis feeding on, 67.

Beauveria, infesting Micrutalis malleifera,

Beauveria bassiana, infesting Haplothrips tritici in Bulgaria, 348; infesting Oiketicus kirbyi in Costa Rica, 638; infesting Melolontha spp. in Czechoslovakia, 187; infesting Leptinotarsa decemlineata in U.S.S.R., 447; infesting insects in U.S.A., 210, 509; infesting other insects, 164, 284, 541; experiments with Eurygaster integriceps and, 115; penetration of, into larvae of L. decemlineata, 196; field use of, 164, 447, 541; and insecticides, 115, 447, 541.

Beauveria tenella, infesting Pristiphora erichsonii in Pennsylvania, 210. beckii, Lepidosaphes (Mytilococcus).

Beech, pests of, in Germany, 11, 376; transplantation of colonies of Formica lugubris to areas of, in Italy, 615.

Beer, as attractant for wasps, 422. Bees, in relation to honeydew require-

ments of Formica polyctena, 105. Bees, Alkali (see Nomia melanderi).

Bees, Honey, effects of Bacillus spp. on, 333, 582; mites infesting, 295, 296, 342, 504, 505, 617; relation of Acarapis woodi to Isle of Wight disease of, 296, 342; question of resistance to A. woodi in strains of, 617; other natural enemies of, 29, 38, 422, 551; effects of insecticides on, 234, 276, 349, 374, 394, 412, 423, 432, 515, 550, 640; enzyme activity of homogenates of, on phosphorus compounds, 213.

Bees, Wild, Acarapis spp. on, in U.S.A., 342; relation of *Eucalyptus* to, in pollination of lucerne, 234. (See also Megachile, Nomadopsis, Nomia and

Trigona.)

Beeswax, absorption of chlordane vapour

by, 640.

Beet, pests of, in Alberta, 402, 468; aphids on, in Belgium, 249; pests of, in Britain, 549; pests of, in Bulgaria, 59, 347, 348, 536, 541; Lygus spp. on, in Finland, 103; aphids on, in France, 605, 669; pests of, in Germany 253, 376, 377, 439, 440, 441, 442, 545, 546; Tetranychus telarius on, in Israel, 613; Cleonus mendicus on, in Italy, 533; pests of, in U.S.S.R., 15, 191, 609; pests of, in U.S.A., 17, 131, 330, 331, 335, 419, 463, 630; Chaetocnema tibialis on, in Yugoslavia, 181; aphids and virus diseases of, 17, 131, 249, 253, 331, 463, 482, 545, 546, 549, 605, 669; other vectors of yellows virus of, 17; effect of yellows virus on sugar content of, 17; development of resistance in, to vellows virus, 17; cultural practices in relation to infestation of, by aphids, 605, 606; feeding habits of Myzus persicae on leaves of, 482; relation of seedling age to injury by L. rugulipennis to, 103; effects of, on development of Agrotis yspilon, 346; T. multisetis feeding on, 67; oviposition of T. telarius on, 547; influence of, on telarius on, 547; influence of, on resistance of T. telarius to acaricides, 548; forecasting abundance of pests of, 15; systemic activity of insecticides in, 249; seed treatment of, 253, 605, 606,

609, 670; toxicity of adhesives and insecticides to, 330, 402, 605, 670; residues of heptachlor in, 402.

Beet, Sea-kale (see Chard).

Beet, Silver (see Chard).

Begonia semperflorens, Pegomyia bicolor on, in Italy, 661.

behmani, Mymar (Anaphes).

behrii, Etiella.

Belgium, aphids and virus yellows of beet in, 249; Macrosiphum avenae transmitting virus disease of cereals in, 375; Leptinotarsa decemlineata on potato in, 681, 683; Steneotarsonemus fragariae on strawberry in, 445; Operophtera brumata on orchard trees in, 85; forest pests in, 85, 606; Plutella maculipennis in, 430; Tetranychus telarius in, 688.

Bell-bird (see Manorina melanophrys).

bella, Sturmia.

bellinus, Typhlodromus.

Bemisia tabaci, on Citrus in Aden, 80; in Greece, 352; transmitting virus disease of sweet potato in Israel, 130; in Sudan, 94; on cotton, 94, 352; insecticides against, 94, 352; favoured by DDT, 94, 95.

Bentonite, as diluent for BHC dusts, 99;

killing Coleoptera, 99.

Benzedrine (see dl-α-Methylphenethylamine).

Benzoxazole, study of analogues of, inhibiting growth of *Ostrinia nubilalis* and *Penicillium*, **68**.

Ber (see Ziziphus jujuba).

berlesei, Caloglyphus; Prospaltella.

berlesiana, Prolasioptera.

Bermuda, Clastoptera undulata on Casuarina and other plants in, 230; Pseudaulacaspis pentagona on oleander in, 229; Carulaspis minima on Juniperus bermudiana in, 229; indigenous and introduced natural enemies of pests in, 229, 230.

Bermuda Cedar (see Juniperus bermud-

iana).

Bermuda Grass (see Cynodon dactylon). Bessa harveyi (see Ptychomyia selecta). Bessa selecta (see Ptychomyia).

betae, Pegomyia; Pemphigus.

Betaine, in diet of Corcyra cephalonica,

Betula (Birch), Cryptorhynchus lapathi on, in Hungary, 307; pests of, in New York, 19; activity of systemic insecticides in, 19.

betulae, Pulvinaria.

BHC, against ants, 29, 589; against aphids, 59, 84, 188, 217, 352, 468, 498, 537, 624, 633, 667, 668, 674, 677; against Coccids, 120, 557; against Mirids, 7, 76;

against other Hemiptera, 28, 57, 241, 316, 352, 593; ineffective against Aleurolobus barodensis, 498; against Bruchids, 28, 99, 178, 560, 561; against Curculionids, 16, 51, 114, 118, 180, 184, 317, 347, 500, 540, 541, 559, 560, 561, 591, 609, 685; selection for resistance to, in Sitophilus granarius, 128; against Elaterids, 352, 433; against Leptinotarsa decemlineata, 247; resistance to, in strains of L. decemlineata, 252, 440; against Melolonthids, 171, 439, 493, 541, 678, 682, 683; against Scolytids, 7, 31, 451, 555, 606, 643, 644; against other Coleoptera, 27, 31, 40, 49, 51, 58, 61, 104, 118, 138, 173, 201, 225, 237, 244, 297, 316, 349, 396, 436, 489, 497, 500, 587, 593, 608, 647, 674, 685, 687; against Hylemya spp., 50, 103, 226, 254, 362, 568, 674; question of resistance to, in Hylemya, 388; metabolism of, in Musca domestica, 64; against Trypetids, 98, 676; against Diptera, 179, 537, 623; against Noctuids, 16, 27, 179, 345, 347, 409, 538, 539; against Pyralids, 16, 51, 79, 297, 374, 471, 498, 503, 553; against other Lepidoptera, 16, 79, 99, 198, 302, 309, 317, 347, 352, 409, 441, 534, 539, 555, 560, 596, 661; against Tenthredinids, 10, 56, 362, 540, 541, 544, 659, 666; against Vespula germanica, 422; against grasshoppers and locusts, 170, 172, 351, 383, 498, 657; factors affecting resistance to, in locusts, 201; against other Orthoptera, 33, 352; against termites, 77, 180, 553, 577, 627; against Thysanoptera, 345, 352, 499, 503; against mites, 264, 352, 532, 544, 545, 628, 675; mites favoured by, 633, 666; Scutigerella immaculata favoured by, 433; effects of, on injurious soil microfauna, 553; effects of, on insect fauna of pine, 683; effects of, on beneficial insects, 8, 180, 302, 627, 682; distribution of resistance to, in agricultural pests, 284; study on penetration of insect cuticle by, 43; factors affecting toxicity of, 118, 170, 180, 201, 356, 657; in aerosols, 10, 27, 247, 537, 676, 682; in smokes, 178, 195, 683; fumigant action of, 120, 247, 374, 623; vapour toxicity of, 27, 180; in bait-sprays, 98; in dusts, 7, 10, 27, 28, 29, 40, 49, 51, 57, 58, 59, 61, 76, 79, 99, 103, 114, 178, 179, 184, 217, 237, 241, 244, 297, 302, 309, 316, 317, 345, 347, 349, 351, 362, 409, 471, 498, 499, 503, 534, 537, 538, 539, 540, 541, 544, 553, 555, 560, 568, 587, 596, 606, 608, 627, 659, 661, 670, 682, 683; in granules, 396, 568; in

sprays, 7, 16, 28, 29, 31, 33, 56, 57, 79, 80, 84, 98, 128, 138, 170, 179, 180, 188, 195, 198, 213, 217, 225, 252, 264, 297, 317, 324, 351, 362, 383, 441, 451, 468, 498, 503, 532, 541, 544, 545, 555, 557, 559, 568, 593, 600, 633, 643, 644, 657, 666, 668, 675, 677, 683; comparison of high- and low-volume applications of, 670; as deposit, 195, 316, 374; applied from aircraft, 7, 383, 540, 541, 659, 683; seed treatment with, 50, 65, 103, 184, 226, 568, 589, 609, 627, 674; trunk applications of, 451; treatment of planting water with, 388; treatment of planting material with, 77, 297, 362, **557**, **661**, **674**; use of, in trap-logs, 606; treatment of nests with, 422, 627; soil treatment with, 77, 171, 180, 254, 309, 345, 347, 352, 396, 433, 439, 468, 493, 534, 544, 553, 555, 568, 577, 578, 609, 661, 678; persistence of, in soils, 77, 83, 416, 567, 578; treatment of textiles with, 587; treatment of timber with, 31, 489, 647; treatment of thatched roofs with, 596; tests and uses of, against pests of stored products, 28, 49, 51, 99, 120, 178, 179, 195, 213, 225, 347, 374, 436, 500, 560, 561, 591, 685; systemic activity of, 83, 146; solutions of, 118, 383, 537; emulsified solutions of, 56, 57, 77, 170, 172, 188, 252, 297, 316, 324, 489, 555, 441, 677, in wettable-powder suspensions, 28, 56, 57, 77, 80, 179, 362; question of tainting of vegetables and fruit by, 229, 247, 544; phytotoxicity of, 50, 65, 104, 297, 627, 675; other effects of, on plants, 562, 678; residues of, on plants and plant products, 16, 323, 324, 326, 578; residues of, in tissues and eggs of fowls, 326; analysis of residues of, 91; tolerance limits for, 323, 326; and Beauveria bassiana, 541; and chlorbenside, 674; combination of creolin and, 675; and DDT, 10, 61, 99, 179, 198, 247, 252, 297, 345, 349, 409, 441, 503, 538, 539, 540, 608, 674, 676; and dieldrin, 441; and fungicides, 50, 103, 264, 674; and oil, 31; and parathion, 532; synergists with, 587, 687: and whitewash, 374; as lindane, 7, 28, 31, 51, 57, 65, 79, 84, 98, 99, 128, 138, 146, 172, 178, 213, 217, 225, 226, 229, 252, 254, 264, 317, 323, 326, 374, 388, 396, 416, 436, 439, 451, 468, 471, 497, 498, 500, 503, 544, 555, 560, 562, 568, 589, 591, 609, 623, 627, 628, 633, 647, 666, 667, 674, 685; yisomer of, 7, 10, 27, 28, 29, 31, 43, 49, 50, 51, 57, 61, 64, 65, 77, 79, 80, 84, 91, 98, 99, 103,

118, 120, 128, 138, 146, 170, 172, 173, 178, 179, 180, 184, 195, 201, 213, 217, 225, 226, 229, 237, 247, 252, 254, 264, 297, 316, 317, 323, 324, 326, 351, 374, 383, 388, 396, 416, 436, 439, 451, 468, 471, 489, 497, 498, 500, 503, 532, 538, 539, 544, 545, 553, 555, 559, 560, 561, 562, 568, 577, 587, 589, 591, 593, 608, 609, 623, 624, 627, 628, 633, 647, 657, 661, 666, 667, 668, 674, 675, 678, 685, 687.

Bhendi (see Hibiscus esculentus).

bicinctus, Aphrodes.

bicolor, Callitula; Cataglyphis; Hypolithus;

Pegomyia; Perilitus. bifasciata, Comperiella.

bifasciculatus, Chrysomphalus.

biguttata, Lissonota.

biguttula, Empoasca (Chlorita).

bilineata, Aleochara; Hexagenia.

bilituratus, Melanoplus.

biloba, Autographa.

bimaculatus, Tetranychus (see T. telarius).

binubila, Apomecyna.

Bioassay, of insecticides, 146, 300, 339, 397, 412, 417, 484, 680; reviews of uses and methods of, 90, 300.

bioculatus, Perilloides (Perillus).

Biological Control (of insects), reviews and surveys of status and development of, 90, 490, 586, 596, 688, 692; symposium on ecological effects of, 262; survey of data on use of *Bacillus thuringiensis* in, 140; integration of chemical control with, 90, 148, 222; cultural practices and type of crop in relation to use of, 199, 200; equilibrium in commercial cultures of beneficial insects for use in, 282; (of noxious plants), 260, 386, 452, 587, 588, 692.

Biotin, in diet for Corcyra cephalonica, 627.

Biotrol, preparation of Bacillus thuringiensis (q.v.), 421.

biplaga, Earias. bipunctata, Adalia.

bipunctella, Cicadulina.

Birch (see Betula).

Birch Leaf Miner (see Fenusa pusilla).

Birds, Hofmannophila pseudospretella breeding in nests of, in Britain, 290; destroying noxious insects, 13, 15, 166, 363, 460, 538, 565, 598, 638, 683; harmful lizards destroyed by, 229; destroying beneficial insects, 104, 229, 565; occurrence of ants and, 141; flight habits of locusts and, 423; attracted to excretions of Psyllid, 233; effects of insect control on ecology of, 262; study on influence of, on forest fauna, 494; Bacillus thuringiensis not

pathogenic to, 282; effects of agricultural chemicals on, 217, 433, 683.

Birdsfoot Trefoil (see *Lotus corniculatus*). Biscirus, predacious on Sminthurus viridis in Victoria, 34.

bispinosus, Dendrothrips (Scirtothrips).

bisselliella, Tineola.

Bis(2,3,3,3-tetrachloropropyl)ether (see Octachlorodipropyl Ether).

Biston marginata, bionomics of, on Camellia oleifera in China, 592.

Bitumen Emulsion, treatment of fossae with, for storage of grain, 178.

bivittatus, Melanoplus.

Blackberry (see Rubus spp.).

Black-currant Gall Mite (see Cecido-phyopsis ribis).

blancardella, Lithocolletis.

blanchardii, Parlatoria.

Blastothrix sericea, parasitising Eulecan-

ium spp. in France, 683.

Blattella germanica, infesting insect cultures in California, 339; toxicity of extract from Bacillus larvae to, 332; metabolism of insecticides in, 248, 263, 476, 512, 513; used in tests of carbamate insecticides, 516; measures against, 339, 573.

Blattisocius tineivorus, in stored grain in Canada, 154; predacious on Sitotroga cerealella in cultures, 376; acaricides

against, 376.

Blepharidopterus angulatus, predacious on Panonychus ulmi in England, 672; effects of fungicidal sprays on populations of, 672.

Blepharipoda (see Sturmia).

Blissus leucopterus, on cereals, sugarcane and grasses in Colombia, Peru and Venezuela, 98.

Blondelia nigripes, survey of data on, parasitising Choristoneura murinana in

France and Germany, 168.

Blood, Dried, in baits for fire ants, 464. Blood Meal, in baits for *Gryllotalpa*, 442.

Blueberry (see Vaccinium).

Bluegrass, Kentucky (see *Poa pratensis*). Body Louse (see *Pediculus humanus humanus*).

boetica, Cosmolyce (Lampides).

bohemica, Sturmia (Drino).

boisduvalii, Diaspis.

Bollworm (see Heliothis zea).

Bollworm, Pink (see Pectinophora gossy-

piella).

Bombyx mori, infection of Galleria mellonella with polyhedrosis virus of, 496; acceleration of embryonic development of, by parathion, 437; used in tests of insecticides, 171; method of rearing, 606.

bonariensis, Hyperodes.

Borax, treatment of flannel with, ineffective against *Anthrenocerus australis*, 587; treatment of timber with, against other Coleoptera and termites, 485, 648.

Boric Acid, ineffective against Vespula germanica, 422; treatment of flannel with, ineffective against Anthrenocerus australis, 587; treatment of timber with, against other Coleoptera, 648.

bordagei, Apanteles.

Bordeaux Mixture, against Cydia spp., 20, 555; ineffective against Dysmicoccus brevipes, 557; repellent effect of, on Leptinotarsa decemlineata, 305; against Trypetids, 47, 273, 274, 438; effects of sprays containing, on beneficial arthropods, 36; and insecticides, 20, 36, 47, 135, 438, 555, 651; toxicity of insecticides decreased by, 274.

borealis, Lygus (Liocoris).

Boron, treatment of timber with, against Coleoptera, 648; survey of data on use of compounds of, as wood preservatives, 648.

Borrelina aporiae, 683.

Borrelinavirus, 196, 401, 427.

Borrelinavirus reprimens, 400.

bostrichorum, Coeloides.

Bostrychus capucinus, nutritional physiology of, infesting timber, 356.

Boswellia serrata, Coleoptera in logs of, in India, 448; DDT treatment of timber of, 448.

botanephaga, Sesamia nonagrioides.

Bothynoderes (see Cleonus). botrana, Lobesia (Polychrosis).

Bougainville, food-plants of *Leptoglossus* australis in, 235.

Box (see Buxus sempervirens).

Boxiopsis madagascariensis, gen. et sp.n., on cacao in Madagascar, 355.

brachialis, Bruchus.

Brachymeria, parasitising Earias spp. in India, 37.

brachyntera, Thecodiplosis.

Brachypterolus pulicarius, Linaria vulgaris controlled by, in Canada, 386.

Brachyrhinus (see Otiorhynchus). Brachytydeus caudatus (see Tydeus).

Bracon brevicornis, bionomics of, parasitising Depressaria apiella in Czechoslovakia, 301, 302; recorded on roses, 302; factors affecting rate of increase of, 299; destroyed by dusts, 302.

Bracon greeni, parasitising Earias spp.

in India, 37.

Bracon terebella, attacking parasitised Cephus spp., 161; in Sweden and

U.S.S.R., 161; introduced into Canada, 161.

Bracon variator, parasitising Tortricids in U.S.S.R., 112.

Bran, use of, in experiments with spores of *Beauveria bassiana*, 284; residues of insecticides on, 270.

branneri, Trigona hyalinata.

brasiliensis, Aphycus (Metaphycus); Eutinobothrus.

Brassica campestris, Autographa nigrisigna on, in Japan, 79.

Brassica juncea, pests of, in Britain, 318, 373; injury by Meligethes aeneus to, 373.

Brassica nigra, susceptibility of, to Coleoptera in Britain, 318; in relation to infestation of crucifers by Hylemya brassicae in Wisconsin, 516.

Brassica oleracea (see Cabbage).

brassicae, Brevicoryne (Hyadaphis); Hylemya (Chortophila, Erioischia); Mamestra (Barathra); Pieris.

Braunsia, species of, parasitising Cydia funebrana in U.S.S.R., 112.

Brazil, ants in, 120; Citrus pests in, 120, 491, 558, 644, 645; mites in relation to leprosis of orange in, 556; Acroleucus coxalis on pimento in, 356; Dysmicoccus brevipes on pineapple in, 557; pests of other fruits in, 120, 555, 646; Homalinotus coriaceus on coconut in, 559; pests of coffee in, 120, 121, 555, 557, 558, 643; pests of cotton in, 557, 643, 645, 646; new Pyralid on groundnuts in, 145, 646; Sciara spp. infesting mushrooms in, 644; Sternocolaspis quatuordecimcostata on sugar-cane in, 646; Feltia subterranea on tobacco in, 357; pests of vegetables in, 357, 554, 556; pests of stored products in, 119, 558, 644; beneficial insects in, 150, 491.

Bread, question of tainting of, by insecticide treatments of stored grain, 270; question of phosphine residues in, 483; in diets for mites, 518.

brethesi, Psychidosmicra. brevicomis, Dendroctonus.

brevicornis, Bracon (Microbracon); Ephialtes (Epiurus); Nematocerus.

Brevicoryne brassicae (on crucifers), in Britain, 199, 200, 320; in Holland, 608; in New Zealand, 74, 75, 167, 505; in U.S.A., 69, 172, 401, 508; transmission of viruses by, 133, 197; relation of diversity of plant species to populations of, 69; rôle of wavelength in food-plant selection by, 200, 201; resistance to, in varieties of rape and swede, 167; trap catches of, 199, 320; relation of populations of, to other cabbage pests,

172; natural enemies of, 401; fungus infesting, 401; unaffected by *Bacillus thuringiensis*, 75; integration of insecticides with biological control of, 148; tests and uses of insecticides against, 74, 505, 508, 608; resistance to parathion in strains of, 148; sampling methods for, 620.

Brevipalpus, on orange in Israel, 684; toxicity of di(p-chlorophenyl)-trifluoromethylcarbinol to, 684.

Brevipalpus californicus, on lemon in Sicily. 299.

Brevipalpus chilensis, bionomics and control of, on vines in Chile, 495.

Brevipalpus obovatus, on Citrus in Aden,

Brevipalpus phoenicis (on Citrus), in Aden, 80; in Italy and Spain, 299. brevipes, Dysmicoccus (Pseudococcus). brevipilosus, Myzus (see Capitophorus

minor).

brevis, Cryptotermes; Hoplocampa; Tanyrhynchus.

Brick Dust, as carrier for insecticides, 50. Bricks, treatments against termites attacking straw in, 180, 181.

Brinjal (see Solanum melongena).

Britain, apple pests in, 379, 670, 671, 672; mites and virus diseases of black currant in, 70, 287, 432, 671; pests of other soft fruits in, 430, 431, 670, 671; Cicadellids transmitting virus diseases of clover and strawberry in, 686; pests of beans and peas in, 287, 431, 550; pests and virus diseases of beet in, 549; pests of crucifers in, 39, 198, 287, 318, 373, 430; aphids and virus diseases of potato in, 320, 321; Trialeurodes vaporariorum on tomato in, 198; pests of other vegetables in, 430, 431, 432; pests of broom in, 30, 656; pests of cereals in, 50, 51, 288, 319, 320, 321, 420, 430, 433; pests of grasses in, 71, **321**, **430**, **487**, **488**; pests of carnations in, 431; Merodon equestris on Narcissus bulbs in, 288; Xyleborus morigerus in, on orchids originating from New Guinea, 625; Hepialus lupulinus on ornamental plants in, 430, 431; guide to diseases and pests of horticultural plants for inspection officers in, 425; Bupalus piniarius on pines in, 550; other virus diseases of plants in, 70, 71; pests of stored products in, 91, 295, 374, 548, 549, 657, 694; Hofmannophila pseudospretella infesting textiles and houses in, 290; Anobium punctatum infesting furniture and building timber in, 489; accidental introduction of termites into, 321; Gregarines infesting

cultures of *Tribolium* in, 582, 583; Anthrenus verbasci in, 249; Formica aquilonia described from, 104; Tyria jacobaeae in, 260, 452; high-altitude catches of insects in, 199; incidence of Acarapis woodi in honey bees in, 617; part of work on terrestrial Acari of, 41; fungus infesting Plutella maculipennis in, 198; natural enemies of pests in, 30, 52, 91, 161, 379, 489, 493, 522, 549, 672; restrictions on use of seed treatments in, 433; reports on hazards associated with use of toxic chemicals

in agriculture in, 432, 433.

British Columbia, pests of apple in, 388, 394, 395, 524; Psylla pyricola on pear in, 158, 395, 411; pests of loganberry and other Rubus spp. in, 395, 529; pests and virus diseases of strawberry in, 529, 692; pests of potato in, 424, 526, 529; pests of other vegetables in, 134, 529; forest pests in, 126, 207, 208, 385, 386, 395, 423, 424, 457, 458, 529; Camnula pellucida on rangelands in, 158; pests of ornamental plants in, 529; Melanoplus bilituratus in, 155; Acarapis sp. infesting honey bees in, 295: Typhlodromus caudiglans misidentified as T. rhenanus in, 522; beneficial insects in, 126, 158, 386, 457; new bacterial diseases of Malacosoma pluviale in, 342.

Broccoli, pests of, in U.S.A., 222, 335,

404.

bromicola, Stenodiplosis.

2-Bromomercurithiophen, laboratory tests of toxicity of, to insects, 202, 203; effect of temperature on toxicity of, 202, 203.

6-Bromopiperonal Chrysanthemate, toxicity of, to Estigmene acraea, 480.

1-Bromopropane, vapour toxicity of, to

Phylloxera vitifoliae, 572.

Bromus, pests of, in U.S.S.R., **542**, **596**. Bromus inermis, relation of mites and thrips to silver top of, in Alberta, **156**; new Cecidomyiid infesting, in U.S.S.R., **542**; in tests of food-plant preferences of Thymelicus lineola, **162**.

Bromus riparius, new Cecidomyiid infest-

ing, in U.S.S.R., 542.

Bromus unioloides, survival of Chloriona orizicola on, 132.

Brontispa longissima, distribution of, in Pacific area, 81; on coconut and other palms, 81.

Broom (see Sarothamnus scoparius).

Bruchidius algiricus, in Algeria and Morocco, 249; on broad beans and flax, 249.

Bruchidius atrolineatus, infestation of

field and stored cowpeas by, in Nigeria, 49, 50; measures against, 50.

Bruchidius incarnatus, in Algeria and Tunisia, 249; on broad beans, flax, chick-pea, Dolichos and peas, 249.

Bruchidius lentis (see Bruchus).

Bruchidius perparvulus, bionomics of, on plants and stored seed of crimson clover in Italy, 182; question of fumigants against, 183.

Bruchophagus (see Eurytoma).

Bruchus brachialis, on hairy vetch in Oregon, 211; effects of insecticides against cereal pests on populations of, 211.

Bruchus chinensis (see Callosobruchus). Bruchus lentis, on flax in Algeria, 249.

Bruchus pisorum (pisi), bionomics of, on pea in Bulgaria, 58, 537; parasites of, 58.

brumata, Operophtera (Cheimatobia). bruneri, Lachnosterna (Cnemarachis, Phyllophaga).

brunnea, Maladera (Serica).

brunneus, Ephialtes (Epiurus); Lyctus.

brunnipes, Macrocoma.

Brussels Sprouts, *Brevicoryne brassicae* on, in Holland, **608**; question of insecticide residues in, **608**.

Bryobia, on apple in Poland, 676; dormant sprays against eggs of, 676.

Bryobia arborea (see B. rubrioculus).
Bryobia cristata, on grasses, 444; differences between bionomics of congeners and, 444.

Bryobia kissophila, on ivy, 444; differences between bionomics of congeners and, 444.

Bryobia praetiosa, auct. (see B. rubrio-culus).

Bryobia praetiosa Koch, on apple in Poland, 677, 678; infesting houses in U.S.A., 419, 519; on grasses, 444, 519; effects of plant nutrition on development of, 519; differences between bionomics of congeners and, 444; acaricides against, 419, 677, 678.

Bryobia redikorzevi, on apple in Bulgaria,
60, 111; on fruit trees in U.S.S.R.,
242; bionomics of, 242; sprays against,
60; effects of DDT on fecundity of, 111.

Bryobia rubrioculus (on fruit trees), in France, 8; in Germany, 439; in Holland, 444; in Nova Scotia, 259, 523; differences between bionomics of congeners and, 444; factors affecting populations of, 259, 444; predators of, 122, 522, 523; ovicides against, 8; favoured by methyl-DDD, 523; effects on predacious mite of acaricides against, 439.

Bucarpolate (see 2-(2-n-Butoxyethoxy)ethyl 3,4-Methylenedioxybenzoate).

Bucculatrix ainsliella, bionomics parasites of, on red oak in Michigan, 327.

Bucculatrix thurberiella, on cotton in U.S.A., 508, 638, 639; natural enemies of, 638, 639; tests and uses of insecticides against, 508, 516.

Budworm, Spruce (see Choristoneura

fumiferana).

Buffel Grass (see Cenchrus ciliaris).

Building Timber, Coleoptera in, 101, 112, 142, 253, 355, 489, 646; termites attacking, 33, 76, 180, 296, 416, 485; treatments of, 253, 254, 296, 485, 486; survey of data on treatments of, 646. Bulan, mixture of Prolan and (see Dilan).

Bulb Fly, Wheat (see Hylemya coarctata). Bulgaria, pests of cereals in, 109, 348, 537; pests of fruit trees in, 58, 59, 60, 110, 111, 348, 540; vine pests in, 58, 60, 110, 348, 538, 539; Harpalus rufipes on strawberry in, 59; Curculio elephas on chestnut in, 60; pests of beet in, 59, 347, 348, 536, 541; Heliothis spp. on mint in, 59; Megaselia pygmaeoides on mushrooms in, 59; pests virus diseases of solanaceous vegetables in, 236, 345, 366, 539; pests of other vegetables in, 58, 59, 60, 346, 348, 349, 536, 537; pests of leguminous forage crops in, 59, 60, 537, 538, 539, 540; Thrips tabaci and virus disease of tobacco in, 345; pests of stored products in, 57, 347; primary food-plants of Aphis fabae in, 536, 537; Melolontha spp. in, 355; annotated list of new or rare insects in, 59; beneficial insects in, 58, 59, 111, 346, 367, 537, 540; fungus diseases of insects in, 348, 541.

buoliana, Rhyacionia (Evetria).

buolianae, Ephialtes (Epiurus, Scambus); ${\it Lissonota}.$

Bupalus piniarius, on pines in Britain, 550; in Holland, 205; diapause in, 205; parasites of, 205, 379; sampling method for, 550: DDT aerosol against, 550; characteristics and world distribution of, 507.

Buprestis haemorrhoidalis, infesting conifers and building timber in Finland,

101.

Buprestis rustica, infesting building timber

in Finland, 101.

Burma, Aonidiella aurantii on Citrus in, 491; parasites of Coccids in, 266, 491. burrelli, Allotropa.

bursarius, Pemphigus.

buscki, Drosophila.

Busseola, characters of pupae of, 292.

Busseola fusca (on cereals), in Kenya, 179; in Nigeria, 292; in S. Rhodesia, 48; bionomics and parasites of, 292; fungus disease of, 292; relation between moisture and pupation in, 48; effects of, on yields of millet and maize, 293; measures against, 292, 293; characteristics and world distribution of, 507.

Bustomus setifer, bionomics of, on vines

in S. Africa, 650.

Bustomus setulosus, bionomics of, on

vines in S. Africa, 650.

2-(2-n-Butoxyethoxy)ethyl 3.4-Methylenedioxybenzoate, as synergist with pyrethrum, 2, 3.

β-Butoxy β'-Thiocyanodiethyl Ether, toxicity of turnip constituent and, 484.

sec.-Butyl trans-4(or 5)-Chloro-2-methylcyclohexane-1-carboxylate (see Medlure).

Butyl 1,3-Dimethylbutyl Acetal of Acetaldehyde, toxicity of, to Estigmene acraea, 480.

sec.-Butvl 6-Methyl-3-cyclohexenecarboxylate, in baits for Trypetids, 643.

n-Butyl p-Nitrophenyl Ethylphosphonate, anticholinesterase activity and toxicity of, to Musca domestica, 407.

3-tert.-Butylphenyl Dimethylcarbamate, hydrolysis and anticholinesterase activ-

ity of, 635.

m-sec.-Butylphenyl Methylcarbamate. tests of toxicity of, to insects, 515, 516.

m-(or 3-)tert.-Butylphenyl Methylcarbamate, tests of toxicity of, to insects, 515, 516; hydrolysis and anticholinesterase activity of, 635.

buxi, Monarthropalpus.

Buxus sempervirens, Monarthropalpus buxi on, in Italy, 694.

Byrsocrypta ulmi, on maize in Italy, 349; unknown in Israel, 349.

Byturus urbanus (tomentosus, auct.), on Rubus spp. in Britain, 671; malathion sprays against, 671.

Cabbage, Pieris brassicae on, in Austria, 184; Hylemya brassicae on, in Britain, 287; pests of, in Bulgaria, 60, 346, 349; pests of, in Canada, 161, 162, 211, 385, 525; Ceutorhynchus pleurostigma on, in Holland, 673; Trichoplusia ni on, in India, 424; pests of, in Japan, 78, 79, 449; pests of, in New Zealand, 75; P. brassicae on, in Poland, 545; Eurydema ornatum on, in Rumania

57; pests of, in U.S.S.R., 114, 244, 245, 362; pests of, in U.S.A., 172, 222, 269, 508, 510; nature of transmission of virus disease of, by Brevicorvne brassicae, 133; effects of H. brassicae on biochemistry of, 287; detection of growth inhibitors in, 22; distribution of P. rapae in fields of, 449; relations between populations of pests on, 172; hemp not protecting fields of, from P. brassicae, 545; treatment of, with nematode against pests, 162, 163; treatments of seedlings of, 362; factors affecting systemic activity of insecticides in, 508; as food for Schistocerca gregaria, 170.

Cabinets, for rearing insects, 339.

Cable, Lead, damaged by Dicranura erminea menciana in China, 693.

Cacao, aphids on, in Dominican Republic, 144; Mirids on, in Ghana, 651; pests of, in Ivory Coast, 7; Xyleborus morigerus on, in Java, 625; new Mirid on, in Madagascar, 355; pests of, in New Guinea, 76, 196, 234; pests of, in Nigeria, 294, 651, 652; termites attacking seedlings of, 294; swollen-shoot virus of, 651, 652; die-back of, 652; question of value of BHC sprays against pests of, 7, 8.

Cacao (Stored), question of infestation of, by Ahasverus advena, 657; Cadra cautella in, 644; residues of methoxy-

DDT in. 644.

Cacoecia crataegana (see Archips). Cacoecia rosana (see Archips).

cacoeciae, Trichogramma.

cadaverinus, Dermestes (see D. ater).

Cadra cautella, in stored cacao in Brazil, 644; in stored groundnuts in Gambia, 178; intercepted on oil-seed cake in Germany, 11; in stored maize in Kenya, 51; in harvested dates in Morocco, 54; susceptibility of sexes and stages of, to pyrethrins, 3, 4; insecticides against, 51, 54, 644.

caeruleus, Korynetes. caesar, Zenillia (Eumea). caespitum, Tetramorium, caffeina, Leucoptera.

Cages, for rearing insects, 290, 390, 496, 633; escape of Tachinid from, 53; for collection of excreta from weevil, 575; for use in pollination studies, 620. Calacarus citrifolii, on Citrus, granadilla and arum lily in S. Africa, 83.

calamistis, Sesamia.

Calandra (see Sitophilus).

Calcalia, Tyria jacobaeae feeding on, 260. calcitrator, Collyria.

Calcium Ammonium Nitrate (as fertiliser). effects of, on populations of spruce

pests, 253.

Calcium Arsenate, against Lepidoptera, 80, 138; against Rhagoletis pomonella, 273, 274; against Leptinotarsa decemlineata, 683; effects of, on natural enemies of pests, 378, 683; effects of, on growth of cotton, 220, 632; in dusts, 138; in sprays, 80, 220, 273, 274, 632, 683; and lime, 80.

Calcium Carbonate, effect of, on toxicity

of insecticide dusts, 118.

Calcium Cyanide, treatment of rubber trees with, against termite, 77; treatment of wasps' nests with, 422.

Calendra (see Sphenophorus).

California, almond pests in, 405, 482; Monellia costalis on pecan in, 151; pests of walnut in, 128, 151, 276, 482, 491; avocado pests in, 200, 378; Otiorhynchus cribricollis spreading from blackberry to globe artichoke in, 509; pests of Citrus in, 22, 147, 267, 378, 509, 519, 561, 688; *Drosophila* in lemon packing houses in, 573; new Lonchaeid on currant in, 128; Parlatoria oleae on olive in, 227; Drosophila damaging grapes in, 265; pests of strawberry in, 217, 563; pests of other fruits in, 70, 128, 265, 271, 402, 481, 482; virus diseases of fruits in, 70; forest pests in, 23, 124, 125, 221, 340, 341, 355, 385, 386, 642: Lepidoptera on cotton in. 508; pests of cereals and grasses in, 421, 508, 509, 520, 580; lucerne pests in, 66, 67; pests and virus disease of Capsicum in, 216, 636, 637; Myzus persicae and Circulifer tenellus transmitting virus diseases of melon in, 408; Eotetranychus lewisi on poinsettia in, 517; pests and virus diseases of beet in, 17, 131, 331, 419, 463, 630; pests of other vegetables in, 67, 131, 480, 508, 509, 569; virus diseases of vegetables in, 131, 480; Tenebrionids infesting stored-grain products in, 420; other insects in, 66, 150, 151, 265, 279, 353, 491, 563; ants associated with aphids and Coccids in, 419, 519; question of climatic factors limiting establishment of Dacus dorsalis in, 563; question of Typhlodromus rhenanus in, 522; pests infesting insect cultures in, 339; mites infesting honey bees in, 295; fungi infesting insects in, 509, 570; natural enemies and biological control of insects in, 21, 22, 24, 66, 67, 124, 150, 151, 200, 221, 222, 227, 266, 353, 378, 386, 400, 491, 519, 521, 564, 630, 641, 688; quarantine regulations in, 403;

tolerance limits for insecticides in, 267.

californica, Plusia (Autographa).

californicus, Apanteles; Aphycus (Meta-

phycus); Brevipalpus.

Caligo eurilochus, bionomics of, on banana in Ecuador, 267, 274; natural enemies of, 267, 268, 274.

Caliothrips, insecticides against, on cotton

in Brazil, 645.

Caliroa limacina (cerasi), on hawthorn and wild cherry in Quebec, 26; on fruit trees, 26; bionomics and parasite of, 26.

Calla, oviposition of Tetranychus telarius on, 547.

Calliephialtes (see Ephialtes).

Calligypona pellucida (marginata), on cereals in Finland, 102; on grasses in France, 373; other food-plants of, 102; stage or variety of oats in relation to, 103; method of rearing, 373; synonymy of, 349.

Calligypona propinqua, on Bermuda grass in Israel, 349; experiments on transmission of maize virus by, 349.

Calligypona striatella, transmitting virus disease to maize and grasses in Israel, 349; on maize in Italy, 350; transmitting cereal diseases in Japan, 350; bionomics of, 350; considered a synonym of C. marginata, 349.

Callimorpha quadripunctaria, parasite of,

87.

Calliptamus palaestinensis, factors affecting development of, in Israel, 142, 143. Callistephus (Aster), yellows virus of, in U.S.A., 271.

Callistephus chinensis, transmission of stolbur virus to, by Aphrodes bicinctus, 304.

Callitula bicolor, parasitic on Oscinella

frit in Germany, 45.

Callosobruchus chinensis, cuticular structure in relation to contact toxicity of insecticides to, 314; effects of heat treatment on survival and fecundity of, 449.

Callosobruchus maculatus, in stored beans in Angola, 99; in stored beans in India, 317, 500; in stored cowpeas in Nigeria, 49, 50; types of sack in relation to infestation by, 501; measures against, 50.

callosus, Phlyctinus; Sphenophorus

(Calendra).

Calocoris, bionomics of species of, on peach and other plants in Italy, 310. Calocoris fulvomaculatus, 310.

Calocoris norvegicus norvegicus, 310. Calocoris norvegicus vittiger, 310. Calocoris trivialis, 310.

Caloglyphus berlesei, in stored grain in U.S.A., 518; food preferences of, 518. Calomycterus setarius, experimental infestation of, by Beauveria bassiana, 284

Calonectria rigidiuscula, association of Mirids with, on cacao in Nigeria, 652. Calosoma maderae auropunctatum, predacious on Apamea anceps in U.S.S.R.,

Calotermes (see Kalotermes).

Calotropis, Lygaeus pandurus on, in India, 315.

Camarota curvinervis, in Bulgaria, 109, 110; in Yugoslavia, 249; bionomics of, on cereals and grasses, 110, 249.

cambiicola, Dioryctria.

Camel-thorn (see Alhagi camelorum).

Camellia oleifera (oleosa), Biston marginata on, in China, 592.

Cameroons, British (see Cameroun Republic and Nigeria).

Cameroun Republic, new weevil on Cola in, 355; predacious ant in, 28.

Camnula pellucida, on rangelands in British Columbia, 158; effects of temperature on development of, 158; infected by Nosema locustae, 582.

campbelli, Eriococcus. campestris, Gryllus.

Camphene, Chlorinated, in dusts against Stylopalpia costalimai, 145; soil treatment with, against larvae of Feltia subterranea, 357.

Campogaster exigua, parasitising Sitona in France, 414; introduced into Manitoba against S. cylindricollis, 414.

Campoletis oxylus, 130.

Camponotus, relation of, to fungus disease of aphids in Congo, 610.

Campoplex, parasitising Rhyacionia rigidana in Ohio, 20.

Campoplex atramentaria, parasitising Eucosma nigricana in Czechoslovakia, 602; effects of sprays on populations of, 603.

Campoplex crassifemur, parasitising Neurotoma nemoralis in Bulgaria, 540.

Campoplex rothi, parasitising Eucosma nigricana in Czechoslovakia, 602; effects of sprays on populations of, 603.

campyla, Cheumatopsyche. Campylomma diversicornis, on cotton in

Turkey, 82.

Campylomma nicolasi, on cotton in

Turkey, 82.

Canada, pests of cereals and grasses in, 71, 160, 393, 394, 420; Aceria tulipae transmitting wheat mosaic viruses in, 71; pests of leguminous forage crops

in, 72, 390; experiments with Aphrodes bicinctus as vector of phyllody virus of clovers in, 686; forest pests in, 25, 107, 121, 122, 123, 130, 185, 208, 210, 341, 342, 391, 394, 458, 474, 522, 529, 532, 580, 586, 619; Lepidoptera on cabbage in, 161; Psorosina hammondi on apple, pear and Prunus maritima in, 529; pests and virus diseases of strawberry in, 23, 686; timing of grasshopper survey in, 575; other insects in, 161, 257, 318, 377, 429, 585; pests of stored grain in, 154, 424; Acarapis spp. on honey bees in, 342, 505; review of insects and mites of potential importance in, 528, 529; Cordyceps ravenellii infecting Lachnosterna larvae in, 619; virus diseases of sawflies in, 341, 342; natural enemies of pests in, 73, 74, 161, 257, 377, 378, 390, 391, 394, 429, 522, 586, 614, 692, 693; insects in relation to control of weeds in, 260, 386; reviews of biological control in, 586, 692, 693. (See also under individual Provinces.)

Canadian Poplar (see *Populus deltoides*).

canaliculatus, Pleonomus,

Canary Islands, forest pests in, 659; Cryptotermes brevis in, 300; Acarapis woodi infesting honey bees in, 296.

cancellatus, Carabus. canescens, Nemeritis.

Cankerworm (see Alsophila pometaria). Cannabis sativa (Hemp), Ostrinia nubilalis on, in Hungary, 306; mites infesting seed of, in Poland, 364; O. nubilalis on, in U.S.S.R., 14; ineffective in protection of cabbages from infestation by Pieris brassicae, 545.

Cantheconidea furcellata, predacious on Earias spp. in India, 37.

canusella, Evagora (Recurvaria).

Cape Gooseberry (see Physalis peruviana). capensis, Cryptolestes; Nudaurelia cytherea.

capitata, Ceratitis.

capito, Locusta migratoria.

Capitophorus fragaefolii (on strawberry), in Canada, 23, 692; in Germany, 106, 546; in Holland, 192; in U.S.A., 23, 217; on other rosaceous plants, 23; occurrence of apterous males of, 23; transmission of strawberry viruses by, 23, 106, 107, 192, 441, 546, 692; insecticides against, 192, 217.

Capitophorus galeopsidis, on potato in Maine, 570; bionomics of, on currants in U.S.S.R., 595, 596; Galeopsis tetrahit as summer food-plant of, 596; fungus infesting, 570; characters distinguishing

C. ribis from, 596.

Capitophorus jacobi, on Fragaria spp. in U.S.A., 23.

Capitophorus minor, on strawberry in Canada and U.S.A., 23; synonymy of. 23.

Capitophorus potentillae, in N. America,

Capitophorus ribis, on black currant in U.S.S.R., 596; characters distinguishing C. galeopsidis from, 596.

Capitophorus tetrarhodus, in N. America, 23, 692; on rose, 23; transmission of

strawberry viruses by, 23, 692.

Capitophorus thomasi, in Canada, 23, 692; in U.S.A., 23; on strawberry, rose and Potentilla spp., 23; transmission of strawberry viruses by, 23, 692. Capnodis cariosa, on pistachio in Turkey,

609.

Capnodis tenebrionis, characteristics and

world distribution of, 507.

Capnodium walteri, associated Eriococcus orariensis on Leptospermum in New Zealand, 587.

Capsaicin, rôle of, in resistance of solanaceous plants to Coleoptera, 608.

Capsicum, new virus disease of, in Bulgaria, 345: Zonosemata electa on. in Ontario, 529; pests of, in U.S.A,, 216, 630, 636, 637; virus disease of, 636; systemic activity of insecticides in, 216.

Capsicum annuum, Leptinotarsa decemlineata not feeding on, 366.

Capsicum frutescens (see Pimento).

Captan (sprays containing), 264, 324, 439; against Cydia pomonella, 20, 21; against Rhagoletis pomonella, effects of, on populations of Tetranychids, 108, 523; effects of, on arthropods on apple, 159; effects of, on beneficial arthropods, 36, 672; toxicity of insecticides decreased by, 274.

capucinus, Bostrychus.

Carabunia myersi, parasitising Clastoptera undulata in West Indies, 230; introduced into Bermuda, 230.

Carabus. species of, predacious Leptinotarsa decemlineata in Germany, 684.

Carabus auratus, 684.

Carabus cancellatus, 684.

Carabus granulatus, 684.

Caragana, Aphis craccivora on, in Bulgaria, 59.

Caraway (see Carum carvi).

Carbolineum, 15, 60.

Carbon (Charcoal), in substrates for mites, 518; in formulations for plant treatments with systemic insecticides. 218, 277, 278, 398, 399, 401; interaction of, with Ceresan, 399.

Carbon, Activated, as carrier for insec-

ticides, 556.

Carbon, Radioactive, used in study of oviposition by *Gryllus assimilis*, 686; insecticides labelled with, 218, 263.

Carbon Bisulphide (as furnigant), against Chloropsalta viridissima, 609; against Lachnosterna consanguinea, 171; trunk injection with, against larvae of Saperda carcharias, 601; treatment of nests with, against Vespula germanica, 422; against pests of stored products, 215, 267, 274, 347, 466, 499; sorption of, by stored wheat, 215; against termites in pianos, 296; factors affecting toxicity of, 274, 466; effects of, on seed germination, 579; as solvent for p-dichlorobenzene, 609; in mixtures with other fumigants, 215, 499, 579.

Carbon Dioxide, treatment of *Narcissus* bulbs with, against *Merodon equestris*, **42**; measurement of output of, by insects, **40**; anaesthetisation of insects

with, 356, 416.

Carbon Tetrachloride (as fumigant), against pests of stored products, 118, 215, 267, 274, 499; sorption of, by stored wheat, 215; factors affecting toxicity of, 274; effects of, on seed germination, 579; determination of, in air, 496; in mixtures with other fumigants, 118, 215, 499, 579.

Carcelia evolans, bionomics of, parasitising Diparopsis watersi in Nigeria, 53; effects of sprays against D. watersi on populations of, 53; cages for, 53.

Carcelia obesa, parasitising Bupalus piniarius in Holland, 205; diapause of B. piniarius in relation to, 205.

carcharias, Saperda.

Cardiaspina, species of, on Eucalyptus in Australia, 552.

cardinalis, Rodolia. cardui, Vanessa.

carinatus, Hadronotus.

Carinodes, parasitising Oiketicus kirbyi in Costa Rica, 638; parasitising Ceramidia viridis in Ecuador, 353.

cariosa, Capnodis.

Carissa grandiflora, Aonidiella orientalis

on, in India, **491.**

Carnation, pests of, in Britain, 431; Tetranychus cinnabarinus on, in France, 550; Sciara amoena on, in Germany, 254; effect of plant age on systemic activity of insecticides in, 431.

carolina, Dissosteira.

Carolina, North, Anthonomus grandis on cotton in, 464; Hypera variabilis on

lucerne in, 331; Heliothis zea on maize in, 327; Diaphania nitidalis on squash in, 471; forest pests in, 336, 418, 513; pests of stored and field tobacco in, 152; fire ant in, 517; Conotrachelus nenuphar in, 216; natural enemies of pests in, 152.

Carolina, South, cotton pests in, 322, 323; Lepidoptera on tobacco in, 465; Conoderus falli in, 575; Protoparce

sexta in, **635**.

Carolinaia setariae, on grass in Wiscon-

sin, **160.**

β-Carotene, and pigmentation in Schistocerca, 5; effect of infection by Nosema locustae on content of, in Locusta migratoria, 582; in diet for Anthonomus grandis, 404.

Carpets, Hofmannophila pseudospretella

infesting, 290.

carpini, Eotetranychus; Trypopitys

(Priobium).

Carpinus betulus (Hornbeam), Eotetranychus carpini on, in Italy, 183; study on influence of birds on insect fauna of, 494.

Carpocapsa (see Cydia).

Carpophilus hemipterus, flight habits of, in California, 265.

Carposina sasakii (niponensis), on fruit trees and hawthorn in U.S.S.R., 112; parasites of, 112; characteristics and world distribution of, 507.

Carrots, *Psila rosae* on, in Britain, 432; pests of, in Canada, 70, 524, 529; pests of, in Holland, 100, 299, 608, 673; *P. rosae* on, in New York, 226; comparison of injury by *Phytomyza lateralis* and *Psila rosae* to, 674; insect pollination of, 341; aster-yellows virus and, 70, 524; seed treatment of, 100, 101, 226, 299, 608; effects of insecticides on germination of, 101; question of heptachlor residues in, 479.

Carulaspis minima (visci), on Juniperus bermudiana in Bermuda, 229; Coccinellids against, 229.

Carum carvi, Depressaria apiella on, in Czechoslovakia, 301.

Carya (Hickory), Acrobasis spp. on, in Ontario, 529; Ennomos subsignarius on, in U.S.A., 418.

Caryedon fuscus, in stored groundnuts in Senegal, 298.

Caryedon gonagra, infesting harvested groundnuts in Gambia, 28, 178, 179; treatment of stored groundnuts against, 28, 179.

casalis, Haemolaelaps.

Casca occidentalis, parasitising Quadraspidiotus perniciosus in Switzerland, 664.

casei, Piophila.

Casein, in diets for insects, 5, 6, 442, 495,

Casein Hydrolysate, in diet for Anthonomus grandis, 404.

Casinaria, parasitising Oiketicus kirbyi in Costa Rica, 638; synonymy of, 493.

Cassava (see Manihot esculenta).

Cassava Flour, treatment of rubber roots with paste containing insecticides and, against termites, 77.

castanea, Diparopsis. castaneum, Tribolium. castaneus, Hemiteles.

Castnia, damaging Heliconia in Costa Rica and Honduras, 520.

Castor (see Ricinus communis).

Castor Oil, as adhesive for seed treatments, 670; toxicity of, to beet, 670.

Casuarina, Clastoptera undulata on, in Bermuda, 230.

Catabena esula, parasitised by Hyposoter exiguae in California, 66.

Cataglyphis bicolor, predacious on Myiopardalis pardalina in Turkey, 352.

Cataglyphis megalocola, predacious on Myiopardalis pardalina in Turkey, 352.

catalaunalis, Antigastra.

Catalpa, Pseudococcus comstocki on, in Ontario, 164.

catenarius, Cerococcus.

Catocala nymphaea (see Ephesia).

Catocala nymphagoga, natural enemies of, on forest trees in Spain, 381.

Catolaccus cyanoideus, parasitising Pseudaletia unipuncta in Ontario, 130.

Cattle, losses of, attributed to destruction of pasture by Spodoptera exempta in Kenya, 530; transport of Eriococcus orariensis by, 587; insecticide residues in tissues and milk of, 64, 92, 146, 147, 225, 384, 483, 521.

Cattleya, Diaspis boisduvalii on, in Indonesia, 143; damaged by cellulose polymer, 272.

caudatus, Triaspis; Tydeus (Brachytydeus). caudex, Melanotus.

caudiglans, Typhlodromus.

Cauliflower, Plutella maculipennis on, in Britain, 198; Hylemya brassicae on, in Holland, 39; Eurydema ornatum on, in Rumania, 57; pests of, in U.S.A., 328, 508, 510; relation of, to oviposition by H. brassicae, 516; Tetranychus multisetis feeding on, 67; experiments on transmission of virus disease of, by aphids, 197; as food for Locusta migratoria, 172.

caulophaga, Haplomyza.

cautella, Cadra (Ephestia).

cavus, Dibrachys.

Cecidophyopsis (Cecidophyes) ribis (on black current), in Britain, 70, 287, 432, 671; bionomics of, in Holland, 302; transmitting reversion virus, 70, 302; measures against, 287, 302, 432, 671, 672.

Cedar (see Cedrus libani).

Cedar, Bermuda (see Juniperus bermudiana).

Cedar, Siberian (see Pinus cembra).

Cedrus libani, pests of, in Turkey, 355. Celcure, treatment of timber with, against

termites, 485; composition of, 485. Celerio lineata livornica, sprays against,

on vines in Italy, 48.

Celery, Spodoptera exigua on, in California, 509; question of Macrosteles fascifrons transmitting aster-yellows virus to, in Ontario, 70.

Cellosize WP-4400, 272. Cellulose, in diets for locusts, 5.

Cellulose Polymers, toxicity of, to arthropods and mammals, 272; phytotoxicity of, 272; efficacy of mevinphos increased

by, 272. Cement, treatment of fossae with, for

storage of grain, 178. Cenchrus ciliaris, ants preventing establishment of, in Queensland, 589.

Centrifugal Force, effects of, on larvae of Pectinophora gossypiella, 479.

Centrococcus insolitus, new parasite reared from, in India, 91.

Centrosema plumieri, experiments with groundnut virus and, 611.

Centrotus cornutus, on vines in Bulgaria,

Cephalcia (Cephaleia) abietis, on spruce in Czechoslovakia, 196, 601; diapause in, 601; natural enemies of, 601; traps for, 601.

Cephalcia arvensis, on spruce in Czechoslovakia, 601; diapause in, 601; natural enemies of, 601; traps for, 601.

Cephaloglypta murinanae, survey of data on, parasitising Choristoneura murinana in France and Germany, 168.

cephalonica, Corcyra.

Cephalonomia tarsalis, in stored grain in Canada, 154.

Cephus cinctus, on cereals in Canada, 72, 160, 527; parasites of *C. pygmaeus* against, 160, 161; factors affecting resistance of cereals to, 72, 526, 527; effects of freezing on larvae of, 551.

Cephus pygmaeus, in Canada, 161; in Sweden and U.S.S.R., 160; parasites of, 160, 161.

Ceracia mucronifera, bionomics of, parasitising Anacridium aegyptium in France, Spain and Tunisia, 313.

Ceramidia viridis, on banana in Ecuador, 353; parasites of, 353; sprays against, 353.

cerasi, Caliroa (see C. limacina); Monoctonus; Myzus; Rhagoletis.

cerasivoranus, Archips.

Cerasus (see Prunus).

Ceratitis capitata, in Brazil, 643; in France, 668; in Gaza Strip, 97; in Morocco, 435; in Spain, 98; in Yugoslavia, 239; on apricot, 98; on Citrus, 97; on coffee, 643; on fig, 239; on peach, 239, 668, 669; on pear, 668, 669; on persimmon, 239; bionomics of, 239; parasite of, 435; traps for, 643; baits and attractants for, 98, 383, 384, 643; insecticides against, 98, 240, 603, 668, 669.

Ceratitis rosa, reared from pomegranate

fruits in S. Africa, 531.

Ceratocystis ulmi (causing Dutch elm disease), relation of Scolytids to, 124, 217; in Canada, 124; in U.S.A., 217. Ceratomegilla fuscilabris, predacious on

aphids in New York, 401.

ceratoniae, Ectomyelois (Myelois).

Cercopis sanguinolenta, on vines in Bulgaria, 110.

Cercospora, mixed sprays against, on banana, 463.

cerealella, Sitotroga.

cerealis, Eremnus. cerealium, Limothrips.

Cereals, Pseudaletia unipuncta on, in N. America, 393, 394; aphids on, in Australia, 684; Macrosiphum avenae on, in Belgium, 375; Hylemya coarctata on, in Britain, 433; pests of, in Bulgaria, 348; pests of, in Canada, 70, 524, 584-586; damaged by Lamellicorn larvae in China, 594; Hydrella griseola on, in Denmark, 354; aphids on, in Egypt, 684; Hepialus lupulinus on, in France, 7; Hyalesthes obsoletus on, in Hungary, 307; pests of, in India, 498, 684; aphids on, in Italy, 684, pests of, in Japan, 350, 684; aphids on, in Jordan, 684; pests of, in Kenya, 180, 530; Pentatomids on, in Middle East, 654, 655, 656; Pentatomids on, in Morocco, 382, 653, 654, 655; aphids on, in New Zealand, 684; Lepidopterous stem borers of, in Nigeria, 292, 293; aphids on, Pakistan, 684; Eurygaster integriceps

on, in Persia, 653; Trigonotylus rufi-cornis on, in Sweden, 354; Stenodema laevigatum on, in Turkey, 352; pests of, in U.S.S.R., 13, 14, 15, 243, 244, 344, 345, 363, 543, 596; pests of, in U.S.A., 95, 160, 520, 639; Phyllotreta vittula on, in Yugoslavia, 181; FAO report on control of Pentatomids on, 300; virus diseases of, 95, 350, 375, 684; feeding behaviour of aphids in relation to spread of virus disease of, 160; forecasting abundance of pests of, 15; destruction of débris of, in control of stem borers, 293; use of X-rays for detection of stem borers in, 495; restriction on use of seed treatment of, 433; (products of), in baits for fire ants, 464; in diets for Ahasverus advena, 657. (See Maize, Wheat, etc.) Cereals (Stored), Cephalonomia tarsalis

in, 154; Coleoptera in, 2, 3, 10, 11, 17, 154, 155, 178, 196, 205, 206, 295, 300, 344, 347, 414, 415, 420, 500, 529, 548, 558, 632, 658; disturbance of, in relation to temperature requirements of Ptinus spp., 344; effect of type of grain on development of Sitophilus spp. in, 205, 206; Lepidoptera in, 10, 11, 596; relation of Sitotroga cerealella to infestation of, by Aspergillus spp., 414, 415; mites in, 91, 149, 154, 155, 424, 518, 529; insects intercepted in importations of, in Germany, 10, 11; study on heating of, by insects, 548; effects of percussion on insects in, 632; methods for detection of infestation of, 571; structure and use of fossae for, in Malta, 177, 178; efficacy of electrical radiation against pests in, 374; insecticide treatments of, 178, 195, 483, 558; question of phosphine residues in, 483.

Ceresan 100 (and 200) (mixtures of ethylmercury 2,3-dihydroxypropyl mercaptide and ethylmercury acetate), effects of treatment of cotton seed with phorate and, on incidence of *Pythium*, 398.

Cerococcus catenarius, on coffee in Brazil, 558; systemic insecticides against, 558. Cerodontha denticornis, on grasses in New Zealand, 231.

Ceron N-4S, 272.

Ceroplastes, in China, 362; on Citrus and tea, 362; parasites of, 362.

Ceroplastes destructor, new parasite of, in Natal, 491; on Citrus in New South Wales, 35; dispersal of, 35.

Ceroplastes floridensis japonicus, on Citrus and tea, 362; question of introduction of parasites of, into U.S.S.R. from China, 362.

Ceroplastes rubens, on litchi in Hawaii, 463.

Ceroplastes rusci, characteristics and

world distribution of, 507.

Ceroplastes sinensis, in China, 362; on Leptospermum in New Zealand, 588; question of introduction of parasites of, into U.S.S.R., 362; on Citrus and tea, 362.

Cerosipha forbesi, in British Columbia, 692; on strawberry in Holland, 192; strawberry viruses not transmitted by,

692.

Cetema, species of, on grasses in England,

Ceutorhynchus, on cruciferous oil-seed crops in France, 373; natural enemies

of, 373.

Ceutorhynchus assimilis, on mustard in Britain, 318, 373; on rape in Germany, 440; susceptibility of varieties of mustard to, 318; climatic factors in relation to oviposition and development of, 440; experiments on factors in relation to diapause in, 205; sprays against, 373, 374.

Ceutorhynchus pleurostigma (on crucifers), in Austria, 184; in Holland, 673; bionomics of strains of, 673; factors affecting development and feeding activity of, 205; races of, 205; characteristics and world distribution of, 507; inserticides account 184, 673

insecticides against, 184, 673. Ceutorhynchus quadridens, on radish in

Germany, 303.

Ceylon, Melanagromyza phaseoli on beans in, 484; Nephantis serinopa on coconut in, 169; pests of tea in, 448, 449, 553, 554; Myzus persicae transmitting new virus disease of tobacco in, 81; natural enemies of pests in, 553, 554; virus disease of Natada nararia in, 448.

Chaconin, experiments with, on resistance of Solanum to Leptinotarsa decemli-

neata, 318.

Chad Republic, Diparopsis watersi on

cotton in, 313.

Chaetanaphothrips orchidii, cause of red rust of bananas in Honduras, 462; on banana in Queensland, 503; insecticides against, 463, 503.

Chaetocnema, characteristics and world distribution of species of, 507.

Chaetocnema aridula, bionomics of, on cereals and grasses in U.S.S.R., 243, 244; parasite of, 244; dusts against, 244.

Chaetocnema confinis, on sweet potato in Louisiana, 131.

Chaetocnema hortensis, bionomics of, on cereals and grasses in U.S.S.R.,

243, 244; parasite of, 244; dusts against 244.

Chaetocnema pulicaria, on maize in Maryland, 325; soil treatments against 325.

Chaetocnema tibialis, on beet in Yugo-slavia, 181.

Chaetococcus bambusae, new parasite of, in Brazil, 491.

chalcites, Plusia.

Chalcoides (see Crepidodera).

Chalia, on cacao in Dutch New Guinea, 76.

chalybii, Melittobia.

Charcoal (see Carbon).

Chard (Sea-kale Beet), new Agromyzid on, in Queensland, 295; infected by yellows virus of beet, 17.

Chauliops fallax, bionomics of, on soy bean in China, 593; sprays against,

593.

Cheese, *Piophila casei* infesting premises for storage of, in Switzerland, 100; development of *Necrobia rufipes* on, 436.

Cheiloneurus, species of, as hyperparasites of Planococcus citri in Israel, 614.

Cheimatobia (see Operophtera).

Chelonella (see Chelonus).

Chelonus, parasitising Nacoleia octasema in Indonesia, 354; parasitising Griselda radicana in Quebec, 157; parasitising Tortricids in U.S.S.R., 112.

Chelonus annulipes, parasitising overwintered larvae of Ostrinia nubilalis

in U.S.S.R., 246.

Chelonus heliopae, egg-parasite of Scrobipalpa heliopa in India, 236.

Chelonus striatigenas, parasitising Nacoleia octasema in Indonesia, 354; introduced into Fiji, 354.

Chelonus sulcatus, parasitising Carposina

sasakii in U.S.S.R., 112.

Chenopodium album, infected by beetyellows virus, 17. Chenopodium murale, infected by beet-

yellows virus, 17.

Chermes, on conifers in Ukraine, 112; taxonomy and distribution of European species of, 185; review of predators of, 185; sampling methods for, 185, 186.

Chermes abietis (on spruce), in Quebec,

157; in Ukraine, 112.

Chermes laricis, on conifers in Ukraine, 112.

Chermes merkeri, in Germany and Sweden, 185; on silver fir, 185; factors affecting populations of, 186; predators of, 186; sampling methods for, 185; status of, 185, 186; misidentified as C. piceae, 185, 186.

Chermes nordmannianae, on Abies alba in Denmark, 674; insecticides against, 674.

Chermes nordmannianae schneideri, physiological effects of feeding by, on silver fir, 45.

Chermes nuesslini (see C. nordmannianae). Chermes pectinatae, on conifers in Ukraine, 112.

Chermes piceae (on Abies), in Canada, 185, 394, 456, 522; in Holland, 185; in Switzerland, 186; in U.S.A., 185, 209, 513; bionomics of, 209, 456, 513; factors affecting populations of, 186; natural enemies of, 25, 394, 456, 522; biological and silvicultural control of, 456, 457; forms of, 185; C. merkeri misidentified as, 185, 186.

Chermes piceae typica, physiological effects of feeding by, on silver fir, 45. Chermes pini, on pines in Ukraine, 112. Chermes strobi, on pines in Ukraine, 112. Chermes tardoides, on conifers in Ukraine,

112.

Chermes tardus, on spruce in Ukraine, 112.

Chermes viridis, on conifers in Ukraine, 112.

Cherry, Neurotoma nemoralis on, in Bulgaria, 540; pests of, in France, 84; Rhagoletis cerasi on, in Germany, 679, 680; R. cerasi on, in Italy, 137, 311, 535; pests of, in Ontario, 163, 529; pests of, in Poland, 188, 676; Lyonetia clerkella on, in Switzerland, 663; Tetranychus viennensis on, in Turkey, 177; pests of, in U.S.S.R., 15, 242; factors affecting attraction of Popillia japonica to, in Virginia, 577; Scaphytopius acutus as vector of virus disease of, 23; varietal susceptibility of, to R. cerasi, 188; forecasting of infestation of, 84; systemic treatment of, with schradan, 188; insecticide residues on, 311, 680.

Cherry, Wild (see *Prunus*).

Cherry Plum (see Plum, Cherry).

Chestnut, Curculio elephas on, in Bulgaria, 60; pests of, in France, 84, 435; pericarp structure in relation to infestation of, by C. elephas, 60.

Cheumatopsyche campyla, nuisance value

of, in Iowa, 262.

Cheyletus, predacious on Sitotroga cerealella in cultures, 376; acaricides

against, 376.

Cheyletus eruditus, in stored grain in Britain, 91, 549; in stored seed of flax and hemp in Poland, 364; predacious on other mites, 91, 364, 549; environmental requirements of, 91.

Chick-peas (see Cicer arietinum).

Chickweed (see *Stellaria media*). Chicory (see *Cichorium intybus*).

Chile, Brevipalpus chilensis on vines in, 495; Acarapis woodi infesting honey bees in, 296; predacious mite in, 125. chilensis, Brevipalpus.

Chilo phaeosema, stem borer of cereals

in Nigeria identified as, 292.

Chilo suppressalis, in Japan, 451; in Malaya, 78; on rice, 78, 451; resistance to attack by, in Oryza ridleyi, 78; alimentary metabolism in, 495; effects of larval density on stages of, 451; parasites of, 451; characteristics and world distribution of, 507.

Chilo zonellus, on maize and sorghum in Kenya, 179; recovery of sorghum from attack by, 179; characteristics and world distribution of, 507; DDT

sprays against, 179.

Chilocorus kuwanae, predacious on Quadraspidiotus perniciosus, 170; introduced into Pakistan, 170.

Chilotraea, bionomics of species of, on

rice in Israel, 350.

Chilotraea argyrolepia, on maize and sorghum in Kenya, 179; stem borer of cereals in Nigeria identified as, 292. Chilotraea bandra, 350.

Chilotraea infuscatella, on sugar-cane in India, 297, 553; measures against,

297, 553.

China, Citrus pests in, 362, 491; Aonidiella orientalis on papaya in, 491; pests of other fruit trees in, 501, 502, 591, 592, 625; cereal pests in, 502, 594, 624, 625; pests of rice in, 501, 625; pests of sugar-cane in, 623; Pectinophora gossypiella on cotton in, 502; Ceroplastes spp. on tea in, 362; pests of soy bean in, 593, 594; forest pests in, 501, 593, 623, 624; Biston marginata on Camellia oleifera in, 592; Abraxas miranda on Euonymus in, 503; Aonidiella aurantii in, 21; foci and forms of Locusta migratoria in, 13; Sitophilus granarius in, 693; Tribolium in, 583; Dicranura erminea menciana damaging cable in, 693; beneficial insects in, 362, 491, 591, 592, 593, 623, 624, 625, 664; identity of parasites of Unaspis yanonensis in, 491; key to pupae of Lepidopterous pests in, 501.

China Aster (see Callistephus chinensis). chinensis, Callosobruchus (Bruchus).

Chionaspis, new parasite of, in Natal, 491.

Chipman 6200 (see Amiton Toluene-p-sulphonate).

Chironomus oryzae, on rice in China, 625; seed treatments against, 625.

Chirothrips pallidicornis, effects of, on seed yields of Dactylis glomerata in

New Zealand, 505.

Chlorbenside, against Chermes nordman-674: against Tetranychids. nianae. 242, 444, 678; toxicity of, to winter eggs of Panonychus ulmi, 685; resistance to, in strains of P. ulmi, 672; in dips, 674; in sprays, 242, 444, 674, 678; and BHC, 674; and DDT, 674; and malathion, 242, 678.

Chlordane, against ants, 419; toxicity of, to aphids, 498; against Curculionids, 109, 114, 314, 317, 609, 673; tests and uses of, against other Coleoptera, 173, 314, 371, 497, 526, 587, 647; against Hylemya spp., 40, 226, 254, 608, 673; against other Diptera, 288, diverse effects of, against Lepidoptera, 277, 553; toxicity of, to Orthoptera, 173, 498; against sawflies, 544; against termites, 77, 180, 553, 577, 627; toxicity of, to Tetranychids, 419, 628; effects of, on beneficial insects, 40; toxicity of, to honey bees, 640; laboratory tests of toxicity of, to insects, 202, 203; effect of temperature on toxicity of 202, 203; in aerosols, 419; in dusts, 109, 114, 314, 553, 587, 673; in granules, 641; in sprays, 419, 544, 641; seed treatment with, 226, 484, 608; soil treatment with, 40, 180, 254, 288, 371, 526, 553, 570, 577, 609; persistence of, in soils, 77, 83, 527; treatment of rubber roots with, 77: nest treatment with, 627: treatment of selfing bags with, 277; treatment of textiles with, 587; treatment of timber with, 647; toxicity of, to wheat, 627; residues of, 91, 339, 570, 641; absorption of, by beeswax, 640; question of metabolism of, 64; α isomer of, 202, 203; and DDT, 419; and o-dichlorobenzene, 419; with piperonyl butoxide, 587; and pyrethrum, 587.

Chlorfenson, against Cydia pomonella, 21; against *Eriophyes pyri*, 191; against Tetranychids, 159, 183, 217, 440, 633, 677, 678; development of resistance to, in strains of Panonychus citri, 509, 510; effects of, on orchard pests and their natural enemies, 36; in dusts, 217; in sprays, 36, 159, 183, 191, 217, 440, 633, 677; and Aramite, 183, 633; and DDT, 21; and ferrous

sulphate, 36.

Chloridea (see Heliothis).

Chlorine, Radioactive, insecticides labelled with, 321, 472.

Chloriona cubana, on grasses and rice in Mexico, 518; new parasite of, 518; experiments on transmission of hoja blanca virus by, 518, 689, 690.

Chloriona orizicola (on rice), in Mexico, 518; in Puerto Rico, 257; in U.S.A., 132, 405, 508; experiments with other food-plants and, 132; as vector of hoja blanca virus, 132, 257, 405, 508, 518; new parasite of, 518; question of use of systemic insecticides against, 405.

Chloris gayana, Eulepida mashona infest-

ing, in S. Rhodesia, 38.

Chlorita (see Empoasca).

Chlorobenzilate (in sprays), Phyllocoptruta oleivora, 409, 410, 645; against Tetranychids, 28, 108, 288, 289, 684; susceptibility or resistance to, in strains of Tetranychus spp., 194, 512; against other mites, 439, 626; toxicity of, to Cydia pomonella, 21; Chrysomphalus ficus favoured by, 645; effects of, on beneficial arthropods, 36, 439; evaluation of spray programmes of, 376; effects of, on yields of oranges, 645; toxicity of DDT decreased by, 21; and fungicides, 108; and limesulphur, 36.

Chlorocide (see Chlorbenside).

P-Chloro-2,4-dioxa-5-methyl-P-thiono-3phosphabicyclo[4.4.0]decane, in sprays against Therioaphis maculata, 468.

2-(p-tert.-Butylphenoxy)-2-Chloroethyl 1-methylethyl Sulphite, against Cydia pomonella, 21; against Tetranychids, 28, 183, 217, 264, 324, 477, 517, 628, 633; susceptibility or resistance to, in strains of Tetranychids, 510, 512: against mites in insect cultures, 575; in dusts, 217; in sprays, 28, 183, 217, 264, 265, 324, 477, 517, 633; and BHC 264; and chlorfenson, 183, 633; and DDT, 21, 264; and fungicides, 264, 265, 324; and Tedion, 477, 633.

2-Chloromethyl 2,2-Dichlorovinyl Methyl Phosphate, in tests against Lymantria

dispar, 514.

Chloronaphthalene, treatments of timber with preparations containing, against Hylotrupes bajulus, 254, 647.

Chlorophan (see Dimethyl 1-Methoxy-2,2-dichlorovinyl Phosphate).

m-Chlorophenyl Methylcarbamate, toxicity of, to Musca domestica, 634.

p-Chlorophenyl Phenyl Sulphone, against Tetranychus telarius, 217; in dusts, 217; in sprays, 217.

Chlorophyll, growth of locusts unaffected by, 5; metabolism of, in Tetranychus

telarius, 108.

Chloropicrin (as fumigant), toxicity of. to Coleoptera, 267; factors affecting toxicity of, to Tribolium confusum, 274;

against pests of stored products, 347; treatment of nests with, against Vespula germanica, 422; determination

of, in air, 496.

6-Chloropiperonyl Chrysanthemate, against Pectinophora gossypiella, 409; ineffective against Heliothis zea, 409; experiments with synergists and, against Alphitobius laevigatus, 687; toxicity of, to other insects, 2; in dusts, 2; in sprays, 409; treatment of stored wheat with, 2.

Chlorops hordei (see Meromyza saltatrix). Chlorops oryzae, on rice in Japan, 78; effects of rice growth or variety on

larval development of, 78.

Chlorops pumilionis, on cereals in U.S.S.R., 113; measures against, 114; characteristics and world distribution of, 507.

Chloropsalta viridissima, on vines in Syria, 608; soil fumigants against, 609.

Chlorthion, against aphids, 59, 608; toxicity of, to Eurygaster, 382; against Trypetids, 98, 268; tests of, against pests of stored products, 213; in baitsprays, 98; in sprays, 59, 213, 608; antagonistic action of sesamex with, 63.

choaspes, Asolcus (Microphanurus) (see

A. vassilievi).

Chokecherry (see *Prunus virginiana*). Cholesterol, in diets for insects, **5**, **214**, **404**, **626**.

Choline, action of, on growth of larvae of *Corcyra cephalonica*, **41**; in diet for

Anthonomus grandis, 404.

Cholinesterases, action of chlorinated hydrocarbons on, 62; action of phosphorus compounds on, 62, 273, 385, 395, 396, 407, 424, 475, 512, 513, 604; bioassay of mevinphos in presence of inhibitors of, 397; action of carbamate compounds on, 634, 635; factors affecting inhibition of, 62, 63, 64, 634, 635; effects of insecticide-induced neuroactive substance on activity of, 63; detection of insecticide residues in Gleditsia by inhibition of, 273.

Choristoneura fumiferana (on conifers), in Canada, 25, 121, 122, 130, 168, 208, 391, 455, 457; in U.S.A., 24, 121, 122, 149, 169, 208, 337, 411, 419, 461; climatic factors in relation to breeding cycles of, 208; factors affecting outbreaks of, 25; overwintering and diapause in, 130, 149; parasites of, 24, 337, 391, 457; vertical distribution of parasites of, 24; review of parasites of, in relation to those of other Tortricids, 169; estimation of egg numbers of, 419; aircraft application of sprays

against, 121, 411, 455; other measures against, 25, 26, 461.

Choristoneura murinana (on Abies), in Czechoslovakia, 602, 603, 683; in France and Germany, 168, 169; parasites of, 603, 683; review of parasites of, in relation to those of other Tortricids, 169.

Choristoneura pinus (on conifers), in Canada, 474; in U.S.A., 212, 467, 474; feeding and oviposition habits of, 467; effects of tree location on degree of

parasitism of, 212.

Chorizagrotis auxiliaris, in Alberta, 640; in U.S.A., 214, 640; on cereals, 214; factors affecting damage to mustard by, 640; food-plant preferences of, 214; factors influencing oviposition by, 214.

Chortophaga viridifasciata, nematode infesting, in Missouri, 576.

Chortophila (see Hylemya). Chou Moellier (see Kale).

Chremylus elaphus, parasitising Rhyacionia buoliana in Argentina, 144.

Christolia, parasitising Oiketicus kirbyi in Costa Rica, 638.

Chromaphis juglandicola, in California and France, 151; on walnut, 151; natural enemies of, 151.

Chromaphon, 497.

Chromatography, separation of constituents of insecticide residues and mixtures by, 140, 397, 562. (See also Gas and Paper Chromatography.)

Chromium Acetate, as constituent of

Celcure (q.v.), 485.

Chromobacterium prodigiosum, experiments with, against Eurygaster integriceps, 115.

Chrotogonus, in India, 174; effects of temperature on susceptibility of, to insecticides, 174.

Chrysanthemum, pests of, in Oklahoma, 339; Tetranychus multisetis feeding on, 67; influence of, on susceptibility of T. telarius to methyl-parathion, 548; as reservoir of plant viruses, 271; study on systemic activity and residues of phorate in, 395.

Chrysanthemum carinatum, transmission of viruses to, by Aphrodes bicinctus, 304.

Chrysanthemum cinerariaefolium, as source of pyrethrum (q,v), 140.

chrysocephalus, Psylliodes.

Chrysocharis orchestis, parasitic on Leucoptera scitella in Italy, 136.

Chrysocraspeda olearia, on Eugenia cumini in India, 88.

Chrysomela decemlineata (see Leptinotarsa).

chrysomelina, Epilachna.

Chrysomphalus, new parasite of species of, on Citrus in Brazil, 491.

Chrysomphalus aonidum (see C. ficus).

Chrysomphalus bifasciculatus, on greenhouse plants in Oklahoma, 339.

Chrysomphalus corticosus (see Melanas-

pis).

Chrysomphalus ficus (on Citrus), Brazil, 120, 645; bionomics of, in Israel, 611; question of parasitism of, by Aphytis lingnanensis, 21, 22; sprays against, 120; effects of sprays against Eriophyid on populations of, 645.

Chrysopa, predacious on cotton pests in Arizona, 639; predacious on Psylla pyri in Poland, 365.

Chrysopa lanata, predacious on stages of Eulia loxonephes in Argentina, 144. Chrysopa perla (predacious on aphids), in Bulgaria, 59, 537; in Yugoslavia, 240.

chrysorrhoea, Euproctis.

chrysostictos, Nythobia (Horogenes).

Cicadella viridis (see Tettigella).

Cicadulina arachidis, question of transmission of groundnut virus by, in Gambia, 298.

Cicadulina bipunctella, on maize and rice

in Philippines, 96.

Cicadulina similis, question of transmission of groundnut virus by, in in Gambia, 298. cicatricosa, Glypta.

Cicer arietinum, Diptera on, in Turkey, 351: Bruchidius incarnatus on, 249.

cicerina, Liriomyza.

Cichorium intybus, Phytomyza lateralis on, in Holland, 673; as reservoir of stolbur virus in Yugoslavia, 240; question of dimethoate residues in, 674.

Cienfuegosia affinis, 518.

Cienfuegosia sulphurea, cotton pests developing on, in Texas, 518.

Cigarettes, Lasioderma serricorne in, 57, 58; determination of insecticide residues in smoke of, 228.

ciliatus, Dacus,

cilicrura, Hylemya,

cincticeps, Nephotettix.

cinctipennis, Closterocerus.

cinctipes, Exetastes.

cinctus, Cephus; Emphytus.

cinerascens, Cnephalia (Isomera).

cinnabarinus, Tetranychus.

cinnamomeus, Aradus.

Circulifer opacipennis, map of distribution of, 1.

Circulifer tenellus, 636; in U.S.A., 408, 563, 630; on beet, 630; transmitting virus disease of melons, 408; map of distribution of, 1; parasites of, 630; DDT against, 563.

circumflexus, Myzus.

Cirrospilus flavicinctus, parasitising Bucculatrix ainsliella in Michigan, 327. cistellata, Apsvlla.

citrella, Phyllocnistis.

citri, Diaphorina; Panonychus; Planococcus (Pseudococcus); Scirtothrips.

citricidus, Toxoptera (Aphis). citriculus, Pseudococcus.

citrifolii, Calacarus. citrina, Aonidiella.

citrinus, Aspidiotiphagus.

Citron, Tetranychus multisetis feeding on,

Citrus, pests and tristeza disease of, in Aden, 80; Eriophyids on, in S. Africa, 83; pests of, in Brazil, 120, 491, 556, 558, 644, 645; Aonidiella spp. on, in Burma, 491: Coccids on, in China, 362, 491; Rhynchocoris humeralis on, in Formosa, 591; pests of, in Gaza Strip, 97; pests of, in India, 80, 173, 491; pests of, in Israel, 22, 611, 612, 614, 684; pests of, in Italy, 181, 299; pests of, in Mexico, 22, 280; Coccids on, in Morocco, 382; Ceroplastes destructor on, in New South Wales, 35; Coccus hesperidum on, in Nigeria, 531: Phyllocoptruta oleivora on, in Peru, 560; Aonidiella aurantii on, in Siam, 491; Brevipalpus phoenicis on, in Spain, 299; pests of, in Turkey, 82; pests of, in U.S.S.R., 242, 362; pests of, in U.S.A., 22, 147, 221, 267, 378, 409, 465, 509, 519, 552, 561, 564, 688; new Tortricid feeding on, 230; Tetranychus multisetis feeding on, 67; relation of ants to populations of Coccids on, 519; aphids and tristeza disease of, 80, 97; experiments with Metcalfa pruinosa as vector of tristeza virus of, 465; mites in relation to leprosis of, 556; ecological survey of pests of, 552; biological control of pests of, 199; effects of acaricides on yields of, 645; study of distribution of oil-emulsion deposits on, 221; insecticide residues in fruits of, 561, 574.

Cladius pectinicornis, on strawberry, rose and other plants in U.S.S.R., 361; bionomics of, 361; insecticides against, 362.

Clastoptera, BHC against, on jute in British Guiana, 28.

Clastoptera undulata, bionomics of, on Casuarina and other plants in Bermuda.

230; parasite of, 230; sprays against, 230.

clavicornis, Rhopalophorus.

clavifer, Helopeltis.

Clays, volatility of insecticides in, 416; as carriers for insecticide dusts, 50. (See also Attaclay and Soil.)

Cleonus mendicus, on beet in Italy, 533;

sprays against, 310, 533.

Cleonus punctiventris (on beet), in Bulgaria, 541; in U.S.S.R., 609; Beauveria bassiana against, 541; insecticides against, 541, 609; characteristics and world distribution of, 507.

Clepsis peritana (see Ptycholoma).

clerkella, Lyonetia.

clitellatus, Pachynematus,

Clivina impressifrons, on maize in Illinois, 420; insecticides against, 420.

Closterocerus, egg-parasite of Caliroa limacina in Quebec, 26.

Closterocerus cinctipennis, relation of oviposition habits of Neodiprion swainei to egg-parasitism by, in Quebec, 459.

Clostridium brevifaciens, 389; infecting Malacosoma pluviale in British Columbia, 342; in sprays against M. americanum, 343.

Clostridium malacosomae, infecting Malacosoma pluviale in British Columbia, 342; in sprays against M. americanum, 343.

Clothes Moth (see *Tineola bisselliella*).

Clover, Cicadellids on, in Britain, 686; Aphis craccivora on, in Bulgaria, 59; Aphrodes bicinctus on, in Canada, 686; A. bicinctus on, in Hungary, 307; pests of, in U.S.S.R., 114, 686; phyllody virus of, 686; in grass mixture, 430; BHC residues on, 323, 324, question of fumigation of stored seed of, against Bruchidius perparvulus, 183. (See Trifolium spp.)

Clover, Sweet (see Melilotus).

Clysiana (Clysia) ambiguella (on vines), in Bulgaria, 60, 538; in France, 8, 9, 434, 494, 667, 668; in Germany, 108; bionomics of, 538; Bacillus thuringiensis ineffective against, 9; insecticides against, 8,9 108, 434, 494, 538, 667, 668

Clysiella (see Clysiana).

CMC Cellulose Gum 7L, 272.

Cnemarachis bruneri (see Lachnosterna). Cnemodon elongatus, predacious Eriosoma lanigerum in Nova Scotia,

388. Cneorrhinus plagiatus, on pines in U.S.S.R., 116.

parasitising cinerascens, Apamea anceps in U.S.S.R., 363.

coagulata, Homalodisca.

coarctata, Hylemya (Leptohylemyia).

Cobalt, Radioactive, effects on insects of γ-radiation from, 568, 607; use of, in determination of predators of *Rhya*cionia buoliana, 393.

Coccidencyrtus, keys to females of species

of, 490; redefinition of, 490.

Coccidencyrtus infuscatus, parasitising Quadraspidiotus juglansregiae California, 491.

Coccinella novemnotata, predacious on

Aphis pomi in Wisconsin, 415.

Coccinella septempunctata, in Bulgaria, 59, 537; in Poland, 365; in U.S.S.R., 243; predacious on aphids, 59, 243, 319, 537; predacious on Psylla pyri, 365; experiments on prevention and induction of diapause in adults of, 319.

Coccinella transversoguttata, predacious on aphids in New York, 401.
Coccophagus basalis, bionomics of, parasitising Coccids, 150, 151; introduced into California from Brazil, 150.

parasitising Coccophagus eritreaensis, Coccus mangiferae in Israel, 613.

Coccophagus inaron, parasitising Eulecanium coryli in France, 683.

Coccophagus lycimnia, parasitising Eulecanium corni in France, 683.

Coccophagus pulchellus, parasitising Eulecanium coryli in France, 683.

Coccophagus quaestor, reared Toumeyella numismatica, 529; in Manitoba and Mexico, 529.

silvestrii, parasitising Coccophagus Eulecanium coryli in France, 683.

Coccophagus trifasciatus (see Lounsburyia). Coccus hesperidum, in Nigeria, 531; in U.S.S.R., 597; in U.S.A., 150, 151, 339, 491, 519; on Citrus, 519, 531; on greenhouse plants, 339; parasites of, 150, 151, 491, 597; ants associated with, 519, 531.

Coccus mangiferae, bionomics and natural enemies of, on mango in Israel, 613; other food-plants of, 613; sooty mould associated with, 613.

Coccus pseudomagnoliarum, Aphycus stanleyi parasitising, in California, 491.

Coccus viridis, sprays against, on coffee in Brazil, 557.

cochleariae, Phaedon.

cockerelli, Paratrioza.

Cockroach, German (see Blattella germanica).

Cockroaches, infesting insect cultures in California, 339; metabolism of insecticides in, 64, 513; experiments on action of insecticides on, 62; physiological activity of body fluids from DDT-treated crayfish in, 63; bait for, 339. (See also *Blattella*, *Leucophaea* and *Periplaneta*.)

Cocksfoot (see Dactylis glomerata).

Coconut, pests of, in South-East Asia and Pacific area, 52, 81, 143, 166, 167, 235, 293; Homalinotus coriaceus on, in Brazil, 559; Nephantis serinopa on, in Ceylon and India, 169; Rhynchophorus palmarum as vector of nematode causing red ring disease of, in Trinidad, 575; pests of, in Zanzibar, 29; review of pests and diseases of, 693; biological control of pests of, 199.

cocophaga, Amblypelta.

Codiosoma spadix, infesting building timber in U.S.S.R., 112.

Codling Moth (see Cydia pomonella).

Coelinius (Coelinidea), species of, parasitising Meromyza saltatrix in Japan, 97. Coeloides bostrichorum, parasitising Ips typographus in Czechoslovakia, 142; method of rearing, 142.

Coeloides melanotus, parasitising Hylesinus fraxini in Czechoslovakia, 187;

parasites of, 188.

Coeloides filiformis, parasitising Hylesinus fraxini in Czechoslovakia, 187; parasites of. 188.

Coelostomidia wairoensis, on Leptospermum in New Zealand, 588.

Coelostomidia zealandica, on Leptospermum in New Zealand, 588.

Coenorhinus (see Rhynchites).

coerulescens, Baris.

Coffee arabica, 76, 448, 490. (See also Coffee.)

Coffea canephora, pests of, in Dutch New Guinea, 76.

coffeae, Diarthrothrips; Oligonychus; Saissetia; Zeuzera.

coffearia, Homona.

Coffee, pests of, in Brazil, 120, 555, 557, 558, 643; pests of, in India, 37, 448; pests of, in Ivory Coast, 7, 651; Xyleborus morigerus on, in Java, 625; atlas of pests and diseases of, in Kenya, 490; pests of, in New Guinea, 76, 234; pests of, in Tanganyika, 652, 653; Antestiopsis spp. on, in Uganda, 490; Xyleborus spp. on, in Vietnam, 618; Colletotrichum coffeanum associated with X. morstatti on, 651; review of pests of, 196; sampling method for arthropods on, 490; biological control of pests of, 199; question of value of BHC sprays against pests of, 7, 8; (stored), fumigation of, against Araecerus fasciculatus, 558.

coffeicola, Pediobius (Pleurotropis),

cognata, Aelia; Cyrtozemia. cognatellus, Hyponomeuta. cognatus, Epicaerus.

coheni, Aphytis.

Cola, new weevils on, in Cameroun, Ivory Coast and Guinea, 355.

Colaeus monedula, destroying pupae of Dendrolimus superans in Siberia, 13.

Colaphellus sophiae, larval feeding of, not influenced by photoperiod, 248.

Colaspidema atrum, characteristics and world distribution of, 507.

Colaspinella grandis, dusts against, on cotton in Turkey, 237.

Colaspis flavida, insecticides against, on

rice in Arkansas, 421.

Cold-hardiness, in Coleoptera in stored grain, 344; in Lepidoptera, 4, 424, 447, 502, 538; in *Prospaltella perniciosi*, 681; of eggs of *Panonychus ulmi*, 528.

Cole, pests of, in U.S.A., 222, 401.

colfaxiana, Barbara.

Colias electo, bionomics of, on lucerne in S. Rhodesia, 48.

Colias eurytheme, trans-ovarian transmission of polyhedrosis virus of, 496; method of rearing individuals of, 281.

Colias lesbia, characteristics and world distribution of, 507.

colibri, Athalia (see A, rosae).

Collards, pests of, in U.S.A., 68, 404; ecological studies of invertebrate fauna of, 68.

collaris, Exora.

Colletotrichum coffeanum, Xyleborus morstatti associated with, on coffee in Ivory Coast, 651; mixed sprays against 651.

Collops, predacious on cotton pests in

Arizona, 639.

Collyria calcitrator, in Britain, 161; in Sweden and U.S.S.R., 160, 161; parasitising Cephus pygmaeus, 161; introduced into Canada against C. cinctus, 161.

Colombia, Blissus leucopterus in, 98; Coleoptera in stored beans and maize

in, 560.

Colorado, yellows virus of beet in, 17; Dendroctonus engelmanni on spruce in, 150.

Colorado Beetle (see *Leptinotarsa decem-lineata*).

Colours, reaction of Sitona cylindricollis to, 319.

columbicus, Nasutitermes.

Colutea arborescens, Eurytoma coluteae infesting seeds of, in Hungary, 306. coluteae, Eurytoma.

comes, Erythroneura (Typhlocyba).

commixtalis, Loxostege.

commodus, Acheta.

Comperiella bifasciata. new parasite reared on, in association with Aonidiella aurantii in Burma and Siam, 491; oviposition of Physcus spp. on, 266.

Comperiella lemniscata, sp.n., parasitising Aonidiella orientalis in China,

India and Pakistan, 491.

Compsilura concinnata (parasitising Lymantriids), introduced into British Columbia, 126; in Yugoslavia, 370.

comptana, Áncylis. comstocki, Pseudococcus.

comstockii, Ephialtes (Calliephialtes).

concinnata, Compsilura.

concinnus, Ips. concolor, Opius.

Concrete, treatment of, against termites,

condignella, Evagora (Recurvaria).

confinis, Chaetocnema.

Confused Flour Beetle (see Tribolium confusum).

confusum, Tribolium.

confusus, Anthocoris; Ips; Melanoplus; Physcus: Tarsonemoides.

congelatella, Exapate.

congerens, Formica (see F. nigricans).

Congo (Leopoldville), Aphis craccivora and virus disease of groundnuts in, 609, 610; Rivellia damaging root nodules of groundnut in, 609; predacious ant in, 28; ants associated with A. craccivora in, 610; fungus infesting A. craccivora in, 610.

Congress of Crop Protection, Proceedings of Fourth International, 444, 445, 608,

609, 681–684.

Congress of Entomology, Proceedings of Eleventh International, 91.

Coniesta ignefusalis, bionomics and lighttrap catches of, on cereals in Nigeria, 292, 293; parasite of, 292; measures against, 293.

coniperda, Conophthorus.

conjugella, Argyresthia.

Connecticut, Rhagoletis pomonella on apple in, 400; effects of sprays on arthropod fauna of apple orchards in, 694; pests of forest trees in, 271, 400, 407, 641; wireworms in, 582; parasites in, 271, 400, 407; diseases of Anisota senatoria in, 407; insecticide residues in earthworms in, 641.

connexa, Eriopis.

Conoderus falli, in S. Carolina, 575; infestation of cultures of, by mites, 575; method of rearing, 575.

Conophthorus, bionomics of species of, on conifers in Ontario, 260, 261.

Conophthorus coniperda, on white pine, 261.

Conophthorus resinosae, on red pine. 261. Conotrachelus nenuphar (on apple), in Canada, 154, 159; in U.S.A., 216, 403, 417; fecundity and egg viability of strains and crosses of, 216; soil treatment against, 417; killed by controlledatmosphere storage of fruits, 403.

conquisitor, Itoplectis.

consanguinea, Lachnosterna.

consobrina, Érnestia; Vespula rufa. conspicuus, Typhlodromus.

Contarinia medicaginis (on lucerne), in Bulgaria, 60; bionomics and overwintering of, in Czechoslovakia, 187; effect of temperature on embryonic development of, 494; characteristics and world distribution of, 507.

Contarinia nasturtii, in Germany, 139; factors affecting development of, 139.

Contarinia oregonensis (on Douglas fir), in Canada, 385, 386; in U.S.A., 355, 385, 386; bionomics of, 385; parasites of, 386.

Contarinia pisi (on peas), in Britain, 287; in Holland, 193; delayed emergence of, 287, 288; determination of crop losses caused by, 193.

Contarinia sorghicola (on sorghum), investigation on first records of, in African territories, 684, 685; in Kenya, 179; in U.S.A., 421.

Contarinia tritici, on wheat in Britain, 420; susceptibility of varieties of wheat to Mayetiola destructor and, 420.

convergens, Hippodamia. convexifrons, Allotropa,

convexinotatus, Heterotermes.

Convolvulus, development of Agrotis vpsilon on, in Bulgaria, 346.

Convolvulus arvensis, Hyalesthes obsoletus transmitting stolbur virus to potato from, in Yugoslavia, 240.

Copidosoma deceptor, parasitising Evagora milleri in California, 221; parasitising E. starki in Canada, 586; effects of sprays on populations of, 222.

Copidosoma truncatella (see Litomastix). Copper, build-up of, in vineyard soils,

228.

Copper Oxychloride, against Dacus oleae, 438; effects of, on populations of *Panonychus ulmi*, 108; and parathion, 135, 438.

Copper Sulphate, in sprays against Dacus oleae, 47, 309, 438; repellent effect of, on Leptinotarsa decemlineata, 305; and DDT, 438; and parathion, 309, 438; effects of, on efficiency of parathion, 47, 135; Celcure (q.v.) containing, 485.

Copra, insects infesting, 436, 657; insecticide treatments of, 436.

optotermes curvignathus, attacking rubber trees in Malaya, 76, 77; soil Coptotermes treatments against, 77.

Coptotermes niger, attacking trees in British Honduras, 76.

Copturus aguacatae, bionomics and con-

trol of, on avocado in Mexico, 356. Coraebus rubi, on roses in Italy, 299.

Co-Ral (see Coumaphos).

Corchorus (see Jute).

Corcyra cephalonica, in stored groundnuts in Brazil, 558; parasitised by Antrocephalus in India, 314; Apanteles angaleti reared on, 317; sex of, in relation to that of parasite, 314; nutritional requirements of, 41, 626; phosphine fumigation against, 558.

Cordia myxa, Perissopneumon tectonae on, in India, 174.

Cordyceps aphodii, infesting Aphodius tasmaniae in S. Australia, 166.

Cordyceps militaris, Anisota senatoria infested by, in Connecticut, 407.

Cordyceps ravenellii, Lachnosterna spp. infested by, in Canada, 619.

coriaceus, Homalinotus.

coricopa, Agriophara. Corium malvae, 480.

Corn Borer, South-western (see Zeadiatraea grandiosella).

corni, Eulecanium (Lecanium, Parthenolecanium).

corniger, Nasutitermes.

cornutus, Aculus (Vasates); Centrotus.

corpulenta, Anomala.

corticosus, Melanaspis (Chrysomphalus). Cortinellus edodes, 623.

corvli, Eulecanium.

corylifoliella, Lithocolletis,

Corylus, Eulecanium corni on, in U.S.S.R., 13.

Corylus avellana, pests of, in Turkey, 82, 83.

Corythaica cyathicollis, on solanaceous plants in Brazil, 554; sprays against, 554; neotype and status of, 554.

Corythaica monacha, on Sida in Brazil, 554; characters and status of, 554.

Corythaica passiflorae (planaris), synonymy and status of, 554.

Cosmolyce boetica, characteristics world distribution of, 507.

Costa Rica, Oiketicus kirbyi on banana in, 637, 638; insects infesting Heliconia in, 520; Heliconia as natural host of Pseudomonas solanacearum in, 520; natural enemies and fungus disease of O. kirbyi in, 638.

costalimai, Deltinea; Stylopalpia.

costalis, Monellia.

Cothonaspis (see Trybliographa).

Cotton, pests of, in Argentina, 694; pests of, in Brazil, 557, 643, 645, 646; Diparopsis watersi on, in Chad Republic, 313; Pectinophora gossypiella on, in China, 502; pests of, in Egypt, 149, 608; pests of, in Formosa, 590; pests and diseases of, in Greece, 352; Tetranychids on, in Hungary, 307; pests of, in India, 36, 37, 139, 140, 314, 317, 628, 629; pests of, in Israel, 351, 613: P. gossypiella on, in Mexico, 473, 511; pests and diseases of, in Morocco, 490; D. watersi on, in Nigeria, 53; survey of data on Hemiptera on, in Nyasaland, 352; Sphenoptera gossypii on, in W. Pakistan, 176; pests of, in Peru, 559, 560; pests of, in Sudan, 52, 94; pests of, in Turkey, 82, 177, 237, 352, 495, 552; pests of, in Uganda, 652; pests of, in U.S.S.R., 15, 113; pests of, in U.S.A., 16, 147, 218, 263, 269, 322, 323, 330, 334, 338, 403, 408, 409, 410, 412, 413, 417, 464, 466, 467, 473, 477, 478, 480, 508, 511, 515, 517, 567, 568, 573, 574, 576, 629, 638, 639; D. watersi on, in West Aden Protectorate, 486; other arthropods on, 467, 472, 478, 552, 569; forecasting abundance of pests of, 15; factors affecting infestation of, by insects, 52, 53, 403, 464, 466, 467, 567, 568; feeding and oviposition sites of Anthonomus grandis on, 410; effects of infestation by A. grandis on yields of, 574; larvae of P. gossypiella surviving centrifugal forces during ginning of, 479; effects of stalk shredders for, on overwintering larvae of P. gossypiella, 334. 417: effects on Tetranychus telarius of treatment of, with gibberellin, 410; methods for collecting insects from, 225; biological control of pests of, 490; seed treatment of, 218, 224, 263, 277, 352, 467; effects of seed treatment of. on Pythium infection, 398; methods and results of treatment of, with systemic insecticides, 176, 224, 277, 467, 472, 473, 475; effects of insecticides on development of, 95, 218, 220, 277, 409, 567, 578, 632, 645, 653; toxicity of insecticides to, 263, 352, 399, 467, 643, 645; persistence of insecticide residues on, 16; metabolism of fenthion in, 476; in diets for insects, 406, 407, 474. (See also Gossypium.)

Cottonseed, phosphine fumigation of, against Pectinophora gossypiella, 558; effects of fumigants on germination of. 579; in diet for Anthonomus grandis,

404.

Cottonseed Meal, in diets for Pectinophora gossypiella, 465, 474; insecticide residues in, 578.

Cottonseed Oil, in baits for fire ants, 464; in stabilised formulations of dichlorvos, 562; insecticide residues in. 578.

Cottonwood, Black (see Populus trichocarpa).

Coulometry, in analysis of insecticide residues. 91.

Coumaphos, soil treatment with, against Melolonthid, 493; residues and stability of, in milk, 266.

Coumarin, rôle of, in termination of flight by Sitona cylindricollis, 319.

Coumarone Indene Resin, in formulations of DDT, 29.

Cowpea Aphid (see Aphis craccivora).

Cowpeas, Aphis craccivora on, in India, 498; Coleoptera on, in Nigeria, 49, 50; resistance or susceptibility to pests in strains of, 413, 414; experiments with Rivellia damaging root nodules of, 609; (stored), Bruchids infesting, 50.

coxalis, Acroleucus, CPBS (see Fenson).

Crab Apple (see Pyrus prunifolia).

Crabro, predacious on larvae of Neodiprion swainei in Quebec, 73.

craccivora, Aphis.

Cranberry, Sparganothis sulfureana on, in U.S.A., 337.

crassicornis, Actia.

crassifemur, Campoplex (Limnerium).

crassiuscula, Phobocampe.

crataeganus, Archips (Cacoecia).

crataegi, Aporia.

Crataegus (Hawthorn), insects on, in Canada, 26, 585; Lepidoptera on, in U.S.S.R., 112; experiments with Rhopalosiphum padi on, 585.

Cratichneumon fabricator, parasitising Operophtera brumata in U.S.S.R., 190. parasitising w-album, parasitising Cratichneumon Anisota senatoria in Connecticut, 407.

Cratotechus (see Eulophus).

Crayfish (see Orconectes).

parasitic on Cremastus flavoorbitalis, Antigastra catalaunalis in India, 316.

Cremastus interruptor, established in Ontario as parasite of Rhyacionia buoliana, 73, 74.

Crematogaster, relation of, to fungus disease of aphids in Congo, 610.

Creolin, use of BHC-activated, against Steneotarsonemus fragariae, 675.

Creontiades, species of, on cotton in Turkey, 82.

Creontiades pallidus, on cotton in Turkey, 82.

Creosote, Coal-tar, against termites, 180, 181; soil treatment with, 180; other methods of application of, 181; and fuel oil. 181.

Crepidodera aurata, on willows and poplars in Yugoslavia, 181.

Crepidodera plutus, on willows and poplars in Yugoslavia, 181.

cretica, Sesamia.

cribricollis, Otiorhynchus (Brachyrhinus). Crickets, in U.S.A., 567; nomenclature of, 567. (See also Acheta, Gryllus and Scapsipedus.)

Cricula trifenestrata, bionomics of, on

mango in Pakistan, 88.

crinitus, Sitona.

Criocoris saliens, Atractotomus mali misidentified as, 159.

cristata, Bryobia. cristatus, Trichomalus. crocea, Hemichroa.

Crop Protection, Proceedings of Fourth International Congress of, 444, 445, 608, 609, 681-684.

Crotalaria longithyrsa, experiments with Rivellia and, 609.

cruciferae, Phyllotreta. cruegeri, Pericyma.

Cryptoblabes gnidiella, in Israel, 612; in Turkey, 82; on Citrus, 82, 612; development of, on grapes, 612; bionomics of, 612; relation of Coccids to, 612.

Cryptolestes, intercepted in stored grain in Germany, 11.

Cryptolestes ater (see C. spartii).

Cryptolestes capensis (in stored food-stuffs), in S. Africa, 656; in Britain, 657; in flour mills, 658; bionomics of, 657, 658; infected by Mattesia dispora, 658; considered distinct from C. spartii, 656.

Cryptolestes ferrugineus (in stored foodstuffs), in Canada, 155; in Gambia, 178; bionomics of, in flour mills, 658; factors affecting oviposition rate of, 566; effects of larval density on populations of, 657, 658.

Cryptolestes pusillus, in stored rice in Louisiana, 571; malathion against, 571.

Cryptolestes spartii, on broom in Britain, 656; method of rearing, 656; synonymy of, 656; C. capensis considered distinct from, 656.

Cryptolestes turcicus, infesting flour mills in Britain, 657; bionomics of, 658; factors affecting populations of, 657, 658.

Cryptolestes ugandae, factors affecting populations of, 658.

Cryptomyzus galeopsidis (see Capitophorus).

Cryptomyzus ribis (see Capitophorus).

Cryptophlebia leucotreta, reared from pomegranate fruits in S. Africa, 531; characteristics and world distribution of. 507.

Cryptorhynchus lapathi, estimation of populations of, in France, 83; in Hungary, 307; on willows, 83, 307; other food-plants of, 307.

Cryptostemma calendulaceum, Aphodius tasmaniae on, as component of pastures

in S. Australia, 504.

Cryptotermes brevis, in imported timber in Britain, 76, 321; in British Honduras, 76; in Canary Islands, 300; in Trinidad, 321; in building timber, 76; in furniture, 76, 321.

Cryptotermes cynocephalus, in timber and furniture in Java, 296; insecticides

against, 296.

Cryptotermes domesticus, in timber and furniture in Java, 296; insecticides

against, 296.

Cryptotermes dudleyi, in timber and furniture in Java, 296; insecticides against, 296; characteristics and world distribution of, 507.

Cryptotermes havilandi, infesting building timber in W. Africa, 485; bionomics and dispersal of, 485; measures against,

485, 486.

Ctenicera aeripennis destructor, in Saskatchewan, 155; adult activity of, 155, 156; effects of diet on larvae of, 495; experiments on feeding response of larvae of, to rye-seed extracts, 528; traps for, 155, 156.

Ctenicera tarsalis, destroyed by bacterial

toxins, 582.

Ctenochiton, species of, on Leptospermum in New Zealand, 588.

Ctenomorphodes tessulatus, on Eucalyptus in New South Wales, 231; parasite of, 231; effects of bush fires on populations of, 231.

cubana, Chloriona (Sogata).

Cubé (see Rotenone).

Cucumber, Tyrophagus longior on, in British Columbia, 529; Tetranychus desertorum on, in Japan, 452; mosaic virus of, in U.S.A., 271; effects of insecticides on respiration of, 562.

cucumeris, Epitrix.

Cucumis melo (see Melon).

cucurbitae, Dacus.

Culex pipiens fatigans (quinquefasciatus, auct.), use of larvae of, in tests of insecticides, 516, 634.

Culicoides, review on bionomics and control of, 90.

culinaris, Rhyncolus.

Cultural Practices, effects of: on populations of aphids, 605, 606; on Ostrinia nubilalis, 12, 56, 372; on dispersal of hibernated adults of Sphenoptera gossypii, 176; on other insects, 7, 52, 58, 109, 191, 203, 348, 371, 377, 441; on Tetranychid populations, 439, 444, 477; on incidence of pests of tea and their parasites, 553; on stability of invertebrate populations, 68, 69, 90; on biological control, 200; on recovery of infested lucerne, 210; on distribution and persistence of insecticides in soils, 255, 272.

cunea, Hyphantria.

cuneatus, Melanoplus,

cupreoviridis, Earias.

cupressana, Lozotaeniodes (Eulia).

Cupressus, Lozotaeniodes cupressana on, in Italy, 137.

cupreus, Rhynchites.

Curculio elephas, in Bulgaria, 60; in France, 84, 435; in Spain, 660; on chestnut, 60, 84, 435; on Quercus ilex, 660; estimation of populations of, 84; pericarp structure in relation to infestation of chestnut varieties by, 60; sprays against, 435, 436; characteristics and world distribution of, 507.

Curculio, Plum (see Conotrachelus nenuphar).

curculionis, Bathyplectes.

Currant, Dasiops alveofrons reared from, in California, 128; Quadraspidiotus perniciosus on, in France, 434; experiments on transmission of Q. perniciosus to, 664.

Currant, Black, mites on, in Britain, 70, 287, 432, 671; Aegeria tipuliformis on, in Czechoslovakia, 445; Cecidophyopsis ribis on, in Holland, 302; Capitophorus spp. on, in U.S.S.R., 595, 596; Cecidophyopsis ribis and reversion virus of, 70, 302; toxicity of acaricides to, 672.

Currant, Red, Capitophorus sidis on, in U.S.S.R., 595.

curvidens, Ips (Pityokteines).

curvignathus, Coptotermes.

curvinervis, Camarota. curvipennis, Camarota.

Cuscuta, question of transmission of plant viruses by, 17, 71, 441.

cuscutae, Aphis.

custodiens, Anoplolepis.

cyanescens, Pardalaspis.

cyanoideus, Catolaccus.

cyathicollis, Corythaica.

Cyclamen, oviposition and development of Tetranychus telarius on, 547.

Cyclocephala, on maize in Illinois, 420; insecticides against, 420.

Cyclohexanone, as diluent for solution of

pyrethrins, 3, 4.

5-(1-Cyclohexen-1-yl)-1,5-dimethylbarbituric Acid, effects of, in mixture with rotenone against Epilachna varivestis, 279.

Cycloheximide, treatment of flannel with, ineffective against Anthrenocerus australis, 587.

Cydia, on wild rose in U.S.S.R., 113.

Cydia funebrana, on damson in China, 625; in Poland, 189; in U.S.S.R., 112; on plum, 112, 189; on Prunus japonica, 112; bionomics and parasites of, 112; characteristics and world distribution of, 507.

Cydia molesta, in Brazil, 646; in France, 84, 256, 664, 665; in Italy, 662; in Ontario, 163; on peach, 84, 163, 256, 646, 662, 664, 665; on other fruit trees, 163; bionomics of, 665; factors affecting development and overwintering of, 662, 665; parasites of, 665; estimation of populations of, 84, 665; traps for, 84, 665; insecticides against, 256, 646.

Cydia nigricana, on peas in Britain, 431; cost of sprays against, in relation to

crop losses, 431.

Cydia pomonella (Codling Moth), in Australia, 165; in Austria, 247; in Britain, 671; in Canada, 159, 394; in China, 592; in Germany, 381, 681; in Greece, 239; in Poland, 189; in Switzerland, 664; in U.S.A., 18, 20, 276, 466; on pear, 592; on walnut, 276; bio, nomics of, 592; parasites of, 381, 592-681; relation of Paramyelois transitella to, 276; speed of entry into apples by larvae of, 637; bait- and light-trap catches of, 165; experiments on biological control of, 381; insecticides against, 20, 159, 239, 394, 466, 592, 637, 664, 671; effects of insecticides on eggs of, 256; resistance to DDT in strain of, 394; effects of measures against, on other insects, 18, 248.

Cydia pyrivora (on pear), in Austria, 247; in Rumania, 57; bionomics of, 57, 247; varietal susceptibility of pears to, 247; measures against, 57, 248.

Cydia splendana, on chestnut in France, 435; sprays against, 435, 436.

Cydia fletcherana, damaging Douglas fir in Montana, 460; characters of sexes of, 460.

cylindrica, Sphaerophoria. cylindricollis, Sitona.

Cylindrocopturus eatoni, on pines in California, 642.

cylindrus, Platypus.

Cynara scolymus, Otiorhynchus cribricollis on, in California, 509.

cynocephalus, Cryptotermes.

Cynodon dactylon, damaged by Aceria neocynodonis in Arizona, 396; Calli-gypona spp. on, in Israel, 349; Aelia acuminata attracted to, 653; question of relation of, to virus disease of maize, 349.

cynthia, Attacus.

Cyperus, Sphenophorus callosus on, in Georgia, 152.

Cyperus rotundus, Calligypona striatella on, in Israel, 349.

Cypress (see Cupressus).

Cyprex (see n-Dodecylguanidine Acetate). Cyrtoptyx dacicida, parasitic on Dacus oleae in Italy, 308, 438, 532.

Cyrtozemia cognata, on vegetables and flowering plants in India, 693.

cytherea, Nudaurelia.

Cyzenis albicans (see Monochaeta).

Czechoslovakia, Eriosoma lanigerum on apple in, 678; Aegeria tipuliformis on black currant in, 445; Depressaria apiella on caraway in, 301; forest pests in, 142, 187, 196, 246, 283, 600, 601, 602, 603, 683; pests of leguminous forage crops in, 187, 304, 445; virus diseases of clover in, 304; pests of potato in, 236, 247, 681; Melolontha spp. in, 186; other pests in, 236, 305; natural enemies of pests in, 142, 186, 187, 188, 246, 301, 602, 603, 678, 683; Microsporidia infecting insects in, 105, 187, 283; other diseases of insects in 186, 187.

D

2,4-Dichlorophenoxyacetic 2,4-D (see Acid)

dacicida, Cyrtoptyx (Dinarmus).

Dactylis glomerata, pests of, in Canada, 162, 389, 585, 586; stem-boring Diptera on, in England, 487, 488; pests of, in New Zealand, 505, 506; effect of Chirothrips pallidicornis on seed yield of, 505.

dactylopii, Leptomastix.

Dactylosphaera vitifoliae (see Phylloxera). Dacus ciliatus, characteristics and world distribution of, 507.

Dacus cucurbitae (on cucurbits), in Formosa, 590; in Pakistan, 175; characteristics and world distribution of, 507; baits for, 590; insecticides against, 175, 590.

INDEX

Dacus dorsalis, question of climatic factors limiting establishment of, in California, 563; in Hawaii, 563; characteristics and world distribution of, 507.

Dacus oleae (on olive), in France, 435; in Greece, 237, 238; in Italy, 46-48, 135, 308, 309, 438, 532, 694; in Spain, 99; bionomics of, 532; effects of low temperatures on emergence of, 135; natural enemies of, 308, 435, 438, 532; other mortality factors of, 308; factors affecting infestation of olive by, 46, 238; baits for, 99, 439, 533; repellency of bordeaux mixture to, 47; other measures against, 47, 99, 135, 237, 238, 239, 308, 309, 438, 439, 533, 562, 694. Dacus tryoni, characteristics and world

distribution of, 507.

Dacus tsuneonis, characteristics and world

distribution of, **507.** Dadap (see *Erythrina*).

Dahlbominus fuliginosus (fuscipennis), competition between Habrocytus sp. and, parasitising Diprion pini in Yugoslavia, 31.

Dahlia, as reservoir of plant viruses, 271. Dakota, North, *Tetanops myopaeformis* on beet in, 330.

Dakota, South, Ostrinia nubilalis on maize in, 461.

Dalopius pallidus, on potato and turnip in Nova Scotia, 73.

Damping-off (see Pythium).

Damson, Cydia funebrana on, in China, 625.

dannehli, Cydia (Carpocapsa) (see C. pyrivora).

Dasiops alveofrons, sp. n., in Mexico, 128; in U.S.A., 128, 271; on apricot, 128, 271; on currant, Juglans and sunflower, 128; characters and bionomics of, 271.

Dasychira pudibunda, on apple in Finland, 103; characteristics and world distribution of, 507.

Dasyneura affinis, characteristics and world distribution of, 507.

Dasyneura balsamicola, Agevillea abietis distinct from, 252.

Dasyneura leguminicola, on red clover in Ontario, 71; influence of harvesting on populations of, 71.

Dasyneura mali, on apple in Finland, 103. datanae, Winthemia (see W. quadripustulata).

Date Palms, pests of, in Morocco, **54**, **382**. Dates (Stored), Lepidoptera infesting, **54**; fumigation of, **54**.

Datura stramonium, Leptinotarsa decemlineata on, in Bulgaria, 366; experiments with plant viruses and, 636, 691. dayatchi, Salicicola.

DD-136 (nematode), introduced into Ontario, 162, 207; insects infected by associated bacteria and, 162, 163, 207, 208, 580; field use of, 162, 163, 207.

D-D Mixture (1,2-dichloropropane and 1,3-dichloropropene), as soil fumigant: against *Chloropsalta viridissima*, 609; against *Lachnosterna consanguinea*, 171.

DDD, against Aphis gossypii, 559; against Quadraspidiotus perniciosus, 168; toxicity of, to Aulacophora foveicollis, 173; against Lepidoptera, 20, 46, 144, 267, 514, 557, 558, 559; against Rhagoletis pomonella, 268; against Thysanoptera, 645; against termites, 577; against Hemitarsonemus latus, 645; effects of, on beneficial arthropods, 159; in dusts, 514, 557, 574; in sprays, 20, 46, 144, 159, 168, 267, 268, 558, 559, 574, 645; soil treatment with, 577; applied from aircraft, 574; distribution of deposits of, on tobacco leaves, 574; residues of, 228, 267; tolerance limits for, 267; and DDT, 20, 168; and glyodin, 20.

DDT, against aphids, 188, 320, 333, 468, 498, 569, 633, 637, 674, 677; aphids favoured by, 549, 631; against Cicadellids, 94, 395, 563, 627; against Coccids, 168, 434; against Mirids, 72, 213, 408, 467, 478, 515, 652; against Pentatomids, 57, 115, 653; question of susceptibility of Eurygaster integriceps to Beauveria bassiana in relation to, 115; against other Hemiptera, 38, 39, 94, 95, 235, 241, 395, 437, 523, 593, 676; against Curculionids, 56, 114, 117, 118, 180, 194, 275, 317, 322, 347, 360, 408, 412, 467, 500, 506, 512, 515, 540, 609, 637; resistance to, in strains of Anthonomus grandis, 218, 219; unsatisfactory against Ceutorhynchus assimilis, 373; against Leptinotarsa decemlineata, 247, 447, 539, 683; effects of, on development and susceptibility of L. decemlineata, 363, 364, 365; resistance to, in strains of L. decemlineata, 252, 318, 440; against Melolonthids, 171, 493; ineffective against Melolontha hippocastani, 541; against Scolytids, 217, 460, 489, 641; against other Coleoptera, 28, 38, 40, 58, 61, 118, 173, 179, 202, 203, 211, 244, 265, 297, 314, 316, 324, 349, 418, 448, 497, 502, 600, 608, 647, 687; influence of oxygen uptake on susceptibility of Alphitobius laevigatus to, 201; question of resistance to, in strains of

Popillia japonica, 418; against Drosophila, 147; effects of dietary nitrogen on susceptibility of D. melanogaster to, 638; selection increasing resistance to. in D. melanogaster, 62; against Melanagromyza phaseoli, 173, 484; toxicity of, to Musca domestica, 171, 202, 203; resistance or susceptibility to, in strains of M. domestica, 63; against Trypetids, 18, 47, 98, 137, 175, 188, 268, 273, 274, 311, 438, 535, 668, 676, 680; resistance or susceptibility to, in strains of Rhagoletis pomonella, 400; against other Diptera, 101, 179, 362, 484, 549, 623; against Noctuids, 27, 179, 211, 227, 234, 269, 275, 277, 324, 345, 406, 408, 409, 412, 413, 503, 512, 514, 515, 530, 538, 539, 557, 569, 580, 630, 686; Heliothis zea favoured by, 323; resistance to, in strain of H. zea, 406; experiments on penetration of, in relation to larval age of *H. zea*, 472; against Pyralids, 61, 79, 80, 164, 179, 333, 503, 514, 630; unsatisfactory against Zeadiatraea grandiosella, 21; against Tineids, 198, 302, 347, 409, 430, 441, 525, 536, 557, 596; resistance to, in strains of Pectinophora gossypiella, 473; Leucoptera sp. favoured by, 653; against Tortricids (sens lat.), 18, 19, 20, 21, 26, 57, 60, 108, 121, 122, 239, 248, 337, 360, 398, 411, 413, 425, 455, 461, 466, 554, 555, 592, 646, 659; resistance to, in strain of Cydia pomonella, 394; ineffective against larvae of C. pomonella in apples, 637; Eulia loxonephes favoured by, 144; ineffective against *Trichoplusia ni*, 413; against other Lepidoptera, 35, 46, 48, 55, 85, 92, 99, 111, 159, 161, 220, 226, 246, 267, 269, 336, 399, 431, 448, 503, 513, 514, 525, 534, 535, 539, 550, 659, 661; resistance to, in strains of Estigmene acraea, 480; resistance to, in relation to virus disease of Malacosoma neustria, 359; against sawflies, 10, 32, 56, 139, 163, 362, 540, 541, 544; against Eurytoma amygdali, 110; effects of, on populations of Oecophylla, 29; against Vespula germanica, 422, 423; against termites, 77, 180, 577, 627; diverse effects of, against grasshoppers and locusts, 172, 173, 498; against Thysanoptera, 71, 151, 193, 345, 348, 408, 467, 503, 515, 569, 653; against mites, 264, **281**, **419**, **439**, **626**, **628**; Tetranychids favoured by, 239, 408, 523, 544, 633; effects of, on populations of Tetranychids, 35, 36, 108, 111; effect of, on egg-production by Tetranychids, 439; against Scutigerella immaculata, 433;

effects of, on natural enemies of pests. 35, 36, 159, 217, 227, 302, 434, 439, 512, 523, 560, 627, 653, 683; persistence of parasites of Lymantria dispar in areas treated with, 399; toxicity of, to bees, 412; toxicity of, to larvae of Bombyx mori, 171; effects of, on other terrestrial and aquatic arthropods, 217, 454, 461, 462; metabolism and excretion of, in insects, 64, 148, 218, 248; effects of, on fecundity of survivors, 55; experiments with cockroaches mode of action of, 62, 63; factors affecting toxicity of, 21, 117, 118, 202, 203, 356; distribution of resistance to, in agricultural pests, 284; review of resistance to, in strains of arthropods, 284; genetic basis of resistance to, 147; toxicity of, to birds, 217; toxicity of, to fish, 12, 304; effects of, on growth and physiology of plants, 95, 220, 360, 562, 632, 652; question of use of, on tea, 449; question of systemic activity of, 83, 146; solutions of, 117, 118, 537; emulsified solutions of, 26, 39, 56, 57, 72, 77, 85, 172, 179, 188, 194, 227, 246, 252, 275, 316, 320, 360, 399, 413, 438, 441, 448, 460, 478, 489, 500, 512, 554, 592, 600, 641, 668, 677; in wettablepowder suspensions, 28, 29, 48, 56, 57, 77, 80, 168, 179, 226, 275, 360, 362, 448, 466, 557, 627, 646; in aerosols, 10, 27, 55, 85, 247, 348, 418, 419, 536, 537, 539, 550, 676, 680; in bait-sprays, 98; 574, 580, 596, 608, 627, 633, 637, 659, 661; in granules, 21, 641; in atomised sprays, 85; in smokes, 195; in sprays, 12, 18, 19, 20, 21, 26, 27, 28, 29, 35, 36, 39, 46, 47, 48, 55, 56, 57, 58, 61, 71, 72, 79, 80, 85, 94, 98, 108, 110, 111, 115, 121, 122, 137, 139, 159, 163, 168, 173, 175, 179, 180, 188, 193, 195, 198, 211, 213, 217, 218, 220, 226, 227, 228, 235, 408, 411, 412, 413, 425, 434, 437, 438, 439, 441, 448, 449, 455, 460, 461, 466, 468, 478, 484, 502, 503, 512, 514, 515, 523, 525, 530, 534, 535, 539, 541, 544, 549, 554, 555, 557, 574, 592, 593, 609, 626, 630, 631, 632, 633, 637, 641, 646, 652, 653, 668, 670, 674, 676, 677, 680, 683, 686; as deposit, 195, 273, 316; persistence and toxicity of deposits of,

20, 21, 356, 394; distribution of deposits of, on tobacco leaves. 574: comparison of high- and low-volume applications of, 670; applied by aircraft, 55, 121, 122, 220, 395, 413, 425, 455, 530, 540, 541, 569, 574, 659, 680; seed treatment with, 433; soil treatment with, 38, 71, 77, 171, 180, 275, 347, 431, 493, 512, 577, 578, 627; persistence of, in soil, 83, 228, 272, 416, 567, 578; vapour toxicity of, 27, 416; treatment of selfing bags with, 277; treatment of planting material with, 77, 194, 362, 661: treatment of nests with, 422, 423, 627; tests and uses of, against pests of stored products, 28, 179, 195, 213, 347, 500; treatment of thatched roofs with, 596; treatment of timber with, 448, 489, 647; residues of, in plants and plant products, 267, 398, 439, 513, 578, 680; residues of, in milk and tissues of cattle, 266, 384; residues of, in earthworms, 641; tolerance limits for, 267; analysis of, 91, 356, 620; determination of, in maize meal, 694; labelled with radioactive isotopes, 218, 472; use of, in evaluation of potency of insecticides, 221; and Aramite, 21; and azinphosmethyl, 21, 161, 409, 467, 515; and Bacillus thuringiensis, 514, 525; effects of Beauveria bassiana used in combination with, 447; and BHC, 10, 61, 99, 179, 198, 247, 252, 297, 345, 349, 409, 441, 503, 538, 539, 540, 541, 600, 608, 674, 676; and lead arsenate, 268, 274; and bordeaux mixture, 20; and chlorbenside, 674; and chlordane, 419; and chlorfenson, 21; and chlorobenzilate, 21; and copper sulphate, 438; and coumarone indene resin. 29: 2,4-D, 348; and DDD, 20, 168; and decanoic acid, 217; and Delnav, 21; and demeton, 21, 637; and diazinon, 239, 268; and N, N-di-n-butyl-p-chlorobenzenesulphonamide, 641; and odichlorobenzene, 419; and dieldrin, **441:** and 3,4-diethoxybenzaldehyde, N,N-diethyl-m-toluamide, 218; and 218; and dimethoate, 239; and dimethyl phthalate, 218; and endrin, 413, 515; and EPN, 21; and ethion, 21; and fungicides, 20, 21, 26, 36, 264, 523; and Genite, 21; and Indalone, 218; and Kelthane, 21; and lime, 641; and limesulphur, 20, 36, 544; and malathion, 466; and methyl-parathion, 322, 515; and oils, 20, 472, 631, 686; and oil emulsions, 434, 631; and parathion, 39, 137, 239, 248, 311, 437, 438, 466, 534, 535, 637; and phorate, 21; and pyrethrum, 418, 633; and rotenone, 337, 525;

and Strobane, 406, 412, 413, 515; and sulphur, 20; and Tedion, 21; and Thiodan, 631, 637; and toxaphene, 220, 406, 408, 412, 413, 467, 468, 478, 480, 514, 515, 569, 632, 633, 637; question of synergism of toxaphene with, 218; and trichlorphon, 676; and Trithion, 21, 268; and other phosphorus compounds, 412, 413; effects of, on action of phosphorus compounds, 62; experiments with synergists and, 687.

DDVP (see Dichlorvos). dealbana, Gypsonoma. debachi, Physcus.

2,3,3a,4,5,6,7,7a,8,8-Decachloro-3a,4,-7,7a-tetrahydro-4,7-methanoindene-1-one (Decachlorooctahydro-1,3,4-metheno-2H-cyclobuta[cd]pentalen-2-one), against aphids, 333; against fire ants, 464, 639; against Lepidoptera, 26, 222, 269, 333, 514; ineffective against Cydia pomonella, 466; against pests in insect cultures, 339; harmless to beneficial arthropods, 159; in baits, 339, 464, 639; in sprays, 26, 159, 222, 269, 333, 466, 514; recovery of residues of, from milk, 266.

Decadarchis, infesting dry fruits of

tamarind in India, 499.

Decanoic Acid, in sprays against Scolytus multistriatus, 217; and DDT, 217, 218. decedens, Empoasca.

decemlineata, Leptinotarsa (Chrysomela);
Polyphylla.

deceptor, Copidosoma.

decipiens, Empoasca; Sitona.

deesae, Glyptomorpha (Stenobracon).

Delaware, pests of Capsicum in, 630; pests of potato in, 333; mite fauna of soy-bean fields in, 414; Hypera variabilis on lucerne in, 400; Ostrinia nubilalis on maize in, 509, 511; Bryobia praetiosa infesting houses in, 419; beneficial arthropods in, 400, 414, 509; Perezia pyraustae infecting O. nubilalis in, 511. delicatula, Psoloessa.

Delnav (2,3-p-Dioxane S,S-Bis(O,O-diethyl phosphorodithioate)), against Dermestes, 49; against Hemiptera, 94, 217, 627; Chrysomphalus ficus favoured by, 645; against Lepidoptera, 21, 513; against mites, 217, 515, 645; cross-resistance to, in strain of Tetranychus pacificus, 512; effects of, on yield of oranges, 645; in dusts, 217, 627; in sprays, 94, 217, 515, 645; soil treatment with, 627; treatment of skins with, 49; toxicity of DDT increased by, 21.

Delonix regia (see Poinciana).
Delphacodes striatella (see Calligypona).

Delphinium, Kakimia robinsoni on, in Manitoba, 529.

deltiger, Toxares.

Deltinea costalimai, gen. et sp. n., on Solanum bonariense in Argentina, 230; feeding on orange, 230.

Deltocephalus, on beet in California, 630;

parasites of, 630.

demeijerei, Microtrombidium.

Demeton, against aphids, 249, 278, 279, 331, 468, 511, 550, 559, 605, 608, 624, 637; effects of, on respiration of aphids, 444; against Cicadellids, 395, 559, 625; diverse effects of, against Coccids, 19, 265, 543; against Coleoptera, 138, 511, 637, 673; against Diptera, 45, 253, 625; against Lepidoptera, 21, 136, 273, 441, 642: toxicity of, to Locusta migratoria, 172; against Thysanoptera, 257, 348; ineffective against Frankliniella, 352; against Eriophyids, 191, 241; against Larvacarus transitans, 626; against Tetranychoids, 19, 183, 264, 324, 408; 517, 559, 570; resistance to, in strains Tetranychids, 147, 272, 510, 512, effects of, on beneficial arthropods, 159, 183; toxicity of, to honey bees, 550; metabolism of, in insects, 64; effects of, on growth and respiration of oats and peas, 562; phytotoxicity of, 277, 625; in granules, 277; in sprays, 45, 138, 159, 183, 191, 241, 249, 253, 264, 273, 279, 324, 331, 348, 395, 408, 441, 468, 511, 517, 543, 550, 570, 605, 608, 624, 626, 637, 642; as soil drench, 265; treatment of planting material with, 191, 278, 673; tests of systemic activity and other methods of application of, 19, 136, 249, 257, 273, 277, 279, 352, 559, 625; residues of, 570, 624; tolerance limit for, 570; and 2,4-D, 348; toxicity of DDT decreased by, 21; and other insecticides, 264; and fungicides, 264, 324.

Demissin, experiments with, on resistance of Solanum to Leptinotarsa decemlin-

eata, 318.

demoleus, Papilio.

Dendroctonus barberi, bionomics of, on Pinus ponderosa in U.S.A., 460; natural enemies of, 460; DDT sprays against, 460.

Dendroctonus brevicomis, on Pinus ponderosa in U.S.A., 340; pine-resin vapour in relation to infestation by, 223, 340.

Dendroctonus engelmanni, population fluctuations of, on spruce in Colorado, 150.

Dendroctonus frontalis, review of, on pines in U.S.A., 495.

Dendroctonus jeffreyi, fumigant toxicity of pine-resin vapour to, 223.

Dendroctonus monticolae (see D. pon-

derosae).

Dendroctonus obesus, on conifers in Canada and U.S.A., 532; intercepted on Douglas fir imported into Italy, 532.

Dendroctonus ponderosae, on Pinus ponderosa in Nevada, 340; water supply and resin exudation in relation to

infestation by, 340.

Dendroctonus pseudotsugae (on Douglas fir), in British Columbia, 458; in Washington, 259; factors affecting flight habits of, 127; associated with Scolytus unispinosus, 458; X-ray detection of, in timber, 259.

Dendroctonus valens, attracted to logs infested by Ips confusus in California,

341.

dendrolimi, Rogas.

Dendrolimus pini (on pines), in Germany, 376; in U.S.S.R., 598, 599; factors affecting populations and mortality of, 376, 599; characteristics and world distribution of, 507.

Dendrolimus punctatus, bionomics of, on pine in China, 501, 623, 624; dispersal of larvae of, 501; parasites of, 624, 625.

Dendrolimus superans (sibiricus) (on conifers), bionomics of, in Mongolia, 116; in U.S.S.R., 13, 344; natural enemies of, 13, 116, 344.

dendrolimusi, Telenomus.

Dendrosoter middendorffi, parasiting Ips typographus in Czechoslovakia, 142; method of rearing, 142.

Dendrothrips bispinosus, associated with abnormal growth of coffee in India,

448.

Denmark, Chermes nordmannianae on Abies alba in, 674; Hydrellia griseola on cereals and grasses in, 354; pests of crucifers in, 139; Hylemya antiqua on onion in, 608; Plutella maculipennis in, 198, 430.

dentata, Phanerotoma.

denticornis, Cerodontha; Hyalochloria; Limothrips.

deplanatus, Anacentrinus.

Depressaria apiella (nervosa), bionomics of, on caraway in Czechoslovakia, 301; parasites of, 301; measures against, 302.

depressella, Emmalocera.

Deraeocoris punctulatus, in Turkey, 352. Dermestes, infesting hides and skins in S. Africa, 49; experiments with insecticides against, 49; method of rearing, 494.

Dermestes ater (cadaverinus), intercepted on oil-seed cake in Germany, 11.

Dermestes maculatus, infesting hides and skins in S. Africa, 49; experiments with insecticides against, 49; toxicity of extract from Bacillus larvae to, 332.

derogata, Sylepta.

Apleurotropis lalori mis-Derostenus, identified as species of, 166.

Derris, against Lepidoptera, 42, 179; in dust, 42; in spray, 179.

Desert Locust (see Schistocerca gregaria).

desertorum, Tetranychus.

destructor, Aspidiotus; Ceroplastes; Ctenicera aeripennis; Glycyphagus; Halotydeus; Mayetiola (Phytophaga); Tribolium.

detritus, Ephialtes (Scambus).

Dexia rustica, parasitising larvae of Melolontha melolontha in Czechoslovakia, 187.

Dextrin, in diets for insects, 5, 495.

Dextrose, in relation to feeding response of wireworms, 528; in diets for insects, 389, 626; in rearing medium for Ostrinia nubilalis, 22.

DFP (see Diisopropyl Fluorophosphate). Diabrotica balteata, on sweet potato in

Louisiana, 131.

Diabrotica graminea, experimental transmission of bacterial wilt of banana by, 520.

Diabrotica longicornis, insecticides against, on maize in U.S.A., 396, 420.

Diabrotica speciosa, characteristics and world distribution of, 506.

Diabrotica undecimpunctata howardi, on maize in Illinois, 420; insecticides against, 420.

Diacrisia lubricipeda, on apple and pear in Bulgaria, 59; in U.S.S.R., 115; Apanteles glomeratus not developing in, 115.

Diadegma, status and synonymy of, 493. Diadromus subtilicornis, parasitising Plutella maculipennis in U.S.S.R., 245.

Diaeretus rapae (parasitising aphids), in New York, 401; integration of chemical control with, 148.

N,N-Diallyl-3-(dimethoxyphosphinyloxy)crotonamide, against Musca domestica, 63; with synergists, 63.

Diammonium Phosphate, in baits for Trypetids, 439, 533, 643.

diaperinus, Alphitobius.

Diapetimorpha venturii, parasitising Rhyacionia buoliana in Argentina, 144. Diaphania indica (see Margaronia).

Diaphania nitidalis, varietal susceptibility of squash to, in N. Carolina, 471; dusts against, 471.

Diaphorina citri, characteristics and world distribution of, 507.

Diarthrothrips coffeae, sprays against, on coffee in Tanganyika, 653.

Diaspidiotus perniciosus (see Quadraspidiotus).

diaspidis, Aphytis.

Diaspis boisduvalii, on Cattleya in Indonesia, 143.

Diaspis pentagona (see Pseudaulacaspis). Diataraxia oleracea, in U.S.S.R., 115; feeding habits of, 620; parasites of, 87, 115; method of rearing, 606.

Diatomite, against pests of stored products, 51; as carrier for insecticide dusts, 51.

Diatraea, on sugar-cane in Venezuela, 357; estimation of losses caused by.

Diatraea saccharalis (on sugar-cane), in Jamaica, 289; in Louisiana, 216, 280, 469; diapause in, 217; parasite of, 289; Bacillus thuringiensis against, insecticides against, 280, 289, method of rearing and diets for, 214, 216.

Diatraea venosatus (see Proceras).

diatraeae, Lixophaga.

Diazinon, against aphids, 217, 279, 324, 402, 412, 468, 608, 616, 633, 637, 643, 674; against Cicadellids, 395, 590; against Coccids, 120, 612, 616; ineffective against Eurygaster, 382; against Coleoptera, 180, 265, 323, 471, 493, 497; against Noctuids, 398, 590, 644; against Tineids, 31, 533, 663; against Tortricids (sens. lat.), 239, 248, 337, 466; against other Lepidoptera, 85, 309, 480, 514, 534, 669; against Trypetids, 47, 98, 135, 137, 175, 237, 240, 268, 273, 309, 311, 438, 533, 535, 668; against other Diptera, 388, 674; against Kakothrips pisivorus, 193; toxicity of, to locusts, 172, 383, 657; factors affecting resistance to, in locusts, 201; against Sminthurus viridis, 34; against Tetranychids, 28, 217, 242, 324, 515, 616, 628; toxicity of, to parathion-resistant strain of Tetranychus telarius, 278; cross-resistance to, in strain of T. pacificus, 512; against other mites, 396, 545, 626; effects of, on beneficial arthropods, 159, 312; metabolism of, in insects, 64; factors affecting toxicity of, 274, 657; in dusts, 217, 323, 637, 644, 674; in granules, 323; in bait-sprays, 98; in sprays, 28, 31, 34, 47, 85, 120, 135, 137, 159, 175, 193, 213, 217, 237, 239, 240, 242, 248, 268, 279, 309, 311, 312, 324, 337, 383, 305, 306, 308, 402, 438, 466, 469, 514 395, 396, 398, 402, 438, 466, 468, 514,

515, 533, 534, 535, 545, 590, 608, 616, 626, 633, 643, 657, 663, 668, 669, 674; evaluation of spray programmes of. 376; as deposit, 240, 273; soil treatment with, 180, 388, 493; treatment of planting water with, 388; tests of, against pests of stored products, 213, 471; factors affecting persistence of, 471; humidity in relation to fumigant action of, 471; residues of, in plants and plant products, 238, 309, 398; effectiveness of, improved by copper sulphate, 47; and DDT, 239, 268; and malathion, 268; and oil emulsions, 120, 616.

Dibrachys cavus, parasitising larvae of Ostrinia nubilalis in Quebec, 27.

Dibrom (see Dimethyl 1,2-Dibromo-

2.2-dichloroethyl Phosphate).

1,2-Dibromo-3-chloropropane, vapour toxicity of, to Phylloxera vitifoliae, 572. 1,2-Dibromopropane, vapour toxicity of,

to Phylloxera vitifoliae, 572.

1,3-Dibromopropane, vapour toxicity of, to Phylloxera vitifoliae, 572.

N,N - Di-n - butyl - p - chlorobenzenesulphonamide, in sprays against Scolytus multistriatus, 641; and DDT, 641.

Dicapthon, against Dacus oleae, 135; against Lymantria dispar, 514; sprays, 135, 514.

Dichlone (see 2,3-Dichloro-1,4-naphtho-

quinone).

o-Dichlorobenzene, in aerosols, against Bryobia praetiosa, 419; and chlordane, 419; and DDT, 419.

p-Dichlorobenzene, against Lachnosterna consanguinea, 171; against Chloropsalta viridissima, 609; soil treatment with, 171, 609.

2.3-Dichloro-1.4-naphthoguinone, effects of sprays containing, on arthropods on apple, 159, 274; toxicity of insecticides decreased by, 274.

2,4-Dichlorophenoxyacetic Acid, effects of sprays containing, on Haplothrips

tritici, 348.

N-(3,4-Dichlorophenyl)-N'-2-(2-sulpho-4 - chlorophenoxy) - 5 - chlorophenyl urea, treatment of textiles with sodium salt of, against Hofmannophila pseudospretella, 290.

Di (p-chlorophenyl) - trifluoromethylcarbinol, toxicity of, to mites, 684.

1,2-Dichloropropane, in D-D mixture, 171, 609.

1,3-Dichloropropene, in D-D mixture,

171, 609.

Dichlorvos, against Diptera, 562, 623; in tests against Lymantria dispar, 514; against Popillia japonica, 226, 418; against Trypetids, 268, 643; metabolism of, 64, 213; in aerosols, 418, 623; in baits, 643; in sprays, 212, 514; stabilised formulations of, 561, 562; tests of vaporisers for, 226; uses of, against pests of stored products, 212.

Dichocrocis punctiferalis, on maize in China, 502; cold-hardiness of, 502; characteristics and world distribution

Dichomeris marginella, on juniper in New York, 19; tests of systemic insecticides against, 19.

dichroum, Apion.

Dicladispa armigera, characteristics and world distribution of, 507.

Dicladocerus, parasitising Evagora milleri in California, 222; effects of sprays on populations of, 222.

Dicranura erminea menciana, damaging

lead cable in China, 693.

Didymuria violescens, bionomics of, on Eucalyptus in Australia, 32, 33, 231, 232; parasites of, 32; population studies of, 33, 231, 232; effects of sprays on populations of, 33.

Dieldrin, against aphids, 498, 549; against Coccids, 557, 667; against other Hemiptera, 72, 230, 235, 321, 498, 653; against Curculionids, 16, 114, 180, 219, 280, 317, 331, 338, 360, 373, 374, 417, 500, 509, 533, 636; against other Coleoptera, 38, 49, 171, 173, 265, 310, 316, 323, 324, 373, 374, 449, 489, 493, 497, 526, 533, 554, 570, 608, 644, **651**, **674**, **687**; influence of oxygen uptake on susceptibility of Alphitobius laevigatus to, 201; against Hylemya spp., 50, 226, 433, 568, 608, 674; question of resistance to, in Hylemya spp., 388; against Psila rosae, 101, 226, 432, 608; against Trypetids, 175, 240, 325, 417, 590; against other Diptera, 173, 430, 484, 549, 623; against Noctuids, 27, 210, 211, 269, 347, 383, 514, 530, 686; against Pyralids, 79, 297, 503; against other Lepidoptera, 16, 26, 198, 226, 269, 273, 277, 290, 317, 335, 441, 503, 641, 653; against grass-hoppers, 498, 521; factors affecting toxicity of, to locusts, 164, 172, 608, 657; metabolism and excretion of, in Schistocerca gregaria, 321, 322; against Phasmatids, 33; effects of, on populations of Oecophylla, 29; against termites, 77, 119, 180, 293, 294, 416, 577, 627; against Thysanoptera, 71, 463, 477, 503, 653; effects of, on beneficial insects, 554, 653; toxicity of, to bees, 374, 412; toxicity of, to fish, 304; restriction on use of, in relation to hazards to wild life, 433; review of resistance to, in strains of arthropods, 284; mode of action of, in cockroaches, 62; phytotoxicity of, 50, 101, 503, 562; other effects of, on growth of plants, 360, 627, 674; in baits, 590; in dusts, 49, 114, 293, 294, 323, 347, 514, 568; in granules, 219, 323, 325, 331, 568, 570, 636, 641; in sprays, 16, 26, 27, 29, 33, 71, 72, 79, 165, 173, 175, 180, 198, 226, 230, 235, 240, 269, 273, 310, 317, 321, 330, 335, 338, 360, 373, 374, 383, 430, 441, 449, 463, 477, 498, 503, 509, 530, 533, 538, 539, 549, 554, 557, 568, 608, 636, 641, 644, 651, 653, 657, 667, 686; as deposit, 240; applied by aircraft, 463, 530; seed treatment with, 50, 65, 66, 71, 101, 226, 433, 484, 568, 608, 627, 674; soil treatment with, 38, 50, 71, 171, 180, 219, 226, 280, 293, 294, 324, 325, 347, 432, 493, 526, 568, 570, 577, 578; persistence of, in soils, 77, 119, 339, 416, 417, 432, 527, 578; treatment of planting material with, 77, 119, 297, 557; treatment of selfing bags with, 277; treatment of planting water with, 388; treatment of nests with, 294, 627; uses of, against pests of stored products, 49, 500; treatment of textiles with, 290; treatment of timber with, 489; treatment of concrete with, 416; question of vapour toxicity of, 27; residues of, on plants and plant products, 330, 417, 578; residues of, in milk and tissues of animals, 88, 266, 384, 521; residues of, in earthworms, 641; factors affecting persistence of residues of, 16, 570; analysis and determination of, 88, 91; as metabolite of aldrin, 64, 146, 272, 322, 384, 417; bioassay of, 146, 397, 412; labelled with 36C1, 321; and BHC, 441; and bordeaux mixture, 651; and DDT, 441; and fungicides, 50; and oil emulsion, 463, 667, 686; and parathion, 477; effects of, on action of phosphorus compounds, 62; and sulphuric acid, 321; experiments with synergists and, 63, 687.

Dielmoth, containing dieldrin (q,v_{\cdot}) , 290. 3,4-Diethoxybenzaldehyde, in sprays against Scolytus multistriatus, 217.

2-Diethylaminoethanol, toxicity of pyrethrins unaffected by, 688.

2-Diethylaminoethyl 2,2-Diphenylpentanoate, experiments on synergism of, Alphitobius laevigatus against and Musca domestica, 687.

Diethyl p-Chlorophenyl Phosphate, anticholinesterase activity and toxicity of, to Musca domestica, 634.

Diethyl 2-Chlorovinyl Phosphate, against aphid. 508: against Diprion similis, 418, 419; soil treatment with, 508; factors affecting systemic activity of, 418, 508.

Diethyl 1-(2,4-Dichlorophenyl)-2-chloro-Phosphate, toxicity of, vinvl Estigmene acraea, 480; in sprays against

Myzus persicae, 216.

Diethyl 1-(2,5-Dichlorophenyl)-2-chloro-Phosphate, toxicity of,

Estigmene acraea, 480.

1-(2,5-Dichlorophenyl)-2,2-Diethyl dichlorovinyl Phosphate, toxicity of, to Estigmene acraea, 480.

O.O-Diethyl O-2.4-Dichlorophenyl Phosphorothioate, against Eumargarodes laingi, 265; as soil drench, 265.

Diethyl 2,2-Di(ethylthio)vinyl Phosphate,

toxicity of, to Aphis gossypii, 479. iethyl 1,3-Di(methoxycarbonyl)-1propen-2-yl Phosphate, toxicity of, to Aphis gossypii, 479.

Diethyl 1-(Dimethylcarbamoyl)-1-propen-2-yl Phosphate, toxicity of, to Aphis

gossypii, 479.

Diethyl Dithiophosphorylethylcarbamate,

in sprays against aphids, 402.

Diethyl 2 - Ethoxycarbonyl - 1 - methyl vinyl Phosphate, in sprays against Fenusa pusilla, 19; trunk applications of, 19.

O,O-Diethyl S-2-(Ethylsulphinyl)ethyl Phosphorodithioate, against aphid, 508; soil treatment with, 508; factors affecting systemic activity of, 508.

O,O-Diethyl S-Ethylsulphinylmethyl Phosphorodithioate, anticholinesterase activity and toxicity of, to Musca 512; domestica, metabolite as phorate, 512, 513.

O,O-Diethyl S-Ethylsulphinylmethyl Phosphorothioate, anticholinesterase activity and toxicity of, to Musca domestica, 512; as metabolite

phorate, 512, 513.

O,O-Diethyl S-Ethylsulphonylmethyl Phosphorodithioate, anticholinesterase activity and toxicity of, to Musca domestica, 512; as metabolite phorate, 512, 513.

O,O-Diethyl S-Ethylsulphonylmethyl Phosphorothioate, anticholinesterase activity and toxicity of, to Musca 512; as domestica. metabolite

phorate, 512, 513.

O,O-Diethyl S-2-(Ethylthio)ethyl Phosphorodithioate, against aphids, 278, 281, 331, 401, 468, 508, 510, 511, 605, 645, 670, 680; against Cicadellids, 325, 405, 470; against other Hemiptera,

409, 510; against Coleoptera, 16, 325, 511; against Diptera, 253, 268, 330; against Lepidoptera, 273, 632; against Thysanoptera, 257, 409, 515, 645; against mites, 19, 645; effects of, on natural enemy of Toxoptera graminum, 281; metabolism of, in insects, 64; effects of, on plants, 65, 330, 399, 401, 556, 605, 670; in dusts, 278; in granules, 16, 273, 278, 281, 325, 331, 405, 409, 468, 470, 510, 515, 605, 645, 670, 680; in sprays, 19, 515, 645; seed treatment with, 65, 253, 399, 401, 405, 470, 605, 670; soil treatment with, 273, 278, 281, 325, 331, 405, 409, 468, 508, 510, 511, 605, 645, 670, 680; movement of, in soil, 475; treatment of bulbs with, 278; tests of systemic action and other methods of application of, 16, 19, 257, 278, 330, 405, 470, 475, 508, 556, 632, 633; labelled with ³²P, 475; question of use of fertilisers with, 645.

O,O-Diethyl S-Ethylthiomethyl Phos-

phorodithioate (see Phorate).

O,O-Diethyl S-Ethylthiomethyl Phosphorothioate, toxicity of, to *Musca domestica*, **512**; metabolism of, in cockroaches, **513**; anticholinesterase activity of, **512**.

Diethyl 1-Ethylthio-1-propen-2-yl Phosphate, toxicity of, to *Aphis gossypii*,

479.

- O,O-Diethyl S-Isopropylcarbamoylmethyl Phosphorodithioate (in sprays), against Dacus oleae, 237, 238; against Penthaleus major, 532; residues of, in olives, 238.
- O,O-Diethyl S-Isopropylthiomethyl Phosphorodithioate, tests of systemic activity of, against insects and mites, 19.
- O,O-Diethyl O-p-Methylsulphinylphenyl Phosphorothioate, against Lepidoptera, 409; against Lygus lineolaris, 515; toxicity of, to Rhagoletis pomonella, 268; in dusts, 409; in spray, 515.

Diethyl Naphthalimido Phosphate, in tests against *Lymantria dispar*, **514**.

O,O-Diethyl O-Naphthalimido Phosphorothioate, against Lepidoptera, 466, 514; against Rhagoletis pomonella, 268; in sprays, 268, 466; residues and stability of, in milk, 266; estimation of residues of, 196.

O,O-Diethyl S-p-Oxathian-3-yl Phosphorodithioate, toxicity of, to *Antho-*

nomus grandis, 338.

O,O-Diethyl O-(2-Phenyl-6-(3(2H)-pyridazinonyl)) Phosphorothioate, toxicity of, to *Tetranychus telarius*, 478.

O,O-Diethyl S-Phthalimidomethyl Phosphorodithioate, toxicity of to *Anthonomus grandis*, 338.

O,O-Diethyl S-Phthalimidomethyl Phosphorothioate, toxicity of, to *Tetranychus*

telarius, 478.

O,O-Diethyl S-Propylthiomethyl Phosphorodithioate, tests of, against pests of stored products, 212, 213; in sprays, 212.

- O,O-Diethyl O-2-Pyrazinyl Phosphorothioate, against Coleoptera, 325; diverse effects of, on Lepidoptera, 325, 469; against *Hylemya brassicae*, 510; in granules, 325, 510; soil treatment with, 325, 510; applied in drenches, 510.
- Diethyl m-Sulphurpentafluorophenyl Phosphate, toxicity of, to *Musca domestica*, 634; anticholinesterase activity of, 634.

N,N-Diethyl-m-toluamide, in sprays, against Scolytus multistriatus, 217.

differentialis, Melanoplus,

Digitaria decumbens, toxaphene residues in, 64.

dignus, Telenomus.

Diisopropyl Fluorophosphate, enzymatic hydrolysis of, by arthropod and mammalian homogenates, 213.

Dilan (mixture of Bulan and Prolan), diverse effects of, on Lepidoptera, 406, 466.

700

Dimecron 20, containing phosphamidon (q.v.), 237.

Dimecron-Combi, containing phosphamidon (q.v.), 613.

Dimefox, metabolism of, in insects, 64.

Dimetan, tests of toxicity of, to insects, 202; effect of temperature on toxicity of, 202; colorimetric determination of, 227.

Dimethoate, against aphids, 74, 216, 249, 279, 395, 401, 402, 431, 467, 473, 505, 508, 510, 524, 549, 633; ineffective against Phylloxera vitifoliae in galls, 668; against Tingids, 38, 554; against other Hemiptera, 80, 158, 395, 470, 510, 639; against Curculionids, 180, 338, 467; against other Coleoptera, 418, 511; against Cydia pomonella, 239, 466; against eggs of Operophtera brumata, 669; against other Lepidoptera, 222, 328, 329, 411, 421, 533, 534, 663; ineffective against Lepidoptera, 21, 469; against *Frankliniella*, 515; against Trypetids, 47, 48, 135, 137, 237, 238, 240, 268, 311, 438, 533, 535, 562, 668, 669; against other Diptera, 549, 674; against Sminthurus viridis, 34; against mites, 266, 288, 289, 396, 431,

628; effects of, on beneficial insects, 158, 627; 639; phytotoxicity of, 277, 401, 467; in aerosols, 418; in dusts, 38, 47, 237, 438, 515; in granules, 21, 277, 554, 633, 663, 668, 669, 674; seed treatment with, 277, 401, 467, 470; soil treatment with, 180, 277, 508, 510, 511; as drench, 395; tests of, against pests stored products, 213; tests of systemic activity and other methods of application of, 216, 249, 279, 395, 431, 467, 470, 472, 473, 524, 663; factors affecting systemic activity of, 277, 328, 329, 431, 473, 508; penetration of, into fruits, 669; question of residues and metabolism of, in olives and olive oil, 48, 238, 439, 562; residues of, on other plants, 395, 674; labelled with ³²P, 467, 472, 562; and DDT, 239.

2,4-Dimethylbenzyl Dimethrin (see

Chrysanthemate).

1,1-Dimethyl-3-(p-acetamidophenyl)triazene, in sprays against Trichoplusia ni, 222, 569; in dust, 569. (See also 4'-(Dimethyltriazeno) Acetanilide.)

4-Dimethylamino-3,5-xylyl Dimethylcarbamate, insecticidal and anticholinesterase activities of, 634, 635; and

piperonyl butoxide, 635.

4-Dimethylamino-3,5-xylyl Methylcarbamate, toxicity of, to Lepidoptera, 476, 508, 569, 637; against Liriomyza, 508; against Scolytus multistriatus, 641; against Thysanoptera, 477, 508, 569; beneficial insects destroyed by, 508; insecticidal and anticholinesterase activities of, 515, 516, 634, 635; in dust, 569; in sprays, 477, 569, 637, 641: synergists with, 476, 635.

2,4-Dimethylbenzyl Chrysanthemate, in tests against Lymantria dispar, 514.

Dimethyl 1-Benzyloxycarbonyl-1-propen-2-yl Phosphate, toxicity of, to Aphis gossypii, 479.

Dimethyl 1-(p-Chlorobenzyloxycarbonyl)-1-propen-2-yl Phosphate, toxicity of,

to Aphis gossypii, 479.

S-p-Chlorophenylthio-O,O-Dimethyl methyl Phosphorodithioate (in sprays), against Anthonomus grandis, 338, 408, 515; against Aphis gossypii, 279; against Lepidoptera, 222, 515; ineffective against Trichoplusia ni, 413; against Lygus lineolaris, 515; and DDT, 413.

Dimethyl 1,2-Dibromo-2,2-dichloroethyl Phosphate, against aphids, 279, 324,

573: unsatisfactory against Therioaphis maculata, 468: against Trialeurodes vaporariorum, 573; against Coleoptera, 211, 418, 573; against Diptera, 268, 573; toxicity of, to Lepidoptera, 222, 514, 569; toxicity of, to Blattella germanica, 573; against Frankliniella occidentalis, 569; against mites, 324, 573; enzymatic hydrolysis of, by arthropod and mammalian homogenates, 213; in aerosols, 418; in dust, 569; as fumigant, 573; in sprays, 211, 213, 222, 279, 324, 468, 569; tests of, against pests of stored products, 213; question of residues of, on lemons, 573; and ethyl-DDD, 222; and oil, 211.

1-(2,4-Dichlorobenzyloxycar-Dimethyl bonyl)-1-propen-2-yl Phosphate, toxi-

city of, to *Aphis gossypii*, **479**. O,O-Dimethyl S-2,5-Dichlorophenylthiomethyl Phosphorodithioate (in sprays), against Anthonomus grandis, 338, 408, 412, 515; against Lepidoptera, 408, 413, 515; against Lygus lineolaris, 515; against Frankliniella, 408; against Tetranychus lobosus, 515; T. desertorum favoured by, 408; and DDT, 412. O,O-Dimethyl S-3,4-Dichlorophenylthio-

methyl Phosphorodithioate, toxicity of, to Anthonomus grandis, 338; Geigy

36368 containing (q.v.), 413.

O,O-Dimethyl S-1,2-Di(ethoxycarbonyl)-Phosphorothioate, ethyl action synergists on oxidation of malathion to, 687.

O,O-Dimethyl-diethylthiophosphorylacetic Acid, in sprays against Dacus

oleae, 309.

1,3-Di(methoxycarbonyl)-1-Dimethyl propen-2-yl Phosphate, diverse effects of, in sprays against Lepidoptera, 409,

Dimethyl 1-(Dimethoxyphosphinyl)vinyl Phosphate, toxicity of, to Aphis gossypii, 479.

O,O-Dimethyl S-Dimethylcarbamoylmethyl Phosphorodithioate, against

Stigmella malella, 309.

Dimethyl 1-(Dimethylcarbamoyl)-1propen-2-yl Phosphate, toxicity of, to insects, 63, 418, 419, 479, 515; against Tetranychus spp., 64, 515; in granules, systemic activity and methods of application of, 418, 419; with synergists, 63, 64.

O,O-Dimethyl S-Ethylcarbamoylmethyl Phosphorodithioate, toxicity of, to

Coleoptera, 16, 338; in granules, 16. O,O-Dimethyl S-2-(Ethylsulphinyl)ethyl Phosphorodithioate, toxicity of, Anthonomus grandis, 338.

O.O-Dimethyl S-Isopropylcarbamovlmethyl Phosphorodithioate, in sprays against eggs of Operophtera brumata. 669; ineffective against Phylloxera vitifoliae in galls, 668.

Dimethyl 1-Methoxy-2,2-dichlorovinyl Phosphate, enzymatic hydrolysis of, by arthropod and mammalian homo-

genates, 213.

Dimethyl 1-(\alpha-Methylbenzyloxycarbonyl)-1-propen-2-yl Phosphate, toxicity of, to Aphis gossypii, 479.

O,O-Dimethyl S-Methylcarbamoylmethyl Phosphorothioate, as metabolite of

dimethoate, 563.

- O.O-Dimethyl O-3-Methyl-4-methylsulphinylphenyl Phosphate, metabolism of, in rats, insects and cotton plants, 475.
- O.O-Dimethyl O-3-Methyl-4-methylsulphinylphenyl Phosphorothioate, metabolism of, in rats, insects and cotton plants, 475.

O.O-Dimethyl O-3-Methyl-4-methylsulphonylphenyl Phosphate, metabolism of, in rats, insects and cotton

plants, 475.

O.O-Dimethyl O-3-Methyl-4-methylsulphonylphenyl Phosphorothioate, metabolism of, in rats, insects and cotton plants, 475, 476.

Dimethyl 3-Methyl-4-methylthiophenyl Phosphate, tests and uses of, against Phylloxera vitifoliae in galls, metabolism of, in rats, insects and cotton plants, 475.

O,O-Dimethyl O-3-Methyl-4-methylthiophenyl Phosphorothioate

Fenthion).

O.O-Dimethyl O-3-Methyl-4-nitro-Phosphorothioate, phenyl against Tryporyza incertulas 328, 329; stage of growth of rice in relation to effectiveness of, 328, 329, in sprays, 328; mammalian toxicity of, 329.

O-p-Methylsulphinyl-O,O-Dimethyl phenyl Phosphorothioate, toxicity of,

to Tetranychus telarius, 478.

Dimethyl 1-(m-Nitrobenzyloxycarbonyl)-1-propen-2-yl Phosphate (in sprays), against Alabama argillacea, against Anthonomus grandis, 408, 413; toxicity of, to Aphis gossypii, 479.

Dimethyl 1-(p-Nitrobenzyloxycarbonyl)-1-propen-2-yl Phosphate (in sprays), against Anthonomus grandis, 408, 413; Alabama argillacea, 413; against against Lygus lineolaris, 515.

p-Nitrophenyl Phosphate, Dimethyl oxidation of, increased by sesamex, 63. O,O-Dimethyl S-p-2-Oxathien-x-yl Phosphorodithioate, toxicity Anthonomus grandis, 338.

S-(4-Oxo-benzotriazino-O,O-Dimethyl 3-methyl) Phosphorothioate, as degradation product of Guthion, 228; residues of, in cigarette smoke and tobacco, 228.

Dimethyl Phosphoric Acid, as metabolite

of fenthion in cotton, 476.

Dimethyl Phosphorothioic Acid, as metabolite of fenthion in cotton, 476.

Dimethyl Phthalate, in sprays against Scolytus multistriatus, 217; and DDT,

O.O-Dimethyl S-Phthalimidomethyl Phosphorodithioate, toxicity of, to Anthonomus grandis, 338; in sprays against Pectinophora gossypiella, 557.

O,O-Dimethyl S-Phthalimidomethyl Phosphorothioate, toxicity of, to Tet-

ranychus telarius, 478.

4'-(Dimethyltriazeno) Acetanilide sprays), against Lepidoptera, against other insects, 408. (See also 1,1-Dimethyl-3-(p-acetoamidophenyl)triazene.)

Dimethyl Trichlorophosphinate (see

Trichlorphon).

3-Methyl-5-pyrazolyl Dimetilan (see Dimethylcarbamate).

dimidiatus, Pteroptrix; Rogas.

dimorpha, Eulia.

Dinarmus dacicida (see Cyrtoptyx).

diniana, Semasia (see Enarmonia griseana). Dinitroalkylphenyl Acrylate, acaricidal action of, 10.

Dinitrobenzene Thiocyanate, in mixtures with acaricides, 439.

Dinitroisobutylphenol, sprays of triethanolamine salt of: against Leucoptera, 534; against Tetranychid, 183.

Dinocap (see 2-(1-Methyl-n-heptyl)-4,6-

dinitrophenylcrotonate).

Dinoseb, against Alphitobius laevigatus, 687; influence of oxygen uptake on susceptibility of A. laevigatus to, 201; against Lepidoptera, 111; Bryobia redikorzevi, 60; in sprays, 60, 111; synergists with, 687; (dimethylacrylic-acid ester), against Tetranychus spp., 550.

Diocalandra frumenti, on coconut in Zanzibar, 29; ants not protecting

coconut from attack by, 29.

Dioctes, status and synonymy of, 493.

Dioctyl Sodium Sulphosuccinate, wetting agent with acaricides, 672; efficiency of endrin sprays improved by, 287.

diomphalia, Lachnosterna (Holotrichia).

Dioryctria abietella, D. abietivorella misidentified as, 355.

Dioryctria abietivorella, on Douglas fir in California, 355; misidentified as D. abietella, 355.

Dioryctria cambiicola, on pines in British Columbia, 529.

Dioryctria nivaliensis, on Pinus radiata in Canary Islands, 659.

Dioryctria pseudotsugella, on conifers in British Columbia, 529.

Dioryctria reniculella, on spruce Canada, 529.

Dioryctria rossi, on ponderosa pine in British Columbia, 529.

Dioryctria zimmermanni, bionomics and natural enemies of, on pines in Illinois, 153.

Dioscorea, new Coccid on species of, in New Guinea, 234.

dioscoreae, Planococcus,

2,3-p-Dioxane S,S-Bis(O,O-diethyl phosphorodithioate) (see Delnav).

Diparopsis, characteristics and world distribution of species of, 507. Diparopsis castanea, distribution of, 357,

507; survey of data on, 507.

Diparopsis tephragramma, map of distribution of, 357.

Diparopsis watersi (on cotton), in Chad Republic, 313; in Nigeria, 53; in West Aden Protectorate, 486, 487; map of distribution of, 357; bionomics of, 53, 486, 487; relation of temperature to diapause in, 313; parasite of, 53; populations of, in relation to floods, 487; measures against, 53, 487.

2,2-Diphenylpentanoic Acid, toxicity of pyrethrins unaffected by, 688.

Diplocarpon rosae, combined against Tetranychus telarius and, 264, 324.

Diplogaster, parasitising Pristiphora erichsonii in U.S.A., 210.

Diplosis abietis pectinatae (see Agevillea abietis).

Diplostichus hamatus, parasitising Gilpinia hercyniae in Ontario, 25.

Dipotassium Phosphate, attraction of insects to, 635.

Diprion hercyniae (see Gilpinia).

Diprion pallida (see Gilpinia).

Diprion pallipes (see Microdiprion).

Diprion pini, in Germany, 251; in Spain, 659; in Sweden, 426; in U.S.S.R., 598; in Yugoslavia, 31; on pines, 31, 426, 598, 659; on spruce, 251; characteristics and world diet. istics and world distribution of, 507; competition between parasites of, 31; effects of soil fertilisers on populations of, 251; BHC dust against, 659.

Diprion sertifer (see Neodiprion).

Diprion similis (on pine), introduced into Ontario from Europe, 163; in Wisconsin, 418; systemic insecticides against, 418, 419.

Diprion verticalis (see Gilpinia).

O,O-Dipropyl O-p-Methylsulphinylphenyl Phosphorothioate, toxicity of, to Tetranychus telarius, 478,

dipsacea, Heliothis (Chloridea). Dipterex (see Trichlorphon).

Dirhicnus alboannulatus, parasitising Lepidoptera in Europe, 379; reared on Galleria mellonella, 379; experiments on feeding and fecundity of, in relation to mass-rearing, 379.

dirhodum, Macrosiphum (Metopolophium). Disodium Phosphate, attraction

insects to, 635.

dispar, Lymantria (Porthetria); Xyleborus (Anisandrus).

disparis, Anastatus.

dissimilis, Polia (see Hadena suasa).

Dissosteira carolina, infected by Nosema locustae, 582.

disstria, Malacosoma.

Distantiella theobroma (on cacao), in Ghana, 651; in Nigeria, 652; relation of, to swollen-shoot virus and die-back, 652; insecticides against, 651.

Distribution Maps, of pests, 1, 357.

Di-Syston O,O-Diethyl S-2-(see (Ethylthio)ethyl Phosphorodithioate). Di-Syston Sulphoxide (see O,O-Diethyl S-2-(Ethylsulphinyl)ethyl Phosphoro-

dithioate). Dithane (see Nabam). divareti, Paremydica. divergens, Scaptocoris.

diversicornis, Campylomma. DNC, against aphids, 84, 240, 676, 677; against other Hemiptera, 412, 434, 676; against Lepidoptera, 111, 534, 669; toxicity of, to locusts, 658; against Oscinella frit, 46; against eggs of Saperda carcharias, 600; against Tetranychids, 8, 60, 189, 242, 676; effects of, on parasites, 434, 678; factors affecting toxicity of, 658, 676; pear and peach damaged by, 8; in sprays, 8, 46, 60, 111, 189, 240, 434, 534, 600, 658, 678; as dormant spray, 84, 242, 412, 669, 675, 676, 677; and oil emulsion, 412, 434, 534, 669, 676; and tar distillate, 434, 676, 677; as sodium salt, 675, 676, 678.

Dodder (see Cuscuta).

Dodecachlorooctahydro-1,3,4-metheno-2H-cyclobuta[cd]pentalene, in baits for fire ants, 639.

dodecella, Exoteleia (Heringia).

Dodecylguanidine Acetate (dodine acetate), effects on beneficial Mirids of sprays containing, 672.

n-Dodecylguanidine Acetate (dodine), effects of sprays containing, on arthro-

pods on apple, 20, 21, 159.

Dolichos lablab, Bruchidius incarnatus on, 249; susceptibility of varieties of, to Empoasca fabae, 463; resistance to Epilachna varivestis in strains of, 413, 414.

domestica, Musca,

domesticus, Acheta (Gryllus); Cryptotermes.

dominica, Rhyzopertha.

Dominican Republic, aphids and ants on cacao in, 144.

Doralis fabae (see Aphis). dorsalis, Acarapis; Dacus. dorsatus, Myllocerus.

Douglas Fir (see Pseudotsuga menziesii).

Dow ET-14 (see Fenchlorphos).

Dow ET-15 (see O-Methyl O-2,4,5-Trichlorophenyl Phosphoramidothioate).

Dow M361 (see O-Methyl O-2,4,5-Trichlorophenyl N-Ethyl Phosphoramidothioate).

Draeculacephala portola, on rice in Texas, 405; tests of systemic insecticides against, 405.

DRDK, 29.

Dreyfusia (see Chermes).

Drino (see Sturmia).

Drosophila, in California, 265, 573; in Ontario, 327; damaging figs and grapes, 265; in lemon packing houses, 573; dispersal of, 265; attractants for, 326; Dibrom fumigation against, 573; use of, in bioassay of insecticide residues, 339; method of rearing, 339.

Drosophila affinis, in tomato fields in

Maryland, 640.

Drosophila algonquin, in tomato fields in Maryland, 640.

Drosophila buscki, on tomato in Maryland, 640.

Drosophila falleni, in tomato fields in Maryland, 640; misidentified as D. transversa, 640.

Drosophila hydei, on tomato in Maryland,

Drosophila melanogaster, in U.S.S.R., 62; on fruits in U.S.A., 265, 640; dispersal of, 265; effects of dietary nitrogen on physiology of, 638; traps for, 265; tests of attractants for, 327; toxicity of Dibrom to, 573; susceptibility or resistance to DDT in strains of, 62, 147, 638; use of, in tests and bioassay

of insecticides, 146, 304, 397, 415, 417; labelled with ³²P, 265.

Drosophila pseudoobscura, tests of attractants for, 327.

Drosophila transversa, D. falleni misidentified as, 640.

Drosophila tripunctata, on tomato in Maryland, 640.

Drosophila virilis, in lumber-yards and houses in Japan, 622; tests of attractants for, 327; dichlorvos against, 623.

Drugs, effects of, in mixtures with rotenone against *Epilachna varivestis*, 279.

druparum, Syntomaspis (see Torymus varians).

dubia, Lypha.

dudleyi, Cryptotermes.

duplana, Rhyacionia (Evetria).

duplex, Pseudaonidia.

Dusmetia sangwani, parasitising Antonina graminis in India, 406; established in

Texas, 406; diet for, 406.

Dusts, distribution of, on tobacco leaves, 574; in relation to electrical charges on plants, 123; effects of carriers on toxicity of, 118; lipid absorptivity of, 496; applied from aircraft, 569, 574, 659, 683; (inert), killing Zabrotes subfasciatus, 99; against pests of stored maize, 51; mechanisms for removal of insect wax by, 496.

Dutch Elm Disease (see Ceratocystis

ulmi).

Dyes, marking of Eurygaster integriceps with, 115; use of, in study of deposits from oil-emulsion sprays, 221.

Dylox (see Trichlorphon). Dysaphis (see Anuraphis).

Dysdercus, in India, 173; effects of temperature on susceptibility of, to insecticides, 174.

Dysdercus fasciatus, survey of data on, on cotton in Nyasaland, 352; predator of, 429.

Dysdercus intermedius, survey of data on, on cotton in Nyasaland, 352.

Dysdercus koenigii, on cotton in India, 628; mating habits of, 628; food intake in relation to growth of, 628; uptake and excretion of schradan by, 497.

Dysdercus nigrofasciatus, survey of data on, on cotton in Nyasaland, 352.

Dysdercus peruvianus, on cotton in Peru, 559; characteristics and world distribution of, 506; insecticides against, 559.

Dysmicoccus brevipes, insecticides against, on pineapple in Brazil, 557.

Dyspessa ulula, bionomics of, on garlic in Italy, 661; measures against, 661.

E

E 605 (see Parathion).

E 605 f, containing parathion and methylparathion (q.v.), 45, 647.

E-605 Forte, containing parathion (q.v.), 111.

E 605 Staub, containing methyl-parathion (q.v.), 45.

Eacles magnifica, bionomics of, on coffee in Brazil, 120, 121; insecticides against,

Earias biplaga, on cotton in Morocco, 490; natural enemies and diseases of,

Earias cupreoviridis, in India, 36, 37; on Malvastrum tricuspidatum and Sida cordifolia, 37; bionomics and natural enemies of, 37.

Earias fabia, in Formosa, 590; in India, 36, 37, 140, 594; on cotton, 36, 37, 140, 590; on Hibiscus spp., 36, 37, 594; on other plants, 36; bionomics of, 36, 37; natural enemies of, 37; insecticides against, 590, 594.

Earias insulana, in India, 36, 37, 140, 594; in Israel, 351; in Morocco, 490; on cotton, 36, 37, 140, 351, 490; on *Hibiscus* spp., 36, 37, 594; on other plants, 37; bionomics of, 36, 37; natural enemies and diseases of, 37, 490; characteristics and world distribution of, 507; insecticides against, 351, 594; resistance to endrin in strains of. 351.

Earthworms, insecticide residues in, 641. Easter Lily (see Lilium longiflorum).

eatoni, Cylindrocopturus. ebenerianus, Trinervitermes.

Echinochloa colonum, Chloriona cubana transmitting virus diseases of, 690.

Echinochloa crusgalli, aphids and redstripe disease of, in Yugoslavia, 80.

Echthrus, 638.

Economic Entomology (see Entomology). Ectomyelois ceratoniae, on Citrus in Israel, 612.

Ecuador, banana pests in, 268, 274, 353; natural enemies of pests in, 267, 268, 274, 353.

editrix, Anomis.

Edwardsiana rosae (see Typhlocyba). Egg Albumen, in baits for Gryllotalpa, 442; in diets for locusts, 5, 6.

Egg-plant (see Solanum melongena).

Eggs. BHC residues in. 326.

Egypt, insects and virus diseases of cereals in, 684; cotton pests in, 149, 608; Trogoderma infesting stored products in, 18; termites attacking building materials in, 180; suction-trap catches of Diptera in, 30. (See also Gaza Strip.)

Ekatin (see Thiometon).

Ekatin M (see Morphothion).

Ekatox (see Parathion).

Elachertus, species of, in Britain, 52.

Elachertus agonoxenae, sp. n., parasitising Agonoxena spp. in New Guinea and Queensland, 52; introduced into Fiji against A. argaula, 52.

Elachertus pini, parasitising Dioryctria

zimmermanni in Illinois, 153.

Elaeis guineënsis, systematic list of pests of stored and field, in South-East Asia and Pacific area, 143; question of infestation of kernels of, by Ahasverus advena, 657; review of pests and diseases of, 693.

elaphus, Chremylus.

hyperparasite of Elasmus maculatus, Oiketicus kirbvi in Costa Rica, 638. Eldana saccharina, on cereals in Nigeria,

electa, Zonosemata.

electo, Colias.

Electrical Radiation, factors affecting efficacy of, against pests in stored cereals, 374.

Electrostatic Fields, effects of, on insects, 99, 123, 130, 444.

elegans, Eupelmus.

elephas, Curculio (Balaninus).

Eleusine coracana, not attacked by Laelia exclamationis, 79.

elisus, Lygus.

elliotti, Aulocara.

elliptica, Psylliodes.

Elm, pests of, in Canada, 124, 455; pests of, in Holland, 42; insects associated with, in Portugal, 298; Galerucella luteola on, in Spain, 659; Operophtera brumata on, in U.S.S.R., 190; pests of, in U.S.A., 217, 478; vectors of diseases of, 124, 217, 478. (See also Ulmus.)

Elm Spanworm (see Ennomos subsig-

narius).

elongatus, Cnemodon (Neocnemodon). elutella, Ephestia.

Elytrigia repens (see Agropyrum).

EM-923 (see Genite).

Emmalocera depressella, measures against, on sugar-cane in India, 297, 553.

Emphytus cinctus, on strawberry, rose and other plants in U.S.S.R., 361; bionomics of, 361; insecticides against, 362.

Empoasca, estimation of populations of, on vines in France, 83; on cotton in Peru, 559, 560; on sweet potato in

U.S.A., 131, 474; natural enemies of, 560; measures against, 474, 559, 560. *Empoasca biguttula*, sprays against, on

cotton in Formosa, 590.

Empoasca decedens, on vines in Italy, 661. Empoasca decipiens, bionomics of, on

vines in Italy, 661.

Empoasca fabae (in U.S.A.), on beans, 17, 328, 397, 463; on lima beans, 325; varietal susceptibility of beans to, 328, 397, 463; on potato, 325, 627, 628; statistical analysis of experiments on control of, 399; insecticides against, 325, 628.

Empoasca facialis, survey of data on, on

cotton in Nyasaland, 352.

Empoasca flavescens (on vines), in France, 372; in Italy, 660, 661; overwintering on conifers, 661; light-trap catches of, 372.

Empoasca lybica, on cotton in Sudan, 94; characteristics and world distribution

of, 507; sprays against, 94.

Empoasca pteridis, on vines in Italy, 661. Empoasca solana, on beet in California, 630; parasites of, 630.

Empoasca subrufa (see Zygina).

Empusa (see Entomophthora).

Enarmonia fortunana, on spruce in Quebec, 157.

Enarmonia griseana, in Austria and Switzerland, 100; on larch, 100; factors affecting populations of, 100.

Enarmonia ratzeburgiana, on spruce in

Quebec, 157.

Enarmonia rufimitrana, on fir in Czechoslovakia, 603; parasites of, 603.

Encarsia, parasitising Anomis texana in Peru, 560.

Encolpotis xanthoria, reared from pomegranate fruits in S. Africa, 531.

Endasys erythrogaster (parasitising sawflies), in Czechoslovakia, 246; in Germany, 682; in Yugoslavia, 138; effects of BHC against Melolontha on populations of, 682.

Endothion, against aphids, 605, 668; ineffective against *Brevicoryne brassicae*, 505; against *Ceratitis capitata*, 668; against Tetranychids, 8; effects of, on parasite of *Eriosoma lanigerum*, 312; in sprays, 312, 505, 605, 668; and

ethion, 605.

Endrin, against aphids, 412, 498, 594; against Cicadellids, 94, 395, 590, 594, 627; against other Hemiptera, 94, 174, 498, 515; against Curculionids, 16, 314, 317, 322, 417, 509, 515; susceptibility or resistance to, in strains of Anthonomus grandis, 220, 264; against other Coleoptera, 171, 173, 265, 314,

316, 451, 497, 651; against Noctuids, 16, 161, 179, 211, 269, 351, 383, 406, 413, 465, 487, 503, 514, 515, 538, 539, 590, 594, 644; resistance to, in strains Noctuids. 351: Heliothis favoured by, 323; against Pyralids, 16, 21, 79, 80, 145, 225, 280, 297, 328, 329, 469; against other Lepidoptera, 16, 121, 154, 161, 269, 335, 336, 425, 465, 503, 533, 539, 555, 641, 646, 659; crossresistance to, in strains of Estigmene acraea, 480; toxicity of, to locusts and other Acridids, 172, 174, 498; against Trypetids, 175, 268, 325, 417, 533; against other Diptera, 173, 443, 484; against Thysanoptera, 71, 515; against Cecidophyopsis ribis, 287, 302, 432; against other mites, 432, 545, 628; toxicity of, to predator of Tetranychids, 627; bees unharmed by, 432; effects of temperature on susceptibility of insects to, 174; genetic basis of resistance to, 220; use of, for control of rodents, 417; toxicity of, to fish, 304; in dusts, 71, 121, 145, 161, 314, 412, 514, 627, 644; in granules, 21, 225, 280, 325, 469; in sprays, 16, 71, 79, 80, 94, 154, 173, 174, 175, 179, 269, 287, 302, 322, 328, 330, 335, 336, 351, 383, 395, 412, 413, 425, 432, 451, 465, 484, 487, 498, 503, 509, 515, 533, 538, 545, 555, 500, 504, 641, 646, 651, 650, applied 590, 594, 641, 646, 651, 659; applied from aircraft, 425, 659; as deposit, 316; seed treatment with, 484; soil treatment with, 71, 171, 325, 417, 484, 627; treatment of sugar-cane setts with, 297; experiments on systemic activity of, 432; tainting of vegetables and fruit by, 229; stage of growth of rice in relation to effectiveness of, 328, 329; effects of, on plants, 484, 594; residues of, in sheep, 280; residues of, on plants, 225, 228, 330; factors affecting persistence of residues analysis of residues of, 91; restrictions on use of, 287; bioassay of, 397, 417; thermal isomerisation of, during gas chromotography, 561; effects of additives on, 287; and aldrin, 509; and bordeaux mixture, 651; and DDT, 413, 515; activity of, enhanced by phosphamidon, 351.

Engelmann Spruce (see *Picea engelmanni*). engelmanni, Dendroctonus.

Ennomos subsignarius, egg-masses of, on oaks and hickory in U.S.A., 418; sexual characters of, 620.

Enoclerus quadrisignatus, predacious on Pectinophora gossypiella in Texas, 338; light-trap catches of, 338.

Ent. 21195 (see 6-Bromopiperonal Chrysanthemate).

Entedon erdosi Delucchi, 306.

Entedon erdosianus (erdosi Szelenyi), n.n., 306.

Entoleter Machine, 632.

Entomology, index to literature of American economic, 496; reviews of recent research on, 90; polyglot glossary of terms in, 301; books on (see under Reviews).

Entomophthora, Macrosiphum euphorbiae infested by species of, in Maine, 570.

Entomophthora aphidis (infesting aphids), in Ontario, 459; in U.S.A., 401, 570.

Entomophthora exitialis, infesting Therioaphis maculata in Arizona, 462.

Entomophthora ignobilis, infesting aphids in U.S.S.R., 570.

Entomophthora muscae, infesting Hylemya antiqua in Quebec, 455.

Entomophthora planchoniana, infesting Macrosiphum euphorbiae in Maine, 570. Entomophthora sphaerosperma, infesting Plutella maculipennis in Britain, 198.

Enzyme Systems, in nutrition of Agrotis orthogonia, 69; activity of, in eggs of strains of Tetranychus telarius, 278; in relation to transmission of viruses by Tettigonia viridissima, 443; interaction between insecticides and, 63, 213, 248, 263; rôle of, in resistance of insects to insecticides, 148. (See also Cholinesterases.)

Eotetranychus carpini (on vines), in France, 8, 445; in Italy, 183, 663; on hornbeam, 183; bionomics of, 183; natural enemies and other factors affecting populations of, 183; measures against, 183, 445; favoured by phosphorus insecticides, 8.

Eotetranychus carpini vitis, on vines in France, 256; effects of insecticides and plant nutrition on populations of, 256.

Eotetranychus lewisi, on greenhouse plants in U.S.A., 517; sprays against 517.

Eotetranychus smithi, on vines in Japan, 452; characters of, 452.

Eotetranychus telarius (see Tetranychus). Ephedrus pulchellus, parasitising Anuraphis plantaginea in Holland, 673. ephemeraeformis, Thyridopteryx.

Ephesia nymphaea, on forest trees in Spain, 381, 660; natural enemies of, 381; virus disease of, 660.

Ephestia cautella (see Cadra).

Ephestia elutella, in warehouses in Britain, 374; traps for, 374; tests and uses of insecticides against, 374; susceptibility of, to pyrethrins, 3, 4.

Ephestia kuehniella (see Anagasta).

Ephestia sericaria (see Anagasta kuehniella).

Ephialtes brevicornis, parasitising Archips crataeganus in U.S.S.R., 15.

Ephialtes brunneus, parasitising Cydia funebrana in U.S.S.R., 112.

Ephialtes buolianae (parasite of Lepidoptera), introduced into Canada, 391; in Italy, 535; influence of diet on fecundity and life-span of, 391; effects of electrostatic fields on oviposition by, 130.

Ephialtes comstockii, parasitising Dioryctria zimmermanni in Illinois, 153.

Ephialtes detritus, in Sweden and Canada, 161; parasitising Cephus spp., 161.

Ephialtes inquisitor, survey of data on, parasitising Choristoneura murinana in France and Germany, 168.

Ephialtes pterophori, reared from Ostrinia nubilalis in Quebec, 27.

ephratae, Nasutitermes.

Epicaerus cognatus, characteristics and world distribution of, 507.

Epicometis hirta, on apple in Persia, 237. Epigonatopus plesuis, bionomics of, parasitising Macrosteles fascifrons in Ontario, 70.

Epilachna chrysomelina, characteristics and world distribution of, 507.

Epilachna paenulata, characteristics and world distribution of, 507.

Epilachna varivestis, in Quebec, 454; in U.S.A., 16, 279, 325, 413; on beans, 16, 279, 413, 454; susceptibility or resistance of bean genera and species to, 413; on lima beans, 325; on maize, 325; bionomics of, 454; effects of temperature on, 454; survival of, in nitrogen atmospheres, 416; toxicity of culture filtrates of Myrothecium roridum to, 572; tests and uses of insecticides against, 16, 148, 271, 279, 325.

Epilachna vigintioctomaculata, chemical factors in relation to resistance of

solanaceous plants to, 608.

Epilachna vigintioctopunctata, on eggplant and potato in India, 497; bionomics of, on vegetables in Pakistan, 88; chemical factors in relation to resistance of solanaceous plants to, 608; toxicity of insecticides to, 497.

Epinotia nigricana (see Eucosma). Epiphyas, Australian species of, 232.

Epiphyas eucyrta, 232.

Epiphyas postvittana, in Australia, 232; characteristics and world distribution of, 507; synonymy of, 232.

Epitrix cucumeris (on potato), in India,

500; in Maryland, 325; insecticides against, 325.

Epitrix parvula, on jute in British Guiana.

27; insecticides against, 27.

Epitrix tuberis, on potato in British Columbia, 424, 526; soil treatments against, 526.

Epiurus (see Ephialtes).

EPN, against Coccids, 120, 557; against Diptera, 623; against Lepidoptera, 9, 21, 145, 469, 646; against mites, 626, 643, 645; metabolism of, in insects, 64; in dust, 145; in granules, 469; in sprays, 9, 120, 557, 626, 643, 645, 646; toxicity of DDT increased by, 21; and oil emulsions, 120, 557; antagonistic action of sesamex with, 63.

equestris, Haplodiplosis: Merodon

(Lampetia).

erdosi Delucchi, Entedon.

erdosi Szelenyi, Entedon (see E.erdosianus). Erechtites, development of Tyria jacobaeae on species of, 260.

Erechtites arguta, 260. Erechtites hieracifolia, 260.

Eremnus cerealis, bionomics of, on vines in S. Africa, 650.

Eremnus setulosus, bionomics of, on vines in S. Africa, 650.

erichsonii, Laricobius; Pristiphora.

Erigeron canadensis, Tetranychus telarius overwintering on, in Maryland, 477.

Eriococcus campbelli, question of introduction of, from Australia into New Zealand for control of Leptospermum scoparium, 588.

Eriococcus leptospermi, on Leptospermum scoparium in association with E. orariensis, 588; introduced into New

Zealand, 588.

orariensis, introduced into Eriococcus New Zealand from Australia, 587; use of, in biological control of Leptospermum scoparium, 587; resistance of L. ericoides to, 588; bionomics of, 587; fungus disease of, 587, 588.

Erioischia (see Hylemya).

on Hibiscus esculentus Eriophyes, in Brazil, 357.

Eriophyes avellanae (see Phytoptus).

Eriophyes insidiosus, transmitting peach mosaic in U.S.A., 71.

Eriophyes merwei (see Aceria).

Eriophyes padi, on plum in Austria, 85. Eriophyes phloeocoptes (see Aceria)

Eriophyes pyri, in S. Africa, 83; in U.S.S.R., 191; on apple and quince, 83; on pear, 83, 191; bionomics of, 191; measures against, 191.

Eriophyes pyri mali, bionomics of, on apple in U.S.S.R., 343.

Eriophyes ribis (see Cecidophyopsis).

Eriophyes vitis (on vines), in S. Africa, 83, 298; in Germany, 71; transmission of viruses by, 71, 303; acaricides against. 298; effects of sprays against Planococcus citri on populations of, 298.

Eriopis connexa, predacious on larvae of Eulia loxonephes in Argentina, 144.

riosoma lanigerum (on apple), in California, 481; in China, 591; in Czechoslovakia, 678; in France, 312, Eriosoma 313; in Japan, 35, 36; in Nova Scotia, 387, 388; factors affecting dispersal behaviour of first-instar nymphs of, 481, 482; natural enemies of, 312, 388, 591, 678; tests and uses of insecticides against, 35, 36, 312, 313.

Eritrea, new parasite of Coccids in, 491.

eritreaensis, Coccophagus.

erminea, Dicranura.

Ernestia consobrina, bionomics of, parasitising Mamestra brassicae in U.S.S.R.,

Ernobius abietis, bionomics of, on spruce in Italy, 136; fungal symbionts of, 136; parasites of, 136.

Erodium moschatum, Aphodius tasmaniae on, as component of pastures in S.

Australia, 504. eruditus, Cheyletus.

Erysit (see Chlorfenson).

Erythrina, larvae of Eupterote fabia reared on, 37.

erythrocephala, Acantholyda.

Erythrocera scutellaris, parasitising Hyponomeuta cognatellus, 87; experiments with Galleria mellonella and larvae of,

erythrogaster, Endasys (Stylocryptus). Erythroneura comes, on vines in Bulgaria,

110.

Erythroneura flammigera, on vines in Italy, **661**.

Erythroneura rhamni, on vines in Italy, 660, 661; overwintering on blackberry, 661; misidentified as E. simplex, 660.

Erythroneura simplex, on vines in France, 372; light-trap catches of, 372; E. rhamni misidentified as, 660.

Erythroneura tiliae, on vines in Italy, 661.

Eserine, effect of, in insects, 62.

Estigmene acraea (on cotton in U.S.A.), other food-plants of, 480; natural enemies of, 638, 639; used in tests of insecticides, 480, 634; insecticides against, 16, 480, 508, 516; question of resistance to chlorinated hydrocarbons in strains of, 480.

Estonia, outbreak of Plutella maculipennis in, 430.

esula, Catabena.

Ethanol, in bait for Drosophila, 327.

Ether, anaesthetisation of insects with

carbon dioxide and, 356.

Ethion, against aphids, 279, 324, 604; Aphis pomi favoured by, 394; ineffective against Phylloxera vitifoliae in galls, 668; against Coccids, 561, 667; toxicity of, to Coleoptera, 338, 603; against Lepidoptera, 21, 337, 394, 398, 466, 669; against Tetanops myopaeformis, 402; tests and uses of, against Trypetids, 268, 668; against Sminthurus viridis, 34; against Tetranychids, 281, 324, 394, 603, 604; susceptibility or resistance to, in strains of Tetranychids, 510, 512; toxicity of, to bees, 412; toxicity of, to Syrphid larvae, 605; tests of toxicity of, to other insects, 603; anticholinesterase activity and mammalian toxicity of, 604; effects of, on plants, 281, 394; in granules, 402; in sprays, 34, 268, 279, 324, 337, 394, 398, 466, 604, 605, 667, 668, 669; soil treatment with, 402; persistence of deposits of, 394; residues of, on Citrus fruits, 561; spectrophotometry of residues of, 561; and DDT, 21; and endothion, 605; and methyldemeton, 604, 605; and oil emulsion, 604, 667.

Ethyl 1-Chloro-2-propyl-2,2-dichlorovinyl Phosphate, toxicity of, to Estigmene

acraea, 480.

Ethyl-DDD, against Coleoptera, 500; against Lepidoptera, 26, 161, 222; against Trypetids, 98, 268, 668, 669; Bryobia rubrioculus favoured by, 523; in dust, 161; in sprays, 26, 98, 222, 268, 523, 668, 669; treatment of sacks with, 500; and Dibrom, 222; and lead arsenate, 268; and malathion, 161, 222. Ethyl-Gusathion (see Azinphos-ethyl).

Ethyl Guthion (see Azinphos-ethyl). Ethyl 3-Indoleacetate, in honeydew of 629; *Toxoptera* graminum,

chromatography of, 629.

Ethylmercury Acetate (see Ceresan). Ethylmercury 2,3-Dihydroxypropyl Mer-

captide (see Ceresan).

Ethylmercury-p-toluenesulphonanilide, in seed treatments of cotton with insecticides, 352.

Ethyl p-Nitrophenyl tert.-Butylphosphonate, anticholinesterase activity and toxicity of, to Musca domestica, 407.

Ethyl p-Nitrophenyl Ethylphosphonate, anticholinesterase activity and toxicity of, to Musca domestica, 407.

Ethylquinone, discharge of, by *Tribolium* castaneum, 621.

Ethylene Chlorobromide, fumigant toxicity of, to Coleoptera, 267.

Ethylene Dibromide, alone or in fumigant mixtures against pests of stored products, 118, 267, 274, 579; ineffective against Tribolium castaneum, 118; as soil fumigant against Lachnosterna consanguinea, 171; factors affecting toxicity of, 274; effects of, on seed germination, 579; sorption and retention of, by fruits, 574; determination of, in air, 496.

Ethylene Dichloride, alone or in fumigant mixtures against pests of stored products, 118, 267, 274, 579; factors affecting toxicity of, 274; effects of, on

seed germination, 579.

Ethylene Oxide, fumigant toxicity of, to

Coleoptera, 267.

Etiella behrii, on lucerne in S. Australia. 233: parathion sprays against, 234.

Etiella zinckenella, on leguminous plants in Saskatchewan, 529; in Japan, 496; cold-hardiness of overwintered larvae of. 424.

Eubadizon gracile, parasitising Evagora starki in Canada, 586.

Eucalyptus, pests of, in Australia, 32, 33, 231, 232, 504, 552; termites attacking seedlings of, in Nigeria, 294; resistance of, to attack by Psammotermes spp., 180.

Eucalyptus acmenioides, pests of, in New South Wales, 232.

Eucalyptus blakelyi, Perga affinis on, in

Australia, 504. Eucalyptus camaldulensis, Perga affinis on, in Australia, 504; termites attacking seedlings of, in Nigeria, 294; relation of, to pollination of lucerne, 234.

Eucalyptus dives, 33.

Eucalyptus intermedia, starch content related to infestation of, by Lyctus brunneus in Queensland, 589.

Eucalyptus leucoxylon, relation of, to pollination of lucerne in S. Australia. 234.

Eucalyptus melliodora, Perga affinis on, in Australia, 504.

Eucalyptus punctata, starch content related to infestation of, by Lyctus brunneus in Queensland, 589.

Eucalyptus radiata, 33.

Eucalyptus robertsoni, 33.

Eucalyptus saligna, pests of, in New South Wales, 232, 233.

Eucalyptus tesselaris, starch content related to infestation of, by Lyctus brunneus in Queensland, 589.

Eucarcelia rutilla (see Carcelia obesa).

Eucosma nigricana, on fir in Czechoslovakia, 602; parasite and hyperparasites of, 602.

eucyrta, Epiphyas.

Euderus agromyzae, sp. n., parasitic on Melanagromyza obtusa in India, 497.

Eugenia cumini (jambolana), Chrysocraspeda olearia on, in India, 88.

eugenii, Anthonomus,

Eugenol, as attractant for Popillia japonica,

Eulaelaps stabularis, in stored grain in Canada, 154.

Eulecanium corni, on vines in Austria, 85; in France, 683; on fruit trees in New York, 470; on apple in Poland, 676; bionomics of, on plum, persimmon, filbert, peach and other trees in U.S.S.R., 13; natural enemies of, 683; dormant sprays against, 470, 676.

Eulecanium coryli, natural enemies of, in

France, 683.

Eulecanium pruinosum, identity of Aphycus sp. parasitising, in California, 491. Euleia heraclei, characteristics and world

distribution of, 507.

Eulepida mashona, bionomics of, in pastures in S. Rhodesia, 38; soil treatments against larvae of, 38.

Eulia cupressana (see Lozotaeniodes).

Eulia dimorpha, on Citrus, 558; in Argentina and Brazil, 558; sprays against, 558.

Eulia loxonephes, bionomics of, on apple in Argentina, 144; natural enemies and virus disease of, 144; effects of sprays on populations of, 144.

Eulia sphaleropa, on fruit trees in Argen-

tina, 299.

Eulophus larvarum, parasitising Operophtera brumata in Europe, 190, 378; introduced into Canada, 378.

Eulophus longulus (see Pnigalio).

Eumargarodes laingi, on sugar-cane in Australia, 265; on grasses in U.S.A., 265; soil treatments against, 265.

Eumea caesar (see Zenillia).

Eumegastigmus pistaciae, on pistachio in Turkey, 609.

Eumerus obliquus, reared from pomegranate fruits in S. Africa, 531.

euonymellus, Hyponomeuta.

Euonymus, Hyponomeuta cognatellus on, in Finland, 194; Aonidiella orientalis on, in Pakistan, 491; H. euonymellus not occurring on, 194.

Euonymus europaeus (primary food-plant of Aphis fabae), in Bulgaria, 536, 537;

in Czechoslovakia, 305.

Euonymus fortunei, Abraxas miranda on, in China, 503.

Euonymus japonicus, Abraxas miranda on, in China, 503.

Euonymus latifolius, as primary foodplant of Aphis fabae in Bulgaria, 536.

Eupelmus atropurpureus, attacking parasitised Cephus spp., 161; introduced into Canada, 161; in U.S.S.R., 161.

Eupelmus elegens, parasitising Rhyacionia buoliana in Argentina, 144.

Eupelmus urozonus, in France, 435; in Îtaly, 438, 535; parasite of Dacus oleae, 435, 438; parasitising Pelatea festivana,

Euphorbia marginata, Eotetranychus

lewisi on, in U.S.A., 517.

Euphorbia pulcherrima, Eotetranychus lewisi on, in U.S.A., 517; effects of acaricides on colour of, 517.

euphorbiae, Aphthona; Macrosiphum.

Euphyllura olivina, bionomics of, on olive in Turkey, 81; sprays against, 81.

associated Euphysothrips subramanii, with abnormal growth of coffee in India, 448.

Euplectrus laphygmae, parasitising Spodoptera exempta in Kenya, 531.

Euproctis chrysorrhoea, on elm in Holland, 42.

Euproctis pseudoconspersa, in Japan, 450; effects of crowding on larvae of, 450.

Eupterote fabia, on coffee and associated shade trees in India, 37; bionomics and feeding habits of, 37.

eurilochus, Caligo.

eurinus, Rogas.

European Pine Shoot Moth (see Rhyacionia buoliana).

European Red Mite (see Panonychus ulmi). Eurydema oleraceum, characteristics and world distribution of, 507.

Eurydema ornatum, bionomics of, on cabbage and cauliflower in Rumania,

57; insecticides against, 57.

Eurygaster (on cereals), in U.S.S.R., 363; egg-parasites of species of, in Morocco and Middle East, 654, 655; toxicity of insecticides to, 382; FAO report on control of, 300.

Eurygaster amerinda, on wild oats and other plants in California, 520; bionomics and ecology of, 520; question of infestation of cereals by, 520.

Eurygaster austriaca, on cereals in Morocco, 382, 653, 654; diapause in, 653; new parasite reared from eggs of,

654.

Eurygaster hottentotta, on cereals in Morocco, 382, 653; diapause in, 653.

Eurygaster integriceps (on cereals), in Bulgaria, 109; in Persia, Turkey and other countries of Middle East, 43, 428, 429, 485, 653, 655, 656; in U.S.S.R. 13, 14, 115, 363, 446, 543; on other plants, 429; baking quality of flour affected by, 109; extent of injury by, 446; adult habits of, 429; factors affecting development of, 14, 601, 655, 656; physiological changes in, 656; natural enemies of, 43, 485, 601, 653, 655: characteristics and world distribution of, 507, 543, 601; marking of, 115; labelled with 182Ta, 429; tests of bacteria and fungi against, 115; insecticides against, 115; FAO report on control of, 300; booklet on, 543.

Eurygaster maura, on cereals in Middle East, 485; in Morocco, 382; natural

enemies of, 485.

Eurygaster minidoka, on wild oats and other plants in California, 520; bionomics and ecology of, 520; question of infestation of cereals by, 520.

Eurysthaea scutellaris (see Erythrocera).

eurytheme, Colias.

Eurytoma (Bruchophagus), in S. Australia, 234; in Bulgaria, 60; key to species of, infesting seeds of leguminous plants in Hungary, 305; nomenclature of species of, in U.S.S.R., 564; on lucerne, 60, 234; status of, 305.

Eurytoma amygdali, bionomics of, on almond in Bulgaria, 110; measures

against, 110.

Eurytoma coluteae, on Colutea arborescens in Hungary, 306.

Eurytoma evolans, sp. n., on Medicago minima in Hungary, 306.

Eurytoma gibba, sens lat. (on lucerne), in S. Australia, 234; in Bulgaria, 60.

Eurytoma gibba Boh. (see E. platyptera). Eurytoma kolobovae (on Lotus corniculatus), in Hungary, 305; in U.S.A., 564; characters of, 564.

Eurytoma morio, as parasite and hyperparasite of Hylesinus fraxini in Czecho-

slovakia, 188.

Eurytoma onobrychidis, on Onobrychis viciifolia in Hungary, 305.

Eurytoma ononis, on Ononis spinosa in Hungary, 306.

Eurytoma pini (parasitic on Rhyacionia rigidana), in Ohio, 20; in Ontario, 157; factors affecting parasitism by, 158; characters of, 158.

Eurytoma (Bruchophagus) platyptera (on Trifolium pratense), in Hungary, 306; in U.S.A., 564; characters of, 564.

Eurytoma plotnikovi, on pistachio in Turkey, 609.

Eurytoma roddi (on Medicago spp.), in Hungary, 306; in U.S.A., 564; characters of, 564.

Eurytoma rosae (parasitising Dacus leoea),

in France, 435; in Italy, 308, 438, 532; winter mortality of, 308.

Eurytoma sophorae, on Sophora japonica in Hungary, 306.

eurytus, Tetrastichus (Geniocerus).

Eutetranychus banksi, on castor in India, 627; predator of, 627.

Eutetranychus orientalis, on Citrus in Aden, 80.

eutettixi, Polynema.

Eutinobothrus brasiliensis, characteristics and world distribution of, 507.

Eutrombidium rostratum (trigonum), predacious on grasshopper in India, 500. Euxoa radians, on lucerne in S. Australia, 234.

Euxoa segetum (see Agrotis).

Evagora, status of, 121.

Evagora canusella, on pines America, 121.

Evagora condignella, on pines in N. America, 121.

Evagora milleri, on pines in N. America, 121, 124, 221, 222; parasites and hyperparasites of, 124, 221, 222; effects of sprays on parasites of, 222; integration of biological and chemical control of, 222.

Evagora moreonella, on pines in N. America, 121.

Evagora starki, on pines in N. America, 121, 586; parasites of, 586.

evanescens, Trichogramma.

evasus, Myllocerus.

Evetria (see Rhyacionia).

Evipal (see 5-(1-Cyclohexen-1-yl)-1,5dimethylbarbituric Acid).

evolans, Carcelia; Eurytoma.

Exapate congelatella, on berry fruits in U.S.S.R., 360; DDT sprays against, 360.

exareolata, Nythobia (Horogenes). exclamationis, Laelia.

exempta, Spodoptera (Laphygma).

Exenterus ictericus, parasitising Monoctenus juniperi in Holland, 302.

Exetastes cinctipes, parasitising Mamestra brassicae in U.S.S.R., 244; bionomics of, 245.

exigua, Campogaster; Spodoptera (Laphygma).

exiguae, Hyposoter.

exitiosa, Aegeria (Sanninoidea).

exitiosus, Nasutitermes.

Exora collaris, on onion in Turkey, 82.

Exorista (see Tachina).

Exoteleia dodecella (on pines), introduced into Canada, 391; bionomics of, in Czechoslovakia, 246; in U.S.S.R., 116; attacked by Ephialtes buolianae, 391; DDT sprays against, 246.

exsecta, Formica. externus, Acarapis. extranea, Anabrolepis.

F

Faba vulgaris (see Beans, Broad). fabae, Aphis (Doralis); Empoasca. fabia, Earias; Eupterote. fabricator, Cratichneumon.

Fac 10 (see O,O-Dimethyl S-Isopropylcarbamoylmethyl Phosphorodithioate). Fac 20 (see O,O-Diethyl S-Isopropylcarbamoylmethyl Phosphorodithioate).

facialis, Empoasca. falcatus, Pygostolus. fallacis, Typhlodromus. fallax, Chauliops. falleni, Drosophila. falli, Conoderus.

farinae, Tyroglyphus (see Acarus siro).

farris, Acarus.

fasciata, Prospaltella.

fasciatus, Dysdercus; Oncopeltus.

fasciculata, Oncopera. fasciculatus, Araecerus. fascifrons, Macrosteles. fatigans, Culex pipiens. fecundus, Ooencyrtus.

Felt, use of, in rearing insects, 339.

feltella, Acrobasis.

Feltia subterranea, on tobacco in Brazil, 357; soil treatments against, 357.

femoralis, Adelencyrtus. femorata, Spilochalcis. femurrubrum, Melanoplus.

Fenchlorphos, against Aphis gossypii, 279; against Lymantria dispar, 514; against Frankliniella occidentalis, 569; toxicity of, to Rhagoletis pomonella, 268; absorption, metabolism and excreton of, by Leucophaea maderae, 248; against Tetranychus telarius, 217; in dust, 217; in sprays, 213, 217, 279, 514, 569; tests of, against pests of stored products, 213.

fenestralis, Nythobia (Angitia, Horogenes). Fenson, in sprays against Panonychus

ulmi, 444, 685.

Fenthion, against aphids, 505, 645; ineffective against Brevicoryne brassicae, 74; against other Hemiptera, 408, 515; against Curculionids, 180, 408, 412, 509; against Diptera, 402, 668, 669; diverse effects of, against Lepidoptera, 9, 409, 411, 413, 466, 469; against Thysanoptera, 477; against Tetranychus telarius, 645; metabolism of, and its derivatives and analogues in rats, insects

and cotton plants, 475, 476; anticholinesterase activity of, 475; toxicity of, to beet, 402; in dust, 409; in sprays, 9, 74, 408, 411, 412, 466, 477, 505, 509, 515, 645, 668, 669; soil treatment with, 180, 402; residues of, in milk, 266; labelled with ³²P, 475; and DDT, 412, 413.

Fenusa pusilla, on birch in New York, 19; tests of systemic insecticides against, 19.
 Ferbam, in spray mixtures against Cydia pomonella, 20; in mixtures with acari-

cides, 264, 324, 439, 523.

Feronia melanaria, bionomics of, on strawberry in Britain, 670; sampling methods for, 671.

Ferric Citrate, effects of, on fecundity of *Aptesis basizona*, **155**.

Ferrisiana virgata, on Citrus in Aden, 80.

Ferrous Sulphate, effects of sprays of, on beneficial arthropods, 36; and chlorfenson, 36; and parathion, 36.

ferrugalis, Udea (Pionea). ferrugineus, Cryptolestes.

festivana, Pelatea.

Festuca, stem-boring Diptera on, in Britain, 488.

Festuca elatior, Thymelicus lineola on, in Ontario, 162.

Festuca pratensis, Oscinella frit on, in Germany, 45.

ficus, Aceria; Chrysomphalus. Field Beans (see Beans, Broad).

Fig, mosaic disease of, in Australia and Britain, 70; mosaic disease and pests of, in California, 70; 265; mosaic disease of, in Italy, 70; Ceratitis capitata on, in Yugoslavia, 239, 240; Aceria ficus transmitting mosaic of, 70; (dried), development of Necrobia rufipes on, 436.

Fig Paste, use of, in labelling *Drosophila* with ³²P, **265**.

Fiji, Hymenia recurvalis on Amarantus viridis in, 88; Nacoleia octasema on banana in, 353; pests of coconut in, 52, 81; parasites introduced into, 52, 354.

Filbert (see Corylus).

filiformis, Coeloides.

Fimbriaphis fimbriata, on strawberry in British Columbia, 529.

fimbriata, Fimbriaphis.

Finland, apple pests in, 103, 194; Pristiphora monogyniae on plum in, 103; Hyponomeuta euonymellus on Prunus padus in, 194; Lygus spp. on beet in, 103; pests of crucifers in, 103, 104, 674; Calligypona pellucida on cereals and other plants in, 102, 103; Sitona spp. on peas and leguminous forage crops in, 41, 102; S. sulcifrons on Phleum pratense in, 41; forest pests in, 101, 194; form of H. padellus associated with Sorbus aucuparia in, 194; H. cognatellus on Euonymus in, 194; Coleoptera in timber in, 42, 101; Plutella maculipennis in, 198, 430; review of aphid vectors of plant viruses in, 101; beneficial insects in, 195.

finlandicus, Typhlodromus.

Fir, Balsam (see Abies balsamea).

Fir, Douglas (see Pseudotsuga menziesii).

Fir, Fraser (see Abies fraseri). Fir, Grand (see Abies grandis).

Fir. Pacific Silver (see Abies amabilis).

Fir, Silver (see Abies alba).

Fir, Subalpine (see Abies lasiocarpa).

fiscellaria, Lambdina.

Fish, ecology of, in relation to the use of insecticides, 262; toxicity of insecticides to, 12, 304. Fish Meal, in baits for Gryllotalpa, 442;

Dermestes reared on, 494.

Fish Oil, in baits for fire ants, 464.

fitchii (Sand.), Rhopalosiphum.

fitchii auct., Rhopalosiphum (see R. padi). flammea, Panolis.

flammigera, Erythroneura.

Flannel, use of, in tests of textile protectants, 290, 587.

flava, Hoplocampa; Potamyia; Scaptomvzella.

flaveolatum, Agrypon.

flavescens, Empoasca; Sitona.

flavicauda, Polyblastus (Scopiorus).

flaviceps, Physcus.

flavicinctus, Cirrospilus.

flavicollis, Kalotermes (Calotermes).

flavicornis, Physcus.

flavida, Colaspis.

Apion flavipes, Anthrenus; (see A. dichroum); Reticulitermes.

flavoorbitalis, Cremastus.

flavus, Microterys.

Flax, Bruchids on, in Algeria, 249; pests of, in Poland, 364, 366; Heliothis spp. on, in Oueensland, 588; Aphthona euphcrbiae on, in Rumania, 608; forecasting abundance of pests of, in U.S.S.R., 15; protection of beet with shelter crop of, 606.

fletcherana, Cydia (Laspeyresia).

Floods, effects of: on larvae of Aphodius tasmaniae, 166; on immature stages of Lepidoptera, 182, 569; on infestation of cotton by Diparopsis watersi, 487.

floralis, Hylemya (Chortophila); Nemorilla. Florida, Citrus pests in, 409, 552; insects and diseases of pecan in, 424; pests of maize in, 275, 327, 512; Chloriona

orizicola and hoja blanca disease of rice in, 405; new sawfly on Pinus 462; Micrutalis elliottii densa in. malleifera transmitting curly-top virus of tomato, tobacco and other plants in, 635, 636; Eotetranychus lewisi on poinsettia in, 517; other insects in, 495, 507, 567.

floridensis, Ceroplastes.

Flour, Coleoptera in, 225, 315, 566; effect of grain infestation by Eurygaster integriceps on quality of, 109; treatment of, with insecticides against pests, 225; residues of insecticides in, 270, 483; in baits for insects, 388, 464; in diets for Tenebrionids, 89, 184; use of, as wetting agent in sprays, 463.

Flour Beetle, Confused (see Tribolium

confusum).

Flour Mills, insects infesting, 657, 658,

694: fumigation of, **694**.

Fluorescent Powder, in experiments on feeding habits of locusts, 164. (See also Dyes.)

Fluoroacetamide, in sprays against mites, 287, 432, 671, 672; toxicity of, to black currants, 672; (sodium salt), in sprays against Brevicoryne brassicae, 74.

Folic Acid, in diet of Corcyra cephalonica,

Folidol E 605, containing parathion (q, v), 297.

forbesi, Amphorophora; Cerosipha (Aphis). Forest Fires, effects of: on infestation of spruce by Monochamus oregonensis, 457; on populations of Phasmatids and their natural enemies, 231, 232,

Forest Pests, in Angola, 608, in Argentina, 230; in Australia, 32, 33, 231, 232, 233, 504, 552, 589; in Austria, 100; in Belgium, 85, 606; in Britain, 550; in British Honduras, 76; in Canada, 24, 25, 73, 74, 107, 121, 122, 123, 124, 125, 126, 127, 130, 140, 157, 158, 163, 168, 169, 185, 206, 207, 208, 209, 210, 258, 261, 341, 342, 385, 387, 391, 392, 393, 394, 395, 423, 424, 455, 456, 457, 458, 459, 460, 474, 522, 529, 532, 580, 586, 619, 692, 693; in Canary Islands, 659; in China, 501, 502, 593, 623, 624; in Czechoslovakia, 142, 187, 196, 246, 283, 600, 601, 602, 603, 683, in Denmark, 674; in Finland, 101, 194; in France, 84, 168, 194, 423, 606; in Germany, 10, 11, 44, 106, 107, 140, 141, 168, 185, 186, 248, 250, 251, 252, 253, 375, 376, 380, 386, 494, 682, 683; in Holland, 185; in Hungary, 307; in India, 88, 448, 552, 620; in Italy, 136, 137, 535, 615; in Java, 625; in Malaya, 296; in Mongolia, 116; in Papua, 300;

in Portugal, 298; in Puerto Rico, 257; in S. Rhodesia, 81; in Spain, 353, 355, 381, 551, 659, 660; in Sweden, 185, 426; in Switzerland, 46, 100, 186; in Turkey, 355; in Uganda, 496; in U.S.S.R., 12, 111, 112, 114, 115, 116, 190, 191, 242, 243, 244, 344, 355, 368, 541, 596, 597, 598, 599, 620; in U.S.A., 19, 20, 23, 24, 73, 74, 121, 122, 123, 124, 125, 129, 149, 150, 153, 169, 185, 208, 209, 210, 212, 220, 221, 222, 259, 271, 276, 327, 336, 337, 339, 340, 355, 385, 386, 399, 400, 407, 410, 411, 413, 418, 419, 458, 460, 461, 462, 467, 474, 478, 481, 495, 513, 532, 552, 619, 632, 641, 642; in Yugoslavia, 31, 54, 55, 181, 241, 367-371, 535, 536; factors affecting populations of, 231, 376, 494; Formica spp. in relation to, 615, 616; question of rôle of ingested antibacterial substances in protection of, against bacteria, 580; comparison of survey methods for, 456; biological control of, in relation to types of tree, 199; review of biological control of, 692, 693; portable spray chamber for, 248.

Forficula auricularia, predacious on Phytodecta olivaceus in Britain, 30.

Formaldehyde, inactivation of plant virus by, 133.

Formica, predacious species of, in Italy, 615, 616; classification of groups of,

Formica aquilonia, described from Britain, 104; members of group of, in Germany, 104; in forests in Italy, 615.

Formica congerens (see F. nigricans).

Formica exsecta, members of group of, in

Germany, 104.

Formica lugubris, members of group of, in Germany, 104; in forests in Italy, 615, 616; predacious on Thaumetopoea pityocampa, 615; transplantation of colonies of, 615, 616.

minor pratensoides (see F. Formica

nigricans).

Formica nigricans, members of group of, in Germany, 104; synonymy of, 104.

Formica polyctena, in Germany, 104, 105, 141, 251, 682; in Italy, 615; predacious on Lepidoptera, 251; predacious on Pristiphora abietina, 141, 682; population fluctuations of, 105; destroyed by woodpeckers, 104, 141; diets for, 105; synonymy of, 104.

Formica pratensis, auct. (see F. nigricans). Formica rufa, in Germany, 104; in forests in Italy, 615; members of group of, 104. Formica rufa rufopratensis minor (see F.

polyctena).

Formica truncorum, members of group of, in Germany, 104.

formicarius, Thanasimus. formiciformis, Theocolax.

Formosa, cotton pests in, 590; Rhynchocoris humeralis on Citrus in. 591: Dacus cucurbitae on cucurbits in. Tryporyza incertulas on rice in, 328; identity of *Physcus* sp. parasitising

Coccids in, 491. formosae, Frankliniella. fornicata, Phytodecta. fornicatus, Xyleborus.

fortunana, Enarmonia (Zeiraphera).

Fossae, use and treatment of, for storage of grain in Malta, 177, 178.

Fostion (see Dimethoate).

Foulbrood (of honey bees), 332.

foveicollis, Aulacophora (Rhaphidopalpa). Fowls, Tenebrionids infesting food of, 420; BHC residues in tissues and eggs of, 326; toxicity of azinphos-methyl to, 475: question of residues of azinphosmethyl in foods for, 475.

fragaefolii. Capitophorus (Passerinia.

Pentatrichopus).

Fragaria, Capitophorus jacobi on species of, in U.S.A., 23. (See also Strawberry.) Fragaria vesca, dodder transmitting virus from clover to, in Germany, 441; experiments with aphids and diseases of strawberry and, 106, 107, 192, 441, 546.

fragariae, Ancylis (see A. comptana): Macrosiphum (Sitobion); Steneotarson-

France, pests of beans in, 604, 605; aphids and virus disease of beet in, 605, 669, 670; Coleoptera on cruciferous oil-seed crops in, 373; Hepialus lupulinus on strawberry, cereals and garlic in, 7: pests of maize in, 371, 372, 433, 666; Calligypona pellucida on grasses in, 373; orchard pests in, 8, 83, 84, 255, 256, 257, 312, 379, 433, 434, 435, 550, 604, 664, 665, 666, 668, 669; Dacus oleae on olive in, 435; pests of vines in, 8, 9, 83, 84, 256, 434, 445, 494, 667, 668; Homoptera in relation to virus diseases of vines in, 372; forest pests in, 83, 84, 168, 423, 606; pests of chestnut in, 84, 435; pests of walnut in, 151; review of Coleoptera infesting wood of walnut in, 194; Tetranychus cinnabarinus on 550; carnation in, Thomasiniana lavandulae on lavender in, 436, 606; Xyleborus morigerus introduced into, on orchids from Britain, 625; Eulecanium coryli and E. corni in, 683; Plutella maculipennis in, 430; Reticulitermes lucifugus santonensis in, 649;

beneficial insects in, 7, 84, 151, 168, 257, 312, 313, 372, 373, 380, 390, 391, 414, 435, 491, 665, 683; Opius concolor introduced into, against D. oleae, 435; introduction of parasites of Quadraspidiotus perniciosus into, 379, 433, 434, 435, 681.

frangulae auct., Aphis (see A. gossypii). Frankliniella (on cotton), in Brazil, 645; in Greece, 352; in U.S.A., 408, 515; insecticides against, 352, 408, 515, 645;

Frankliniella formosae, on wheat China, 625.

Frankliniella fusca (in U.S.A.), damaging groundnuts, 151; on cotton, 467, 515; insecticides against, 151, 467, 515.

Frankliniella occidentalis (in U.S.A.), on cauliflower, 508; on cotton, 409; on lettuce, 569; on onion, 477; on potato, 474; insecticides against, 409, 474, 477, 508, 569,

Frankliniella tritici, on roses in Maryland, 212; traps for, 212; insecticides against,

212.

Frankliniella vaccinii, on blueberry in New Brunswick, 71; soil-surface treatments against, 71.

Fraser Fir (see *Abies fraseri*). fraxini, Hylesinus (Leperisinus). frit, Oscinella (Oscinis, Oscinosoma).

Frit Fly (see Oscinella frit).

Froghopper, Sugar-cane (see Aeneolamia varia saccharina).

frontalis, Dendroctonus; Magdalis; Megachile.

frontatus, Microterys.

Fructose, in relation to feeding response of wireworms, 528.

frugiperda, Spodoptera (Laphygma).

frumenti, Diocalandra. frustrana, Rhyacionia.

fuliginosa, Waterstonia.

fuliginosus, Dahlbominus.

fulvicollis, Opius.

fulvicornis, Perilampus.

fulvipes, Xenoschesis.

fulvomaculatus, Calocoris.

fulvus, Physcus.

fumiferana, Choristoneura,

fumiferanae, Apanteles; Glypta.

Fumigants, toxicity of, in relation to respiratory rate of Coleoptera, 267; method of testing pine-resin vapours as, against bark-beetles, 223; apparatus for treating insects with radioactive, 356; sorption of, by stored cereals, 215, 554; effects of, on germination of stored seed, 579; air-flow indicator in sampling of, 424; determination of concentration in mixtures of, 619. fumipennis, Lypha.

fundella, Argyresthia.

funebrana, Cydia (Grapholitha, Laspeyresia).

funesta, Oxythyrea.

Fungi, as symbionts of insects, 136, 650; use of, in measurement of activity of factors in maize resistance to Ostrinia nubilalis, 22, 68; toxicity of filtrates of, to culture *Epilachna* varivestis, 572. (See also Mushrooms.)

Fungi, Entomogenous, infesting aphids, 401, 459, 462, 569, 570, 610; infesting Coccids, 54, 587, 588; question of insecticides in relation to susceptibility of Eurygaster integriceps to, 115; infesting Membracid, 636; infesting Coleoptera, 166, 187, 196, 284, 377, 447, 541, 619; relation of, to development of Ahasverus advena, 657; mechanism of penetration by, into larvae of Leptinotarsa decemlineata, 196; infesting Hylemya antiqua, 455; infesting Lepidoptera, 44, 164, 198, 407, 538, 638; infesting sawflies, 210; infesting termites, 428; laboratory experiments with, 44, 115, 284; field use of, 115, 164, 541; effects of DDT used in combination with, 447.

Fungi, Injurious, resistance in wheat varieties to Cecidomyiid and, 65; relation of insects to, 124, 190, 193, 217, 275, 323, 398, 414, 415, 651, 652; inhibition of, by secretion from Scaptocoris divergens, 565; combined sprays against mites and, 264, 324; effects of sprays against, on beneficial arthropods, 159; relation of, to seed treatment with insecticides, 398; treatment of logs against infection by, 532. (See also

Sooty Mould.)

fur, Ptinus.

furcellata, Cantheconidea.

Furniture, Coleoptera infesting, 355, 489; attacked by termites, 296, 485; treatments of, 486.

furvus, Pediobius.

graminearum, *Fusarium* relation Ostrinia nubilalis to, causing stalk rot of maize in U.S.A., 398.

Fusarium oxysporum, associated with Sitona hispidulus on alsike clover in Oregon, 275.

Fusarium Wilt (of banana), inhibition of, by secretion from Scaptocoris divergens, 565.

fusca, Busseola; Frankliniella.

fuscicollis, Ageniaspis; Agriotes.

fuscilabris, Ceratomegilla.

fuscipennis, Dahlbominus (see D. fuligino-

fuscofemoralis, Psammotermes.

fuscula, Macropsis. fuscus, Caryedon; Ptilinus.

6

gaebleri, Tarsonemoides.

galactina, Aptesis (Microcryptus).

galactinus, Xylocoris.

Galacturonic Acid, in squash variety resistant to *Diaphania nitidalis*, 471.

galeopsidis, Capitophorus (Cryptomyzus).
Galeopsis tetrahit, migration of Capitophorus galeopsidis to, in U.S.S.R., 596.

Galerucella luteola (on elm), bionomics of, in Portugal, 298; in Spain, 659;

DDT against, 659.

Galerucella tenella, characteristics and world distribution of, 507.

Gall Mite, Black-currant (see *Cecido-phyopsis ribis*).

gallaesolidaginis, Gnorimoschema.

gallarumulmi, Anthocoris.

Galleria mellonella, injurious in bee-hives in Turkey, 551; on pine, 130; factors affecting physical properties of haemolymph of, 157; parasites of, 130, 208, 381, 382; development of Dirhicnus alboannulatus in, 379; experiments with parasites of other insects and, 87; Streptococcus faecalis infecting larvae of, 620; infection of, with polyhedrosis virus, 427, 496; larvae of, in baits for Gryllotalpa, 442; coddled larvae of, in diet for Ephialtes buolianae, 391; use of, in experiments with nematode, 162, 163; phosphine fumigation against, 551.

Gambia, insects infesting harvested groundnuts in, 28, 178, 179; question of *Cicadulina* spp. transmitting groundnut virus in, 298; first record of *Contarinia sorghicola* in, 684.

Game-birds, effects of agricultural chemicals on, 433.

gamma, Autographa.

Gamma, Autographa Gammexane, 422.

Gammexane 22, containing BHC (q.v.), 120.

Garlic, Hepialus lupulinus on, in France, 7; Dyspessa ulula on, in Italy, 661; treatment of bulbs of, 661.

Gartolit (see BHC).

Gas Analyser, Infra-red, measurement of carbon-dioxide output of insects by, 40.

Gas Chromatography, of insecticide residues, 91, 92, 577; thermal isomerisation of endrin during analysis by, 561.

Gas-oil, as adhesive for seed treatments, 670; toxicity of, to beet, 670.

Gastrimargus nigericus, bionomics of, in Mali, 620.

Gaza Strip, Citrus pests in, 97.

GC-1283 (see Dodecachlorooctahydro-1,3,4-metheno-2H-cyclobuta[cd]pentalene).

Gebutox-flüssig (see Dinoseb). Geigy 24027 (see Pyrazothion).

Geigy 30493 (see O,O-Dimethyl S-3,4-Dichlorophenylthiomethyl Phosphorodithioate).

Geigy 30494 (see O,O-Dimethyl S-2,5-Dichlorophenylthiomethyl Phosphorodithioate).

Geigy 36368 (mixture of Geigy 30494 and Geigy 30493), in sprays against *Alabama argillacea*, **413**.

Gelatin, in baits for Gryllotalpa, 442; use

of, in tests of repellents, 305.

Gelis, hyperparasite of Thymelicus lineola in Ontario, 162.

Gelis acarorum, parasitising Hoplocampa spp. in Yugoslavia, 138.

Gelis tenellus, parasite and hyperparasite of Rhyacionia buoliana in Argentina, 144, 230.

geminata, Solenopsis.

genalis, Glypta.

General Chemicals 3582 (see Diethyl 1-(2,5-Dichlorophenyl)-2,2-dichlorovinyl Phosphate).

General Chemicals 3583 (see Diethyl 1-(2,5-Dichlorophenyl)-2-chlorovinyl Phosphate).

General Chemicals 3661 (see Diethyl 1, 3-Di(methoxycarbonyl)-1-propen-2-yl Phosphate).

General Chemicals 3707 (see Dimethyl 1,3-Di(methoxycarbonyl)-1-propen-2-yl Phosphate).

General Chemicals 4072 (see Diethyl 1-(2,4-Dichlorophenyl)-2-chlorovinyl Phosphate).

Geniocerus (see Tetrastichus).

Genite, against Cydia pomonella, 21; in sprays against Tetranychids, 419, 515; toxicity of DDT decreased by, 21.

Geocoris, predacious on cotton pests in Arizona, 639.

Georgia, Sphenophorus callosus on Cyperus in, 152; forest pests in, 19; 337, 418; pests of maize in, 152, 277; pests of peach in, 226, 336, 641, 642; virus disease of peach in, 336.

Geotrupes stercorosus (sylvaticus), experiment on cuticular penetration of, by

BHC, 43.

German Cockroach (see *Blattella germanica*).

germanica, Blattella; Vespula. germanus, Xyleborus (Xylosandrus).

Germany, pests of beet in, 253, 377, 439, 440, 441, 442, 545, 546; virus diseases of beet in, 253, 545, 546; pests of crucifers in, 254, 303, 440; Oscinella frit on cereals and grasses in, 45, 255; Aporia crataegi on fruit and other trees in, 380, 683; apple pests in, 9, 10, 107, 257, 376, 381, 439, 440, 441, 681; Rhagoletis cerasi on cherry in, 679, 680; Myzus persicae on peach in, 547; pests of vines in, 108, 109, 255; Eriophyes vitis, aphids, nematodes and dodder transmitting virus of vines in, 71; aphids and virus diseases of strawberry in, 546: dodder transmission of clover virus to Fragaria vesca in, 441; aphids and virus diseases of potato in, 255, 546, 547, 680; Leptinotarsa decemlineata on potato in, 252, 377, 440, 543, 684; Sciara amoena on carnation in, 254; Mamestra brassicae on poppy in, 11; aphids and virus disease of tobacco in, 303; forest pests in, 10, 11, 44, 105, 106, 107, 140, 141, 168, 185, 186, 248, 250, 251, 252, 253, 375, 376, 441, 494, 682, 683; Myzus ascalonicus infesting stored onions in, 492; insects intercepted in importations of grain and oil-seed cake into, 10, 11; Hylotrupes baiulus infesting timber in, 253, 254; resistance of timbers from, to termites, 649; pests of textiles in, 377; distribution of primary food-plants of Aphis nasturtii in, 255; other insect pests in, 44, 139, 140, 430, 442, 649, 688; Tetranychus telarius in, 688; book on pests and diseases of vegetables, pulses, fruits, herbs and mushrooms in, 425; Carabid populations in fields and hedges in, 377; Formica spp. in, 104, 105; woodpeckers destroying F. polyctena in, 104; insects taken by bats in, 684; natural enemies of pests in, 10, 45, 106, 140, 141, 168, 251, 252, 257, 319, 375, 376, 377, 380, 381, 386, 439, 442, 494, 681, 682, 683, 684; Prospaltella perniciosi established in, Quadraspidiotus perniciosus, against **681**; Protozoa infecting *Tortrix viridana* in, 105; virus disease of A. crataegi in, 683.

germari, Aelia.

germinationis, Opomyza.

Gesarol (see DDT).

Ghana, Mirids on cacao in, 651; first record of *Contarinia sorghicola* in, 685; susceptibility or resistance of hardwoods to termites in, 649.

ghorfii, Asolcus.

gibba, sens. strict., Eurytoma (Bruchophagus) (see E. platyptera).

Gibberellin, effects on populations of Tetranychids of treatment of plants with, 410.

gifuensis, Macrocentrus.

gigas, Hyponomeuta.

Gilpinia abieticola, on spruce in Germany, 107; population fluctuations of, 107.

Gilpinia hercyniae (on spruce), in Canada, 24, 25, 163, 209, 341; in Germany, 107; population fluctuations of, 107; parasites of, 24, 25, 209; pathogenicity of Bacillus cereus to, 283; effects and transmission of virus diseases of, 25, 341, 342.

Gilpinia pallida, on spruce in Germany, 251; on pine in Sweden, 426; effects of soil fertilisers on populations of, 251. Gilpinia polytoma, on spruce in Germany,

107; population fluctuations of, 107. Gilpinia verticalis, on pine in Sweden, 426. Ginkgo biloba, factors affecting attraction of Popillia japonica to, in Virginia, 577.

Gipsy Moth (see Lymantria dispar).

giraulti, Anagrus.

glabra, Thaumatomyia.

glabrata, Ametastegia.

glabratus, Hylastes (Hylurgops).

Gladiolus, as reservoir of plant viruses, 271.

glasgowi, Haemolaelaps.

Glass Fibre, lining of fossae with, for

stored grain, 178.

Gleditsia, damaged by Homadaula albizziae in Kentucky, 273; Eulecanium corni on, in U.S.S.R., 13; systemic activity of insecticide in, 273.

Glenea lefebvrii, measures against, on cacao in Dutch New Guinea, 76.

Glischrochilus quadrisignatus quadrisignatus, bionomics of, in relation to maize and populations of Ostrinia nubilalis in Iowa, 329.

Globe Artichoke (see Cynara scolymus). glomeratus, Apanteles.

Glossary, of terms in entomology, 301.

gloverii, Lepidosaphes.

Glucose, in diets for locusts, 5.

Glue, protection of timber with, against termites, 486.

Glutamic Acid, in saliva of *Aphis pomi*, **95**; relation of, to oviposition by *Tetranychus telarius*, **547**.

Glutamin, in honeydew of Aphis pomi, 95; relation of, to oviposition by

Tetranychus telarius, 547.

Glycaspis, on Eucalyptus in New South Wales, 232, 233; new species of, 232, 233.

Glycaspis (Alloglycaspis) baileyi, sp.n., bionomics of, on Eucalyptus in New South Wales, 232, 233; birds attracted to excretions of, 233; sooty mould associated with, 233; natural enemies of, 233.

Glycine, in relation to feeding response

of wireworms, 528.

glycinivorella, Leguminivora (Grapholitha). Glycogen, metabolism of, during flight

by aphids, 203.

Glycyphagus destructor (in stored cereals and seeds), in Canada, 155, 424; in

Poland, 364.

Glyodin, in spray mixtures against orchard pests, 20, 21, 159, 274; effects of sprays containing, on beneficial arthropods, 36; toxicity of insecticides decreased by, 274.

Glypta, parasitising Choristoneura spp. in Europe and N. America, 169; parasitising *Pelatea festivana* in Italy,

535.

Glypta cicatricosa, parasitising Tortrix

viridana in U.S.S.R., 598.

Glypta fumiferanae, parasitising Choristoneura spp. in U.S.A., 24, 212, 337; vertical distribution of, 24; effect of tree location on degree of parasitism by, 212.

Glypta genalis, parasitic on Aphelia

viburniana in Germany, 376.

deesae, parasitising *Glyptomorpha* Raphimetopus ablutella in India, 119. gnidiella, Cryptoblabes.

Gnophothrips piniphilus, on jack pine in

Ontario, 529.

Gnorimoschema, type species and status of, 235.

gallaesolidaginis, Gnorimoschema 235. Gnorimoschema heliopa (see Scrobipalpa). Gnorimoschema instabilella (see Scrobipalpa).

Gnorimoschema ocellatella (see Scrobi-

palpa).

Gnorimoschema operculella (see Phthorimaea).

Gogu (see Hibiscus cannabinus).

gonagra, Caryedon.

gonostigma, Orgyia.

Gooseberry, Cape (see Physalis peruviana). Goryphus nursei, parasitising Earias spp. in India, 37.

gossypiella, Pectinophora (Platyedra).

gossypii, Aphis; Sphenoptera.

Gossypium, Pectinophora gossypiella on species of, in Mexico and U.S.A., 511. (See also Cotton.)

Gossypium hirsutum, resistance in races and varieties of, to infestation by Pectinophora gossypiella, 269.

Gourd, Aulacophora foveicollis on, in India, 173.

gracile, Eubadizon; Thelairosoma.

gracilis, Telenomus.

graminea, Diabrotica.

graminicola, Nysius.

graminis, Antonina; Typhlodromus.

graminum, Siteroptes; Toxoptera (Schizaphis).

granarium, Macrosiphum (see M. avenae); Trogoderma.

granarius, Aphodius; Sitophilus (Calan-

Granary Weevil (see Sitophilus granarius). Grand Fir (see Abies grandis).

grandiosella, Zeadiatraea.

grandis, Anthonomus; Asolcus; Colaspinella.

Granosan M (see Ethylmercury-ptoluenesulphonanilide).

granulatus, Carabus.

Grapefruit, Myllocerus dorsatus on, in India, 173; Coccids on, in Israel, 611, 612, 614; Anastrepha ludens on, in Mexico, 280; distribution of oilemulsion deposits on, 221.

Grapholitha funebrana (see Cydia).

Grapholitha glycinivorella (see Leguminivora).

Grapholitha molesta (see Cydia).

Graphosoma semipunctatum, method of rearing, 654.

Grass, Bermuda (see Cynodon dactylon). Grass, Buffel (see Cenchrus ciliaris).

Grass, Guinea (see Panicum maximum). Grass, Pangola (see Digitaria decumbens).

Grass, Quack (see Agropyrum). Grass, Rhodes (see Chloris gayana).

Grass, Rye (see Lolium spp.).

Grasses, Spodoptera exempta on, in Africa, 530; Pseudaletia unipuncta on, in N. America, 393, 394; pests of, in Australia, 34, 165, 166, 424, 504, 589, 684; pests of, in Britain, 321, 430, 487, 488; Camarota curvinervis on, in Bulgaria, 110; pests of, in Canada, 71, 73, 156, 158, 161, 162, 389, 523, 524, 584-586; Hydrellia griseola, on, in Denmark, 354; pests of, in Finland, 41, 102; Calligypona pellucida on, in France, 373; oviposition and feeding of Oscinella frit on, in Germany, 45; Aphrodes bicinctus on, in Hungary, 307; pests of, in India, 79, 119, 406, 684; Calocoris spp. on, in Italy, 310; Meromyza saltatrix on, in Japan, 96; S. exempta on, in Kenya, 179, 530, **686:** Chloriona cubana on, in Mexico. 518; pests of, in New Zealand, 231, 505, 506, 684; Lamellicorn larvae infesting, in S. Rhodesia, 38; grasshopper on, in Turkey, 351; pests of, in U.S.S.R., 191, 244, 344, 345, 363, 447, 542, 596; pests of, in U.S.A., 95, 152, 160, 219, 223, 224, 265, 396, 406, 419, 464, 519, 520, 521; Blissus leucopterus on, in Venezuela, 98; Anacentrinus deplanatus migrating to sorghum from, 219; ants preventing establishment of, 589; resistance of, to Chilo suppressalis, 78; preferences of Pentatomids for, 653; overwintering of Sphenophorus callosus on, 152; survey of data on virus diseases of, 684; vectors of virus diseases of, 71, 518, 523, 524, 684, 690; relation of mites and thrips to silver top of, 156; effect of infestation on yields of, 223, 420, 505; cultural practices in relation to infestation of, 488; effects of dieldrin spray against O. frit on 430; seed treatment of, 589; insecticide residues on, 64, 92, 280, 330; suction apparatus for sampling arthropods on, 291; use of X-rays for detecting stem borers in. 495; use of, in rearing wireworms, 575; simulation of cellulose content of, in diets for locusts, 5.

Grasshoppers, in Argentina, 356; in Canada, 575, 631; in U.S.A., 223, 224, 406, 521, 575; effects of, on grass yields, 223; transmitting plant viruses, 443; bionomics of, 521; effects of rainfall on populations of, 224; natural enemies of, 356, 500, 576; infected by Nosema locustae, 582; timing of survey of, 575; question of accuracy of forecast maps for, 631; insecticides against, 521; question of inter-specific fertility of, 694. (See also Calliptamus, Camnula, Chortophaga, Hieroglyphus, Melanoplus and Thisoicetrinus.)

gratiosus, Rhipidothrips.

Greece, pests of apple in, 239; pests and diseases of cotton in, 352; *Dacus oleae* on olive in, 237, 238.

greeni, Bracon (Microbracon).

gregaria, Schistocerca.

Grenadilla (see Passiflora quadrangularis). Grewia asiatica, Perissopneumon tectonae on, in India, 174.

grisator, Sthenias.

griseana, Enarmonia.

Griselda radicana, bionomics and natural enemies of, on spruce in Quebec, 157. griseola, Hydrellia; Leucopis.

grisescens, Lydella stabulans.

griseus, Hyperodes.

Groundnut Butter, in baits for insects, 339, 464.

Groundnut Meal, in baits for ants, 464. Groundnut Oil, in baits for ants, 464.

Groundnut Sweepings, in baits for Ephestia elutella, 374.

Groundnuts, Thysanoptera on, in Alabama, 151; new pests of, in Brazil, 145, 646; pests of, in Congo, 609, 610; Tetranychus telarius on, in Israel, 613; pests of, in Senegal, 298; list of pests of, in South-East Asia and Pacific area, 423; vectors of virus diseases of, 298, 609-611; review of pests and diseases of, 693; toxicity of phorate to, 152; question of heptachlor residues in, 479.

Groundnuts (Stored), insects infesting, 28, 178, 179, 298, 558, 583; experiments on infestation of, by *Ahasverus advena*, 657; list of pests of, in South-East Asia and Pacific area, 423; treatments of, against infestation, 28, 178, 179, 558.

Gryllotalpa gryllotalpa, damaging cotton in Greece, 352; in Germany, 442; insects in diet of, 442; tests of baits for, 442; insecticides against, 352.

Gryllus, American species of, 567.

Gryllus (Acheta) assimilis, in America and West Indies, 567, 687; on forage crops in Argentina, 356; effects of β-radiation on oviposition by, 686, 687; other crickets misidentified as, 567.

Gryllus campestris, species allied to, in America, 567; on cotton in Greece, 352; insecticides against, 352.

Gryllus domesticus (see Acheta).

Gryon (Hadronotus) monspeliensis (parasitising eggs of Pentatomids), in Middle East, 485; in Morocco, 655. guatemalensis, Typhlodromus.

Guava, Perissopneumon tectonae on, in India, 174; Megachile frontalis on, in New Guinea, 234.

Guiana, British, pests of jute in, 27, 28; miscellaneous pests in, 551.

Guiana, Dutch, *Rupela albinella* on rice in, 93; *Trigona* spp. damaging crops in, 144.

guildi, Ips.

Guinea, new weevil on *Cola* in, 355. Guinea Grass (see *Panicum maximum*). Guineacorn (see Sorghum).

gularis, Aphomia.

gurdaspurensis, Odontotermes.

Gusathion (see Azinphos-methyl). Guthion (see Azinphos-methyl).

Gymnetron antirrhini, Linaria vulgaris controlled by, in Canada, 386.

Gynandrophthalma limbata, on pistachio in Turkey, 609.

Gynandrophthalma unipunctata, on pistachio in Turkey, 609.

Gypsonoma dealbana, on hazel in Turkey, 83; effects of sulphur dust on populations of, 83.

Gypsum, effect of, on toxicity of insecticide dusts, 118.

Gypsy Moth (see Lymantria dispar). Gyrolachnus, classification and new species of, 493.

H

H-5727 (see m-Isopropylphenyl Methyl-

carbamate).

Habrocytus, parasitising larvae of Caliroa limacina in Quebec, 26; competition between Dahlbominus fuliginosus and species of, parasitising Diprion pini in Yugoslavia, 31.

Habrolepis aspidioti, sp.n., parasitising Quadraspidiotus perniciosus, 491; in

Eritrea, 491.

Habrolepis obscurus, sp.n., parasitising Coccids in S. Africa, 491.

Habrolepis rouxi, oviposition of Physcus spp. on, 266.

hacupha, Oroscopa.

Hadena sordida, bionomics and ecology of, on wheat and grasses in U.S.S.R., 447; relation of diet to fertility of, 447. Hadena suasa, feeding habits of, 620.

Hadronotus, parasitising *Amblypelta* lutescens papuensis in Papua, 235.

Hadronotus carinatus, parasitising Eury-gaster integriceps in Turkey, 43; study on reproductive capacity of, 43.

Hadronotus monspeliensis (see Gryon). Hadrotettix trifasciatus, on rangelands in Arizona, 224.

haematopus, Penthaleus (see P. major). Haemolaelaps casalis, in stored grain in Canada, 154.

Haemolaelaps glasgowi, in stored grain in Canada, 154.

haemorrhoidalis, Buprestis; Heliothrips.

Hail, damage to apple by, 159.

Hairy Vetch (see Vicia villosa). Halictus, parasitised by Heterostylum robustum in U.S.A., 565.

Halotydeus destructor, characteristics and world distribution of, 507.

Haltica oleracea, recorded on strawberry in Yugoslavia, 242; food-plant range of, 242.

Haltica quercetorum, on oak in Yugoslavia, 181.

hamatus, Diplostichus.

hammondi, Psorosina.

hampei, Stephanoderes (Hypothenemus). Hansenius, predacious on Oecophylla sp. in Zanzibar, 29.

Haplodiplosis equestris, on cereals and Agropyrum repens in Sweden, 101; insecticides against, 101. Haplomyza caulophaga, on sea-kale beet in Queensland, 295.

Haplothrips, associated with abnormal growth of coffee in India, 448.

Haplothrips aculeatus, on wheat in China, 625.

Haplothrips tritici (on cereals), bionomics of, in Bulgaria, 348; in U.S.S.R., 343; migration habits of, 343; natural enemies and fungus disease of, 348; measures against, 348.

hariolus, Phaeogenes.

Harpalus rufipes (pubescens) (on strawberry), in Britain, 670, 671; in Bulgaria, 59; bionomics of, 670, 671; sampling methods for, 671; malathion sprays against, 671.

harti, Hyaliodes.

harveyi, Ptychomyia (Bessa) (see P. selecta).

havilandi, Cryptotermes.

Hawaii, pests of litchi in, 463; Dacus dorsalis in, 563; diseases of Pseudaletia unipuncta in, 283, 428; virus disease of Spodoptera mauritia in. 196.

Hawthorn (see Crataegus).

Hay, persistence of insecticide residues on, 330.

Hazel (see Corylus avellana).

Heat, effects of, on survival and fecundity of Callosobruchus chinensis, 449; miggrating larvae of Trichiocampus viminalis killed by, 392; study on effects of insects on, in stored grain, 548; treatment of *Narcissus* bulbs with, against Merodon equestris, 288. (See also Temperature and Water, Hot.)

Hedera helix, Tetranychus multisetis not feeding on, 67.

hederae, Acaricalus; Aspidiotus.

heimi, Trinervitermes. heinrichi, Homoeosoma.

(Sunflower), Helianthus annuus Pyralid on, in Argentina and Uruguay, 230; Dasiops alveofrons reared from, in Arizona, 128.

helichrysi, Anuraphis (Aphis).

Heliconia, Lepidoptera damaging, Costa Rica and Honduras, Nacoleia octasema on, in New Guinea, 354; as natural host of Pseudomonas solanacearum, 520.

Helicopter (see Aircraft).

heliopa, Scrobipalpa (Gnorimoschema).

heliopae, Chelonus.

Heliothis, key to species of, in Queensland, 588; in U.S.A., 406; feeding preferences of species of, on Nicotiana spp.,

Heliothis armigera, on cotton in Israel, 351; on maize in Kenya, 179; on flax and other plants in Queensland, 588; in Sweden, 503; on solanaceous plants, 503, 588; insecticides against, 351, 503; characters of, 588.

Heliothis assulta, on Physalis and tomato in Queensland, 589; characters of, 588. Heliothis dipsacea, in Bulgaria, 59, 537, 538, 539; on mint, 59; on lucerne, 537, 538, 539; other food-plants of, 538; bionomics of, 538; diseases and natural enemies of, 538; insecticides against, 538, 539.

Heliothis peltigera, on mint in Bulgaria,

Heliothis phloxiphaga, on Mentha spp. in Washington, 324; DDT against, 324. Heliothis punctigera, in Australia, 233, 234, 588; in Sweden, 503; on flax, 588; on lucerne, 233, 234; on solanaceous and other plants, 503, 588; DDT sprays against, 234, 503; characters of, 588.

Heliothis rubrescens, in Queensland, 588; characters of, 588.

Heliothis virescens, on tobacco in S. Carolina, 465; baits for, 465; measures against, 465.

Heliothis zea (in U.S.A.), on cotton, 323, 408, 409, 412, 413, 477, 508, 515, 638, 639; on crucifers, 404; on lettuce, 514; on maize, 227, 275, 277, 327, 328, 508, 509, 580; varietal susceptibility of maize to, 327, 328, 476; Nicotiana spp. resistant to, 406; factors affecting adult life-span of, 563; fluctuations in populations of, 328; natural enemies of, 66, 580, 638, 639; light-traps for, 477; lighttrap catches of, 219, 328; virus and bacterial diseases against, 515, 580; insecticides against, 227, 275, 277, 406, 409, 412, 413, 508, 509, 514, 515, 552, 580; larval age of, in relation to penetration by DDT, 472; resistance to DDT in strain of, 406; effects of insecticides against Anthonomus grandis on populations of, 323; method of rearing, 406: artificial diet for, 517.

Heliothrips haemorrhoidalis, on avocado in Morocco, 383; review of plants infested by, in Puerto Rico, 257; map of distribution of, 1; systemic insecticides against, 257.

Helopeltis antonii, on cacao in Dutch New Guinea, 76; BHC dusts against, 76.

Helopeltis clavifer, on cacao in Dutch New Guinea, 76.

helvolus, Aphycus (Metaphycus).

Hemiberlesia lataniae, large-scale rearing of natural enemies of Coccids on, 229. Hemiberlesia rapax, on Leptospermum in New Zealand, 588.

Hemichroa crocea, pathogenicity of Bacillus cereus to, 283.

hemipterus, Carpophilus; Valgus.

hemisphaerica, Saissetia (see S. coffeae). Hemitarsonemus latus, in Brazil, 120, 643, 645; in British Guiana, 28; on cotton, 643, 645; on jute, 28; as precursor of attack on jute by Tetranychus telarius, 28; on papaya, 120; acaricides against, 28, 120, 643, 645.

Hemiteles, parasitising Operophtera

brumata in U.S.S.R., 190.

Hemiteles castaneus, parasitising Pachynematus scutellatus in Czechoslovakia, 246.

Hemiteles pastranai, hyperparasite of Rhyacionia buoliana in Argentina, 230. Hemlock, Western (see Tsuga heterophylla).

Hemp (see Cannabis sativa).

Hendecaneura shawiana, bionomics of, on

blueberry in New York, 567.

Hepialus lupulinus, in Britain, 430, 431; in France, 7; on anemone, 430; on cereals, 7, 431; on strawberry and other soft fruits, 7, 430, 431; on other plants, 7, 431; bionomics and parasites of, 7; tests and uses of insecticides against, 7, 431

Heptachlor, against ants, 420, 421, 483; against aphids, 420, 421, 498; against other Hemiptera, 72, 627; against Curculionids, 16, 152, 219, 275, 280, 329, 331, 417, 509, 609, 636; against Elaterids, 371, 372, 420, 421; against Melolonthids, 493, 570; against other Coleoptera, 323, 396, 417, 420, 421, 471, 641, 674; against Hylemya, 50, 51, 226, 388, 433, 568, 674; resistance to, in strains of *Hylemya* spp., 388; against Trypetids, 325, 326, against other Diptera, 101, 226, 330, 402; against Lepidoptera, 16, 211, 277, 309, 420, 421, 478, 514, 534; cross-resistance to, in strains of Estigmene acraea, 480; against grasshoppers, 521; against termites, 577; against Thysanoptera, 71; toxicity Schizotetranychus andropogoni, 628; against pests of lucerne, 92; in dusts, 309, 323, 417, 534, 568, 627; in granules, 21, 145, 152, 219, 323, 325, 330, 331, 396, 402, 417, 479, 483, 568, 570, 636; granule size in relation to efficiency of. 396; in sprays, 16, 71, 72, 478, 509, 514, 568, 572, 636, 641; seed treatment with, 50, 51, 65, 66, 101, 226, 329, 433, 568, 609, 674; seed-furrow application of, 330; soil treatment with, 50, 51, 71, 145, 152, 219, 226, 275, 280, 309, 325, 326, 371, 372, 388, 396, 402, 417,

479, 493, 534, 568, 570, 577, 627; persistence and distribution of, in soils, 145, 146, 416, 417, 483, 570, 572, 573; treatment of selfing bags with, 277; treatment of planting water with, 388; pre-flooding treatment of rice-fields with, 329; humidity in relation to fumigant action of, 471; factors affecting persistence of, 471; experiments on systemic activity of, in peas, 146; effects of, on plants, 402, 675; residues of, in plants, 92, 331, **402**, **417**, **479**; residues of, in milk and tissues of cattle, 521; restriction on use of, in relation to hazards to wild life, 433; spectrophotometric determination of, 145; bioassay of, 146; metabolism of, 63, 64, 145, 146; heptachlor epoxide as derivative of, 64, 92, 145, 146, 417; effects of additives on stability of, 145; antagonistic action of sesamex with, 63, 64.

Heptachlor Epoxide, in tests against Tribolium castaneum, 471; as derivative of heptachlor: in insects, 64; in plants, 92, 145, 402, 417, 479; in milk, 92; volatilisation of, from soils, factors affecting persistence of, 471; humidity in relation to fumigant action of, 471; bioassay of, 146; oxidation of, increased by sesamex, 63, 64.

heraclei, Euleia (Acidia).

Graphosoma Heracleum sphondylium, semipunctatum reared on seeds of, 654. Herbs, pests and diseases of, in Germany,

Hercothrips (see Caliothrips).

Hercules 528 (see 2,3-p-Dioxane S,S-Bis (O,O-diethyl phosphorodithioate)).

Hercules 2032 (see O,O-Diethyl S-p-Oxathian-3-yl Phosphorodithioate). Hercules 3004 (see Diethyl 1-Ethylthio-1-

propen-2-yl Phosphate).

Hercules 3273 (see O,O-Dimethyl S-p-2-Oxathien-x-yl Phosphorodithioate).

Hercules 3895 (see Diethyl 2,2-Di(ethylthio)vinyl Phosphate).

Hercules 5727 (see m-Isopropylphenyl Methylcarbamate).

hercyniae, Gilpinia (Diprion).

Heringia dodecella (see Exoteleia).

Hermetia illucens, bionomics of, in New Zealand, 424.

heros, Rhynchites.

hertingi, Myxexoristops.

hesperidum, Coccus.

hesperus, Lygus.

Hessian Fly (see Mayetiola destructor). Heterostylum robustum, parasitising Halticus and wild bees in U.S.A., 564, 565; bionomics and natural enemies of, 565. Heterotermes convexinotatus, infesting timber in British Honduras, 76.

Heterotermes indicola, resistance of timbers to infestation by, 649.

HETP, toxicity of, to Locusta migratoria, 172; containing TEPP, 172.

Hevea brasiliensis (Rubber), systematic list of pests of, in South-East Asia and Pacific area, 423; attacked by termite in Malaya, 76, 77; Amblypelta lutescens papuensis on, in Papua, 235.

5, 6, 7, 8, 9, 9-Hexachloro-1, 4, 4a, 5, 8, 8ahexahydro-1,4,5,8-dimethanophthalazine N-Oxide, in sprays against Otio-

rhynchus cribricollis, 509.

5,6,7,8,9,9-Hexachloro-1,2,3,4,4a,5,8,8aoctahydro -1, 4, 5, 8-dimethanophthalazine Sulphate, in sprays against Otiorhynchus cribricollis, 509.

Hexaethyl Ethylthiomethylidynetriphosphonate, ineffective in sprays against

Lepidoptera, 409.

Hexagenia bilineata, nuisance value of, in Iowa, 262.

Hexamermis, infesting Chortophaga viridifasciata in Missouri, 576; infesting Operophtera brumata in U.S.S.R., 190.

Hexamermis arvalis, sp.n., parasitising Hypera variabilis in New York, 581.

Hexamermis brevis, 190.

Hexobarbital (see 5-(1-Cyclohexen-1-yl)-1,5-dimethylbarbituric Acid).

heydeni, Thorictodes.

hibisci, Typhlodromus.

Hibiscus cannabinus, pests of, in India, 36, 314, 594; varietal susceptibility of, to Agrilus acutus, 594.

Hibiscus esculentus, pests of, in Brazil, 357; pests of, in India, 36, 314, 317, 594; Pectinophora gossypiella on, in U.S.A., 511; Podagrica malvae semi-rufa on, in Yugoslavia, 181; growth stimulation of, by endrin, 594; seed treatment of, 357.

Hibiscus ficulneus, Earias spp. on, in India, 37.

Hibiscus rosa-sinensis, Earias spp. on, in India, 37.

Hibiscus sabdariffa, pests of, in India, 37, 594; varietal susceptibility of, to Agrilus acutus, 594.

Hickory (see Carya).

Hides, Dermestes infesting, in S. Africa, 49; treatment of, with insecticides, 49. Hieroglyphus nigrorepletus (in India), on cereals, 498; mite predacious on, 500; toxicity of insecticides to, 498.

hilaris, Aulacophora; Scymnus.

hippocastani, Melolontha.

Hippodamia convergens (predacious on aphids in U.S.A.), 401; effects of insecticides against aphids on populations of, 279, 281.

hirsutus, Maconellicoccus (Phenacoccus).

hirta, Epicometis.

Hispaniella lauri, parasitising Quadraspidiotus perniciosus in Switzerland, 664. hispidulus, Sitona.

parasitising Histochaeta marmorata, Apamea anceps in U.S.S.R., 363.

Hockeria, parasitising Laelia exclama-

tionis in India, 79.

Hofmannophila pseudospretella, in Britain, 289, 290; damaging textiles in Germany, 377; feeding habits of, 290; treatments of textiles against, 290;

method of rearing, 290.

Holland, carrot pests in, 100, 299, 608, 673; pests of crucifers in, 39, 608, 673; insects on other vegetables in, 193, 673; pests of fruit trees in, 444, 673, 682; Cecidophyopsis ribis transmitting reversion virus of black currant in, 302; aphids and virus diseases of strawberry in, 192; forest pests in, 42, 185, 205; Merodon equestris on Narcissus in, 42; Monoctenus juniperi on juniper in, 302, 303; Anuraphis plantaginea on plantain in, 673; Tetranychus telarius on roses, vines and other plants in, 193, 682, 688; Plutella maculipennis in, 430; natural enemies of pests in, 39, 205, 302, 673, 682.

Holly (see *Ilex*).

Hollyhock (see Althaea rosea).

hololeucus, Niptus.

Holotrichia (see Lachnosterna).

holoxanthus, Aphytis.

Homadaula albizziae, insecticides against, on ornamental trees in Kentucky, 273. Homalinotus coriaceus, BHC spray against, on coconut in Brazil, 559.

Homalodisca coagulata, in Georgia, 336, 337; as vector of virus disease of peach, 336; on oak, 336, 337; overwintering habits of, 336, 337; traps for, 336.

Homalotylus quaylei, parasitising Scymnus predators of Planococcus citri in Israel,

614.

Homoeosoma heinrichi, sp.n., on sunflower in Argentina and Uruguay, 230. Homona coffearia, on tea in Ceylon, 553, 554; Macrocentrus homonae introduced into Ceylon against, 553; sprays against, 554.

homonae, Macrocentrus. hominivorax, Callitroga.

Honduras, pests and leaf-spot disease of banana in, 462, 463, 565, 566; insects infesting Heliconia in, 520.

Honduras, British, termites attacking trees and timber in, 76.

Honey, effects of, on fecundity of Aptesis basizona, 155; sources of, for Formica polyctena, 105; in diets for insects, 29, 105, 208, 290, 314, 379, 391, 406, 422, 563, 624; in rearing media for mites, 641; use of, for labelling Anthonomus grandis with 32P, 472; use of, in study of feeding habits of Pieris brassicae, 51, 52.

Honeydew, rate of excretion of, by aphids, 322; in diets for insects, 208, 209, 391, 422, 612; contamination of apples by, 402: in relation to sooty mould, 415; growth reactions of plants in relation to auxins in, 629; amino acids in, 95; paper chromatography of auxins in, 629.

Hong Kong, new parasite of Chrysomphalus ficus introduced into Israel from, 22.

Hoplocampa brevis, characteristics and

world distribution of, 507.

Hoplocampa flava (on plum), in Poland, 61, 544; bionomics of, in Yugoslavia, 138; natural enemies of, 138; insecticides against, 139, 544.

Hoplocampa minuta (on plum), in Poland, 61, 544; in Rumania, 56; in Yugoslavia, 138; bionomics of, 56, 61, 138; natural enemies of, 138; insecticides

against, 56, 139, 544.

Hoplocampa testudinea (on apple), in Finland, 103; in France, 665, 666; bionomics of, 666; attracted to ultraviolet light, 666; sprays against, 666.

Horcias nobilellus, characteristics and

world distribution of, 507.

hordei, Chlorops (see Meromyza saltatrix). Hormiopus ptericoptophagus, gen. et sp.n., parasitising Ptericoptus sp. in Argentina, 552.

Hornbeam (see Carpinus betulus).

Hornworm, Tobacco (see Protoparce sexta).

Horogenes (see Casinaria and Nythobia). hortensis, Chaetocnema.

horticola, Anomala (Phyllopertha).

hottentota, Villa.

hottentotta, Eurygaster.

House-fly (see Musca domestica).

Houses, Hofmannophila pseudospretella infesting, in Britain, 290: Drosophila virilis in, in Japan, 622; Bryobia praetiosa in, 419, 519.

howardi, Diabrotica undecimpunctata.

howardii, Aphelinus.

humanus, Pediculus.

humeralis, Rhynchocoris; Sitona.

Humidity, effects of: on Coleoptera, 127, 165, 166, 318, 323, 486; on Diptera, 97, 139, 446; on Hemiptera, 566, 601; on Lepidoptera, 4, 5, 48, 56, 291, 368, 427, 665; on locusts, 170, 285, 286; on mites, 67, 91, 549, 583; on beneficial insects, 257, 258, 317, 601; on toxicity of insecticides, 356, 676; on persistence and fumigant action of insecticides, 471; on systemic activity of dimethoate, 472; on insecticides in relation to soils, 272, 416; on toxicity of insecticides to seeds, 218, 331, 332, 579.

humilis, Hyalopteroides; Hyperodes; Irido-

_myrmex.

Hungary, pests of cereals in, 306, 307; Tetranychids on cotton and fruit trees in, 307; Aphrodes bicinctus on clover, lucerne and grasses in, 307; Eurytoma spp. infesting seeds of leguminous plants in, 305; potato pests in, 305, 307; pests of rape in, 308; Ostrinia nubilalis on hemp in, 306; Cryptorhynchus lapathi on deciduous trees in, 307.

Hyadaphis brassicae (see Brevicoryne). Hyalesthes obsoletus, on cereals in Hungary, 307; on potato and other plants in Yugoslavia, 240; transmitting stolbur virus, 240, 307; dusts against, 241.

Hyalinaspis, species of, on Eucalyptus in

Australia, 552.

hyalinata, Trigona.

hyalinipennis, Oxycarenus.

hyalinus, Japananus.

Hyaliodes harti, predacious on Bryobia rubrioculus in Nova Scotia, 523.

Hyalochloria denticornis, eggs of Anomis texana destroyed by, in Peru, 560.

Hyalopteroides humilis, on Dactylis glomerata in New Brunswick, 586.

Hyalopterus amygdali (on peach), in Portugal, 617; in Yugoslavia, 240; bionomics and natural enemies of, 240; sprays against, 240, 617.

Hyalopterus pruni, in Poland, 188; in Yugoslavia, 240; in Wisconsin, 160; on grass, 160; on plum, 188, 240; bionomics and natural enemies of, 240; sprays against, 188.

hydei, Drosophila.

Hydrangea, Tetranychus multisetis not feeding on, 67.

Hydrellia griseola, on cereals and grasses in Denmark, 354.

Hydrellia tritici, on grasses in New Zealand, 231.

Hydrochloric Acid, experiments with, in decomposition of froth masses of Cercopids, 321.

Hydrogen Cyanide (as fumigant), against Aonidiella aurantii, 383; against pests of stored products, 267, 347, 436, 584, 694; effects of, on respiration and activity of Coleoptera, 584; effects of, on germination of seed, 579; apparatus for treating insects with radioactive, 356.

Hydrogen Fluoride, treatment of timber with, against *Hylotrupes bajulus*, **647**,

648; fumigant action of, 648.

Hydrogen Phosphide (as fumigant), against pests of stored products by Delicia process, 347; toxicity of, to Coleoptera, 267; against Galleria mellonella in bee-hives, 551. (See Aluminium Phosphide.)

Hydrogen Sulphide, factors affecting evolution of, from lime-sulphur sprays,

672

Hydropsyche orris, nuisance value of, in Iowa, 262.

Hydroxyethyl Cellulose Polymer, tests and uses of, against *Tetranychus telarius*, 272; phytotoxicity of, 272; efficacy of mevinphos increased by, 272.

Hydroxypropyl Starch, toxicity of, to Tetranychus telarius, 272.

Hylastes glabratus, infesting spruce in Germany, 44; tests of attractants for, 44

Hylastes obscurus (on red clover), in Quebec, 453, 454; in U.S.A., 323, 417; bionomics and adult habits of, 453, 454; root-rot associated with, 323; insecticides against, 323, 417.

Hylastes rufipes, vector of Ceratocystis ulmi on elm, in Canada, 124.

Hylastinus (see Hylastes).

Hylemya antiqua (on onion), in Denmark, 608; in New York, 226; in Quebec, 455; fungus infesting, 455; insecticides against, 226, 608.

Hylemya brassicae (on crucifers), in Britain, 39, 286, 287; in Finland, 103, 104, 674; in Germany, 303; in Holland, 39, 40, 673; in Ontario, 162; in U.S.S.R., 362, 542; in U.S.A., 510, 516, 517, 568; bionomics and adult habits of, 39, 261, 516, 517; factors affecting diapause in, 39, 542; population fluctuations of, 40; life tables for, 286; effects of, on biochemistry of food-plant, 287; natural enemies of, 39, 40, 162, 362; insecticides against, 103, 104, 362, 510, 568, 673, 674; resistance to chlorinated hydrocarbons in strains of, 510; effects of insecticides on populations of predators and, 40.

Hylemya cilicrura, in Ontario, 388; in U.S.A., 420, 568; on maize, 420; on radish, 568; on tobacco, 388; map of distribution of, 357; bait for, 388; insecticides against, 388, 420, 568; resistance to insecticides in strains of,

Hylemya coarctata, on cereals in Britain, 50, 51, 433; characteristics and world distribution of, 507; seed and soil treatments against, 50, 51; restriction on use of seed treatments against, 433.

Hylemya floralis (on crucifers), in Finland, 103, 104, 674; in Germany, 254; seed and soil treatment against, 103, 104, 254.

Hylemya laricicola, bionomics of, damaging larch cones in U.S.S.R., 244.

Hylemya trichodactyla (liturata), tobacco in Ontario, 388; insecticides against, 388; resistance to insecticides in strains of, 388.

Hylesinus fraxini, on ash in Czechoslovakia, 187; parasites and hyperparasites

of, 187, 188.

Hylesinus vestitus, on pistachio in Turkey, 609.

Hylobius abietis (on pine), in Finland, 194; in Spain, 660; characteristics and world distribution of, 506; DDT treatment of seedlings against, 194.

Hylobius radicis, bionomics of, on pines in Ontario, 457, 458; temporary storage

of stages of, 281.

Hylotrupes bajulus, in timber in Germany, 253, 254; treatments of timber against, 253, 254, 646, 647, 648; surveys of data on, 646, 648.

Hylurgopinus (see Hylastes).

Hylurgops (see Hylastes).

Hymenia recurvalis, on Amarantus viridis in Fiji, 88; bionomics of, on Amarantus and other vegetables in India, 498; BHC dusts against, 498.

hyoscyami, auct., Pegomyia (see P. betae). Hyoscyamus niger, Leptinotarsa decemlineata on, in Bulgaria, 366.

Hypera, species of, on clover in U.S.S.R., 114.

Hypera meles, Microctonus aethiops reared on, 391.

Hypera nigrirostris, Microctonus aethiops

reared on, 391.

Hypera variabilis (postica) (on lucerne), in Bulgaria, 60; in France, 390, 391; in India, 119, 317; in U.S.A., 323, 331, 400, 576, 581, 636; bionomics of, 331; flight habits of, 576; factors affecting development of, 323; natural enemies of, 390, 391, 400, 581; adhesive traps for, 576: insecticides against, 317, 331,

Hyperchalcidia soudanensis, parasitic on Coniesta ignefusalis in Nigeria, 292.

Hyperodes bonariensis, food-plant preferences of, on pastures in New Zealand, 505, 506; characteristics and world distribution of, 507; insecticides against, 506; misidentified as H. griseus, 505.

Hyperodes griseus, H. bonariensis mis-

identified as, 505.

Hyperodes humilis, on maize in Florida, 275, 512; insecticides against, 275, 512; effect of DDT on natural enemies of,

Hyperomyzus staphyleae (see Rhopalosiphoninus).

Hyperteles, classification and new species of, 493.

Hyphantria cunea, in Austria, 312; danger of introduction of, into Italy, 311; in Yugoslavia, 241, 368, 682; on mulberry and other plants, 312; bionomics of, 312; factors affecting ecdysis in, 241; natural enemies of, 312, 682; activation of virus in Lymantria dispar by virus from, 368; measures against, 312; form of, 312.

Hyphantria cunea textor, in Europe, 312.

hyphantriae, Meteorus; Rogas. Hypocrita jacobaeae (see Tyria).

Hypolithus abbreviatus, on potato in Nova Scotia, 73; reared on wheat, 73.

Hypolithus bicolor, in Saskatchewan, 155; adult activity of, 155, 156; traps for, 155, 156.

Hypolixus nubilosus, on Citrus in Turkey, 82.

Hyponomeuta cognatellus, on Euonymus in Finland, 194; parasite of, 87.

Hyponomeuta euonymellus, bionomics of, on Prunus padus in Finland, 194, 195; not occurring on Euonymus, 194; parasites of, 195.

Hyponomeuta gigas, on poplar in Canary Islands, 659.

Hyponomeuta padellus, on fruit trees in Bulgaria, 110; form of, on Sorbus aucuparia in Finland, 194, 195; parasites of, 195; biological separation of forms of, 111; classification of group of, 194.

Hyponomeuta padellus malinellus (on apple), in Bulgaria, 110; in Finland, 194, 195; in Germany, 107; in U.S.S.R., 597; bionomics of, 111, 194, 597; parasites of, 111, 195; experiments with Thelohania weiseri and, 105; tests of Bacillus thuringiensis against, 107; sprays against, 111; H. p. padellus not interbreeding with, 111.

Hyponomeuta padellus padellus, sprays against, on plum in Bulgaria, 110; bionomics and parasite of, 111; H. p. malinellus not interbreeding with. 111. Hyponomeuta rorellus, on poplar in

Spain, 659.

Hyposoter exiguae, bionomics of, parasitising Lepidoptera in California, 66; parasite of, 67.

Hypothenemus hampei (see Stephanoderes). Hypsauchenia subfusca, on mulberry in

India, 626.

Hyssopus rhyacioniae, parasitising Dioryctria zimmermanni in Illinois, 153. Hysteroneura setariae (see Carolinaia).

hystrix, Abacarus.

T

IAA (see 3-Indoleacetic Acid). IAE (see Ethyl 3-Indoleacetate). IAN (see 3-Indoleacetonitrile). IBA (see 3-Indolebutyric Acid). Ibertox (see Dinitroisobutylphenol).

Icerya aegyptiaca, characteristics and world distribution of, 507.

Icerya purchasi, on pine seedlings and other plants in Puerto Rico, 257; controlled by Rodolia cardinalis, 257; sprays against, 257.

sarcitorius, parasitising Apamea anceps in U.S.S.R., 363.

ictericus, Exenterus.

idaei, Aphis.

Idaho, Ips oregonis on pine in, 386; mites in stored grain in, 149; mite attacking I. oregonis in, 386.

Idiocerus alkani, sp.n., on pistachio in

Turkey, 81.

Idiocerus stali, on pistachio in Turkey, 91, 609.

ignefusalis, Coniesta.

Ilex crenata, pests of, in New York, 19; activity of systemic insecticides in, 19. Ilex opaca, Acaricalus hederae on, in British Columbia, 529.

ilicis, Paratetranychus (Oligonychus).

Illinois, maize pests in, 420; Dioryctria zimmermanni on pines in, 153; Sitotroga cerealella and fungi infesting stored grain in, 269, 414; natural enemies of D. zimmermanni in, 153.

illucens, Hermetia. imaginis, Thrips. imitata, Melanolophia. immaculata, Scutigerella. impexus, Scymnus (Pullus). impressifrons, Clivina. Inareolata (Dioctes) (see Nythobia). inaron, Coccophagus. incarnatus, Bruchidius. incerta, Orthosia (Taeniocampa). incertulas, Tryporyza (Schoenobius). inclusum, Trogoderma. inconspicua, Magdalis.

Incurvaria rubiella, characteristics and world distribution of, 507.

Indalone, in sprays against Scolytus

multistriatus, 217.

India, Eutetranychus banksi on castor in, 627; pests of cereals in, 79, 314, 498, 627, 684; pests of grasses in, 119, 406, 684; virus diseases of cereals and grasses in, 684; sugar-cane pests in, 119, 171, 297, 497, 498, 553, 628; Citrus pests in, 80, 173, 491; tristeza disease of Citrus in, 80; pests of mango in, 170, 315; pests of mulberry in, 626; pests of other fruits in, 174, 315, 594, 595; Nephantis serinopa on coconut in, 169; pests of coffee in, 37, 448; Oligonychus coffeae on tea in, 288; cotton pests in, 36, 37, 139, 140, 314, 317, 628; pests of brinjal in, 79, 497; pests of Hibiscus spp. in, 36, 314, 594; Syngamia, absuntalia 314, 594; Syngamia abruptalis on mint in, 424; pests of other vegetables in, 173, 317, 424, 495, 498, 499, 500, 693; Chrysocraspeda olearia on Eugenia cumini in, 88; forest pests in, 37, 88, 448, 552, 620; Hypera variabilis on lucerne in, 119, 317; Antigastra catalaunalis on sesame in. 316; Lepidoptera on Sesbania bispinosa in, 316; Scrobipalpa heliopa on tobacco in, 236; pests of ornamental plants in, 36, 37, 693; Acarapis woodi infesting honey bees in, 296; pests of stored products in, 117, 196, 315, 448, 499, 500, 594, 595, 626, 685; Necrobia rufipes in copra imported into Italy from, 436; locusts in, 116, 500; other insects in, 37, 173, 174, 293; natural enemies of pests in, 37, 66, 79, 91, 119, 169, 196, 236, 314, 316, 317, 406, 491, 497, 500, 626.

Indiana, Lepidoptera on cabbage in, 269; Noctuids on mint in, 284; virus disease of Noctuids in, 284; insecticidal activity of soil fungus in, 572.

indica, Atherigona; Margaronia (Diaphania).

indicola, Heterotermes.

indicus, Trinervitermes.

Indigofera suffruticosa, new Agromyzid on, in Queensland, 295.

indigoferae, Melanagromyza. Indochina (see Vietnam).

3-Indoleacetic Acid, in honeydew of aphids, 629; paper chromatography of, 629.

3-Indoleacetonitrile, in honeydew of Toxoptera graminum, 629; paper chromato-

graphy of, 629.

3-Indolebutyric Acid, in honeydew of aphids, 629; paper chromatography of, 629.

Indolepyruvic Acid, in honeydew of aphids, 629; paper chromatography of,

629.

Indonesia, study on Nacoleia octasema and its parasites on banana and other plants in, 353, 354; Diaspis boisduvalii on Cattleya in, 43; resistance of timbers from, to termites, 649. (See also constituent territories.)

Indophenyl Acetate, use of, in tests of esterase activity in eggs of Tetranychus

telarius, 278.

Indopol H-100, in dormant spray against

Psylla pyricola, 412.

Infra-red Gas Analyser, measurement of carbon-dioxide output of insects by, 40.

Infra-red Light, determination of insecti-

cides by, 561, 620.

Infra-red Spectrophotometry, in analysis of Thiodan residues, 92.

infuscatella, Chilotraea. infuscatus, Coccidencyrtus.

Inositol, and pigmentation of locusts, 5; in diet for Anthonomus grandis, 404.

inquisitor, Ephialtes.

Insecticides (including acaricides), standardisation of names for, 356, 497; transliterated list of Russian names for, 214; tests of organic compounds as potential, 407, 478, 479, 480, 515, 572, 634, 635; determination of contact toxicity of, 210; question of vapour toxicity of, 27; relation of structure to toxicity of, 64, 68, 445, 634, 635; comparison of potency of, 221; symposium on mode of action of, 62-64; detoxification and metabolism of, in insects, 64, 90; effects of sublethal doses of, on susceptibility of larvae of Lymantria dispar to polyhedral virus, 368; toxicology of toxaphene group of, 140; reviews and studies of resistance to, in arthropods, 128, 147-149, 252, 262, 284; determination of resistance to, 284, 680; toxicity of, to fish, 304; experiments on effects of, on growth and respiration of plants, 562; tainting of vegetables and fruits by, 228; ecological aspects of use of, 55, 262, 432, 433; hazards of, 40, 300, 432, 433; residual effectiveness of, 16, 444;

residues of, 300, 424; chromatography of residues of, 552, 577; lipid adsorptivity of dusts related to use as diluents for, 496; factors affecting concentration of, in sprays, 571; evaluation of wetting agents for, 140; biological tests of, 444, 599, 680; use of *Drosophila* for identification of, 304; bioassay of, 90, 300, 397; survey of use and development of, 301.

Insectorubin, effect of infection by Nosema locustae on deposition of, in

Locusta migratoria, 582.

Insects, natural enemies of, in tropical Africa, 653; part of a work on agricultural importance of, in Europe, Mediterranean basin and western Asia, 491; book on injurious species of, in Israel, 37; list of, damaging crop plants in Madagascar, 250; associated with dying cacao trees in New Guinea, 196; surveys of, on fodder crops in New Zealand, 75, 506; surveys of, infesting stored products in Pakistan, Persia and Turkey, 236; list of, intercepted in U.S.A., 299; surveys of data on, not occurring in U.S.A., 506-508; relation of, to outbreaks of virus diseases of plants, 271; review of chewing, as vectors of plant viruses, 443; review of damage to manufactured materials caused by, 489; methods for detection of infestation of stored cereals by, 571; estimation of populations of, 83, 84, 620; forecasting occurrence of, 15; estimation of mortalities in successive instars of, 590; vertical distribution of, 29, 30, 199, 578; mass migratory flights by. 199; factors affecting activity of, 30, 423; survival of, in nitrogen atmospheres, 324, 416; study of haemolymph of, in relation to physiological condition, 447; effects of electrostatic fields on, 123; effects of disease on pH of body fluids of, 161; review of bacteria against, 88; question of rôle of ingested antibacterial substances in protection of, against bacteria, 580; book on Microsporidia attacking, 89; book on parasites of leaf-mining, 492; bionomics of Coleoptera predacious on, 494; toxicity of turnip constituent to, 484; survey of measures against, 301; tests of aerosols for use against, in aircraft, 418; topical application of insecticides to, with microcapillary tubes, cuticular structure in relation to toxicity of insecticides to, 43, 314; statistical analysis of results of experiments

on control of, 399; apparatus for treatment of, with radioactive fumigants, 356; basis and development of resistance to insecticides in, 128, 252, 284; standardised methods for determination of resistance to insecticides in, 284; containers for transport of, 25, 496; apparatus for separating and collecting, 224, 291, 356, 694; anaesthetisation of, 356; removal of wax from, 496; cage for pollination studies on, 620; temporary storage of larvae of, 281; methods of rearing, 281, 282; pests infesting cultures of, 339; polyglot glossary of taxonomic terms for, 301; reviews of recent investigations on, 90. insertum, Rhopalosiphum.

insidiosa, auct., Pseudoperichaeta (see P.

roseanae).

insidiosus, Aphycus (Metaphycus); Erio-

phyes.

insolitus, Achrysopophagus; Centrococcus. instabilella, Scrobipalpa (Gnorimoschema). insulana, Earias.

integriceps, Eurygaster. intermedius, Dysdercus.

International Union for Conservation of Nature and Natural Resources, symposium at 8th technical meeting of, 262.

interpunctella, Plodia.

interruptor, Cremastus (Temelucha).

interruptus, Tydeus. intricata, Oncopera. inunctus, Tetrastichus.

invisor, Phaeogenes.

Iodine, ineffective with polyvinyl pyrrolidone, against Agrotis ypsilon, 211.

Ionisation Detector, determination of concentrations in fumigant mixtures

with, 619.

Iowa, Ostrinia nubilalis on maize in, 225, 276, 329; Nitidulid in relation to O. nubilalis and maize in, 329; nuisance value of mayfly and caddisflies in, 262. IPA (see Indolepyruvic Acid).

Iphiaulax, parasitising Oiketicus kirbyi in Costa Rica, 638; parasite of, 638.

Ipomoea carnea, Amblypelta lutescens

papuensis on, in Papua, 235.

Ips, on Pinus ponderosa in U.S.A., 460; as precursor of attack by Dendroctonus barberi, 460; methods of rearing, 390.

Ips concinnus, on pine in U.S.A., 386; mites destroying eggs of, 386.

Ips confusus, on pines in U.S.A., 340, 386; water supply and resin exudation in relation to infestation of pines by, 340; occurrence and determination of attractants for, 340, 341; other barkbeetles attracted by, 340, 341; natural

enemies of, 386, 387, 496; Serratia marcescens in cultures of, 300.

Ips curvidens, on white spruce in Slovakia, 283; new Microsporidian infecting, 283.

Ips guildi, attracted to logs infested by I. confusus in California, 341.

Ips latidens, attracted to logs infested by I. confusus in California, 341.

Ips oregonis, on pine in U.S.A., 386;

mites destroying eggs of, 386.

Ips pini, bionomics of, in pine logs in Canada, **122**, **123**; factors affecting populations of, **123**.

Ips plastographus, on pine in California, 386; mites destroying eggs of, 386.

Ips radiatae (see I. concinnus).

Ips tridens, method of rearing, 390.

Ips typographus (on conifers), in Czecho-

slovakia, 142; in France, 606; in Germany, 44, 386; characteristics and world distribution of, 507; water supply of trees in relation to infestation by, 253; natural enemies of, 142, 386; tests of attractants for, 44; measures against, 606; methods of rearing, 142.

Ips yohoensis, 390. Iran (see Persia).

Iraq, Eurygaster integriceps on cereals in, 655.

Iridomyrmex, preventing establishment of buffel grass in Queensland, 589; insecticides against, 589.

Iridomyrmex humilis, in Bermuda, 229; in California, 519; relation of, to Coccids and beneficial insects, 229, 519.

Iris, Anuraphis tulipae on, in British Columbia, 529.

Iris germanica, Rhadinoceraea micans on, in Yugoslavia, 32.

Iris pumila, Rhadinoceraea micans on, in Yugoslavia, 32.

Iron Sulphate, in combined sprays against *Rhagoletis pomonella*, **273**.

Irrigation, influence of, on systemic activity of insecticides, 508.

Isaria farinosa, infesting Pristiphora erich-

sonii in Michigan, 210.
Ischnurgops ornatus, sp.n., parasitising

Pachynematus scutellatus in Czecho-slovakia, 246.

Isle of Wight Disease (of honey bees), question of cause of, 296, 342.

Isobutyl p-Nitrophenyl Ethylphosphonate, anticholinesterase activity and toxicity of, to *Musca domestica*, 407.

Isodrin, toxicity of, to aphids, 498; against Coleoptera, 317, 323; toxicity of, to Locusta migratoria, 172; against

Spodoptera exempta, 179; against Trypetids, 268, 325; in dusts, 323; in granules, 323; in sprays, 179; soil treatment with, 325.

Isolan, against aphids, 19, 312, 667; against Fenusa pusilla, 19; against Eotetranychus carpini, 183; effects of, on parasite of Eriosoma lanigerum, 312; in sprays, 183, 312; trunk applications of, 19; translocation of, in birch, 19: colorimetric determination of, 227; and phenkapton, 183.

Isomera (see Cnephalia).

Iso-octyl Phenoxy Polyethoxy Ethanol

(see Triton X-100).

2-Isopropoxy-5-methoxyphenyl Dimethylcarbamate, insecticidal anticholinesterase activities of, and 635; and piperonyl butoxide, 635.

2-Isopropoxy-5-methoxyphenyl Methylcarbamate, insecticidal and anticholinesterase activities of, 634, 635; and

piperonyl butoxide, 635.

o-Isopropoxyphenyl Methylcarbamate, in dusts against Heliothis zea, 509; tests of toxicity of, to insects, 515, 516; and piperonyl butoxide, 516.

Isopropyl p-Nitrophenyl Ethylphosphonate, anticholinesterase activity and toxicity of, to Musca domestica, 407.

3-Isopropylphenyl Dimethylcarbamate, and insecticidal anticholinesterase activities of, 634, 635; hydrolysis of, 635; and piperonyl butoxide, 635.

3-(or m-)Isopropylphenyl Methylcarbamate, in dusts against Heliothis zea, **509**; in sprays against *Lygus lineolaris*, 515; toxicity of, to other insects, 476, 480, 515, 516; insecticidal and anticholinesterase activities of, 634, 635; hydrolysis of, 635; colorimetric determination of, 227; and synergists, 476, 635.

o-Isopropylphenyl Methylcarbamate. tests of toxicity of, to insects, 515.

2-Isovaleryl-1,3-indandione, laboratory tests of toxicity of, to insects, 202, 203; effect of temperature on toxicity of, 202, 203.

Israel, pests of cereals and grasses in, 349, 350; virus disease of maize in, 349, 350; pests of Citrus in, 22, 611, 612, 614, 684; Coccids on vines and other fruits in, 613, 614; Noctuids on cotton in, 351; Aulacophora foveicollis on cucurbits in, 613; Aphis gossypii and transmission of mosaic virus of lettuce and zinnia in, 614, 615; Myzus persicae and virus diseases of potato in, 614; vectors of virus diseases of sweet potato in, 130; book on injurious

insects and mites in, 37; relation of climate of, to development of Calliptamus palaestinensis, 142, 143; Byrsocrypta ulmi not recorded from, 349; natural enemies of pests in, 22, 613, 614. (See also Gaza Strip.)

Italy, Lepidoptera on almond in, 38, 181; pests of apple in, 136, 309, 533-535, 616: cherry pests in, 137, 311, 533; pests of cereals and grasses in, 182, 310, 349, 350, 684; virus diseases of cereals in, 80, 350, 684; Citrus pests in, 181, 299; fig mosaic in, 70; pests of peach in, 181, 310, 437, 616, 662; pests of other fruits in, 40, 299, 437; pests of olive in, 40, 46-48, 135, 308, 310, 438, 532, 694; pests of vines in, 40, 48, 183, 310, 660, 661, 663; Bruchidius perparvulus on crimson clover in, 182; forest pests in, 136, 137, 183, 535, 615, 660; pests of vegetables in, 310, 436, 531, 533, 661, 681; Monarthropalpus buxi on box in, 694; Coraebus rubi on roses in, 299; Pegomyia bicolor on begonia in, 661; other species of Pegomyia in, 662; danger of introduction of Hyphantria cunea into, 311; Reticulitermes lucifugus in, 649; Tyria jacobaeae in, 452; Dendroctonus obesus intercepted in, 532; Acarapis infesting honey bees in, 295; pests of stored products in, 182, 300, 426; beneficial insects in, 136, 137, 138, 182, 183, 308, 391, 436, 438, 532, 535, 616, 660; occurrence and transplantation of Formica spp. in forest areas of, 615, 616. Itoplectis alternans (parasitising Tortricids), in Czechoslovakia, 683; in

U.S.S.R., 598. Itoplectis conquisitor (parasitising Lepidoptera), in Minnesota, 212; in Ontario, 162; influence of diet on survival and fecundity of, 208, 209; effect of tree location on degree of parasitism by, 212; effects of electrostatic fields on, 123.

Itoplectis maculator (parasitising Tortricids), in Czechosovakia, 683: survey of data on, in France and Germany,

Ivory Coast, pests of cacao in, 7; coffee pests in, 7, 651; new weevil on Cola in, 355.

Ivy (see *Hedera helix*).

J

Jack Pine (see Pinus banksiana). Jackdaw (see Colaeus monedula). jacobaeae, Tyria (Hypocrita).

jacobi, Capitophorus (Pentatrichopus).

Jam, in diet for Vespula germanica, 422. Jamaica, pests of sugar-cane in, 289; Gryllus assimilis in, 567, 687; Tribolium in, 583; beneficial insects in, 289.

Japan, apple pests in, 35, 36; Coleoptera damaging persimmon in, 450; Eotetranychus smithi on vines in, 452; Pieris rapae on cabbage in, 78, 449, 450; pests of soy bean in, 452, 552; Tetranychids on other vegetables in, 452; rice pests in, 78, 451, 452; pests of other cereals and grasses in, 96, 97, 350, 684; virus diseases of cereals in, 350, 684; Tetranychids on tea in, 452; Paratetranychus ilicis on azalea in, 452; Diptera on pine mushroom in. 623; Drosophila virilis in lumber-yards and houses in, 622; distribution of Etiella zinckenella and Leguminivora glycinivorella in, 496; other insects in, 450, 451, 452, 567; natural enemies of pests in, 36, 78, 97, 451, 491; wilt disease of *Pieris rapae* in, 78.

Japananus hyalinus, on Norway maple and sour cherry in Ontario, 529.

Japanese Larch (see *Larix leptolepis*). Japanese Plum (see Plum, Japanese).

Japanese Red Pine (see Pinus densiflora).

japonica, Popillia.

japonicum, Trichogramma.

japonicus, Ceroplastes floridensis.

Java, *Xyleborus morigerus* on miscellaneous plants in, **625**; termites infesting timber and furniture in, **296**.

Jeffrey Pine (see Pinus jeffreyi).

jeffreyi, Dendroctonus.

Jordan, vectors of virus diseases of cereals in, 684.

jucundus, Apanteles.

juglandicola, Chromaphis.

Juglans major, Dasiops alveofrons reared from, in Arizona, 128.

juglansregiae, Quadraspidiotus (Aspidiotus).

juncii, Lixus.

Juniper (see Juniperus communis).

Juniper Webworm (see *Dichomeris mar-ginella*).

juniperi, Monoctenus.

Juniperus bermudiana, Carulaspis minima

on, in Bermuda, 229.

Juniperus communis, nature of injury by Monoctenus juniperi to, in Holland, 302, 303; Dichomeris marginella on, in New York, 19; activity of systemic insecticides in, 19.

junodi, Kotochalia.

Jute, pests of, in British Guiana, 27, 28.

K

Kakimia robinsoni, on Delphinium in Manitoba, 525.

Kakothrips pisivorus (robustus), bionomics of, on pea in Holland, 193; characteristics and world distribution of, 507; measures against, 193.

Kale, insects on, in New Zealand, 74, 75, 505, 506; pests of, in Virginia, 404.

kalmii, Lygus.

Kalotermes flavicollis, resistance of timbers to infestation by, 649.

Kalotermes tabogae, infesting timber in British Honduras, 76.

kanpurensis, Rogas.

Kansas, pests of cereals in, 65, 276, 575; Therioaphis maculata on lucerne in, 468, 642.

kanzawai, Tetranychus.

Kaolin, against pests of stored products, 51; as carrier for insecticide dusts, 51; effect of, on toxicity of dusts, 118.

Karo Solution, use of, in determination of insecticide residues in soils, 339.

Kelthane, Chrysomphalus ficus favoured by, 645; toxicity of, to Cydia pomonella, 21; against Therioaphis maculata, 468; tests and uses of, against Tetranychus spp., 217, 281, 324, 603, 604, 613, 633, 684; against other Tetranychids, 159, 183, 288, 289, 419, 517, 628, 672; susceptibility or resistance to, in strains of Tetranychids, 193, 266, 278, 510, 512; against other mites, 120, 532, 545, 556, 560, 645, 664; effects of, on beneficial arthropods, 159, 627; effects of, on crop yields, 281, 645; effects of, on colour of poinsettia, 517; in dusts, 217; in sprays, 120, 183, 217, 266, 288, 289, 324, 419, 468, 532, 545, 556, 560, 633, 645, 664, 672; persistence of, on apple leaves, 672; toxicity of DDT increased by, 21; and TEPP, 532; and parathion, 532.

Kentucky, Homadaula albizziae on ornamental trees in, 273; pests of strawberry in, 570; tolerance limits for

insecticide residues in, 570.

Kentucky Bluegrass (see *Poa pratensis*). Kenya, pests of cereals in, 179, 180, 530, 684; *Spodoptera exempta* on pastures in, 179, 530, 686; atlas of pests and diseases of coffee in, 490; pests of stored maize in, 51; *Tribolium* in, 583; parasites of pests in, 180, 531; virus diseases of *S. exempta* in, 531, 686.

Kepone (see 2,3,3a,4,5,6,7,7a,8,8-Decachloro-3a,4,7,7a-tetrahydro-4,7-methanoindene-1-one).

Keratin, digestion of, by larvae of Hofmannophila pseudospretella, Kerosene, soil treatment with emulsion of, against Lachnosterna consanguinea, 171; use of, in topical applications of insecticides and synergists, 687. Khlorofos (see Trichlorphon). kirbyi, Oiketicus. kissophila, Bryobia. kivuensis, Anagyrus. Koeleria cristata, relation of mites and thrips to silver top of, in Alberta, 156. koenigii, Dysdercus. kolobovae, Eurytoma (Bruchophagus). Korscheltellus (see Hepialus). predacious Korynetes caeruleus, Anobium punctatum in Britain, 489. Kotochalia junodi, characteristics and world distribution of, 507. krameri, Leiodinychus. kuehniella, Anagasta (Ephestia). kurdjumovi, Rungsia.

L

Labrorychus, parasitising Lepidoptera in Canada, 27, 162.

Lachnosterna, infection of, by Cordyceps ravenelii in Canada, 619; in U.S.A., 420, 570; on maize, 420; on strawberry, 570; insecticides against, 420, 570.

Lachnosterna bruneri, in Florida, 495. Lachnosterna consanguinea, bionomics and light-trap catches of, on sugarcane in India, 171; insecticides against, 171.

Lachnosterna diomphalia, bionomics of, on soy bean and other crops in China, 593, 594.

Lachosterna morosa, on cereals and other crops in China, 593.

lacteicolor, Apanteles.

kuvanae, Ooencyrtus.

kuwanae, Chilocorus.

Laelia adalia, 79.

Laelia exclamationis, bionomics and parasites of, on Guinea grass in India, 79; rice and ragi not attacked by, 79; insecticides against, 79.

laensis, Pseudodoniella.

laevigata, Atta.

laevigatum, Stenodema.

laevigatus, Alphitobius.

laingi, Eumargarodes.

lalori, Apleurotropis.

Lambdina fiscellaria lugubrosa, on Tsuga heterophylla in British Columbia, 395; aerial spray of DDT against, 395.

Lambdina fiscellaria somniaria, laboratory

experiments with polyhedrosis virus and. 581.

lambens, Sarcophaga (Sarcodexia).

Lampetia (see Merodon).

Lampides boeticus (see Cosmolyce).

lanata, Chrysopa.

lanigerum, Eriosoma.

lapathi, Cryptorhynchus.

Laphygma (see Spodoptera). laphygmae, Euplectrus.

Larch, Enarmonia griseana on, in Austria and Switzerland, 100; Pristiphora erichsonii on, in Canada, 124, 125, 210; Formica spp. in forests of, in Italy, 615: Dendrolimus superans and secondary pests on, in Mongolia, 116; pests of, in U.S.S.R., 112, 244; pests of, in U.S.A., 210, 474.

Larch, Eastern (see Larix laricina). Larch, Japanese (see Larix leptolepis). Larch, Siberian (see Larix sibirica).

laricicola, Hylemya.

laricis, Chermes (Adelges). laricivorus, Taeniothrips.

Laricobius erichsonii (predacious Chermes piceae), introduced into Canada, 456, 522; characters of, 522.

Laricobius rubidus, in Canada, 522; characters distinguishing L. erichsonii from. 522.

Larix (see Larch).

Larix decidua (europaea), Formica spp.

in forests of, in Italy, 615.

Larix sibirica, Dendrolimus superans and secondary pests on, in Mongolia, 116. Larix laricina, Pristiphora erichsonii on, in Canada, 456; Choristoneura pinus

on, in Minnesota, 474.

Larix leptolepis, Pristiphora erichsonii on, in Pennsylvania, 210.

Larvacarus transitans, on Ziziphus jujuba in Pakistan, 176, 626; bionomics of. 176; sprays against, 626.

larvarum, Eulophus (Cratotechus); Tachina

(Exorista).

Lasioderma serricorne, in S. Africa, 650; bionomics of, in Bulgaria, 57; intercepted in Germany, 11; in India, 117; in stored tobacco, 57, 117, 650; on oil-seed cake, 11; effects of nicotine on symbiont of, 650; effects of redrying of tobacco on populations of, 117; measures against, 58.

Lasioglossum, predacious on larvae of Neodiprion swainei in Quebec, 73.

Lasiophthicus pyrastri, predacious aphids in Yugoslavia, 240.

Laspeyresia (see Cydia). lata, Amorphoidea. lataniae, Hemiberlesia.

latens, Petrobia.

lateralis, Phytomyza(Napomyza).

Latheticus orvzae (in stored cereals), in India, 315, 626; new Tenebrionid predacious on, 626; experiments on competition between Tribolium confusum and, 89.

Lathromeris senex, egg-parasite of Bruchus

pisorum in Bulgaria, 58.

Lathyrus pratensis, Sitona suturalis on, in Finland, 102.

latidens, Ips.

latungula, Parabates (Netelia).

latus, Hemitarsonemus.

lauri, Hispaniella.

lavandulae, Thomasiniana.

Lavender, Thomasiniana lavandulae on, in France, 436, 606.

Lawn Armyworm (see Spodoptera mauritia).

Lawns (see Grasses).

Lawsonia alba, Perissopneumon tectonae

on, in India, 174.

Lead Arsenate, in dusts, 138; in sprays: against cotton pests, 559; against orchard pests, 18, 159, 239, 268, 273, **274, 637**; against pests of vines, **9, 48**; against Abraxas miranda on Euonymus, 503; against Lozotaeniodes cupressana on cypress, 138; ineffective against Vespula germanica, 422; effects of, on beneficial arthropods, 36; effects of, soil, 83; and azinphos-methyl, 268; and bordeaux mixture, 36; toxicity of, increased by captan, 274; and DDT, 268; and ethyl-DDD, 268; and fungicides, 36; and lime, 48; and limesulphur, 36; and methyl-demeton, 559; and Sevin, 268.

Lead Cable, damaged by Dicranura erminea menciana in China, 693.

Lebanon, new egg-parasite of Eurygaster integriceps in, 655.

Lebaycid (see Fenthion). Lecanium (see Eulecanium).

Lechriolepis nephopyropa, on pines in S. Rhodesia, 81.

lecontei, Agonoderus; Neodiprion.

lefebvrii, Glenea.

lefroyi, Taeniothrips. leguminicola, Dasyneura.

Leguminivora glycinivorella, in Japan, 496; characteristics and world distri-

bution of, 507; cold-hardiness of overwintered larvae of, 424.

Leiodinychus krameri, in stored grain in U.S.A., 518; food preferences of, 518. Lema melanopa, characteristics and world distribution of, 507.

lemniscata, Comperiella.

Lemon, Myllocerus evasus on, in India,

173; pests of, in Israel, 611, 614; pests of, in Italy, 181, 299; insecticide residues in fruits of, 561, 573, 574.

Lemon Packing Houses, Dibrom fumigation of, against Drosophila, 573.

Lentinus edodes, 623.

lentis, Bruchus (Bruchidius).

Leperisinus (see Hylesinus).

Lepidiota, on cacao in Dutch New Guinea, 76.

Lepidosaphes, new parasite of species of, on Citrus in India, 491.

Lepidosaphes beckii (on Citrus), in Brazil, 120; in India, 491; in Morocco, 382; Texas, 564; bionomics of, 382; parasites of, 382, 491, 564; sprays against, 120.

Lepidosaphes gloverii, map of distribution of, 357.

Lepidosaphes pistaciae, on pistachio in Turkey, 609.

lepidosaphes, Aphytis.

Lepisma saccharina, damaging textiles in Germany, 377.

Leptinins, experiments with, on resistance of Solanum to Leptinotarsa, 318.

Leptinotarsa decemlineata (Potato Beetle), distribution of, in Europe in 1961, 681; in Belgium, 681, 683; in Bulgaria, 366, 539; in Canada, 207, 377; in Czechoslovakia, 247, 681; in Germany, 252, 377, 440, 543, 684; in Hungary, 305; in Italy, 533, 681; in Majorca, 681; in Poland, 366, 681, 683; in Rumania, 366, 681; in U.S.S.R., 15, 447, 543, 681; in U.S.A., 325, 377; in Yugoslavia, 366, 681; map of distribution of, 357; on other solanaceous plants, 366; rôle of chemical factors in food preferences of, 318, 608; as vector of bacterial wilt of banana, 520; diapause in, 6, 204, 543; annual cycle of, 366; relation between soil type and natural mortality of, 683; natural enemies of, 207, 208, 377, 684; penetration of Beauveria bassiana into larvae of. 196: combined effects of B. bassiana and DDT on, 447; tests of repellents against adults and larvae of, 305; tests and uses of insecticides against, 247, 252, 310, 325, 366, 533, 539, 683; susceptibility or resistance to insecticides in strains of, 252, 363, 364, 365, 440.

Leptins, experiments with, on resistance of Solanum to Leptinotarsa, 318.

Leptoconops, review of bionomics and control of, 90.

Leptocorisa varicornis, measures against, on rice in India, 316.

Leptoglossus australis, bionomics and food-plants of, in New Guinea and Pacific Islands, 235; sprays against, 235.

Leptohylemyia (see Hylemya).

Leptomastidea abnormis, parasitising Planococcus citri in Israel, 614.

Leptomastix dactylopii, parasitising Planococcus citri, 614; introduced into Israel from Canada, 614.

leptospermi, Eriococcus.

Leptospermum, Coccids on, in New Zealand. 587, 588.

Leptospermum ericoides, resistance of, to Eriococcus orariensis in New Zealand, 588.

Leptospermum scoparium, Coccids and biological control of, in New Zealand, 587, 588.

lepturoides, Omophlus.

lesbia, Colias.

Lethane 384 (see β-Butoxy β'-Thio-

cyanodiethyl Ether).

Lethrus apterus, on plants in Bulgaria, 58; bionomics of, 58; measures against, 58.

Lethum australiense, 345.

Lethum australiense var. necrotica n., 345.

Lethum rhizospilum, 133.

Lettuce, Hepialus lupulinus on, in Britain, 431; Macrosteles fascifrons on, in Canada, 70, 524; Aphis gossypii on, in Israel, 614, 615; Penthaleus major on, in Italy, 531; pests of, in U.S.A., 131, 470, 514, 569; transmission of virus diseases of, 70, 131, 467, 470, 524, 614, 615; treatments of, with systemic insecticides, 470.

Leucania loreyi, dusts against, on maize in U.S.S.R., 345.

Leucinodes orbonalis, on brinjal in India, 79; characteristics and world distribution of, 507; sprays against, 79, 80.

Leucophaea maderae, survival of, in various atmospheres in relation to space-vehicle use, 324, 325, 416; absorption, metabolism and excretion of insecticides by, 43, 248.

leucophthalmus, Philaenus.

Leucopis, predacious on aphids in Nova Scotia, 388.

Leucopis americana, predacious on aphids in Nova Scotia, 388.

Leucopis griseola, predacious on Aphis fabae in Bulgaria, 537.

Leucopis obscura, predacious on Chermes piceae, 456; introduced into Newfoundland, 456.

Leucopoecila (see Spanogonicus).

Leucoptera caffeina, on coffee in Tanganyika, 652, 653; parasites of, 652,

653; effects of sprays on populations of, 653.

Leucoptera merycki, on coffee in Tanganyika, 652, 653; parasites of, 652, 653; effects of sprays on populations of, 653.

Leucoptera scitella (on apple), in China, 625; in Italy, 136, 309, 533, 534; bionomics of, 136, 534; parasites of, 136; insecticides against, 136, 534.

leucopterae, Parahormius.

leucopterus, Blissus.

leucotreta, Cryptophlebia (Argyroploce). lewisi, Eotetranychus.

lichtensiae, Aphycus (Metaphycus).

Life Tables, sampling methods for, for

Ostrinia nubilalis, 258.

Light, effects of: on food-plant selection by aphids, 200, 201; on movements of first-instar nymphs of Eriosoma lanigerum, 481; on Oncopeltus fasciatus, 518; on egg production by Trialeurodes vaporariorum, 198; on resistance of wheat varieties to Cephus cinctus, 526; on diapause in Cladius pectinicornis, 361; on populations of Curculionids, 206, 207, 318, 319; on parasitism of Dendrolimus punctatus, 624; on moulting of larvae of Hyphantria cunea, 241; on habits of Oecophylla, 29; on behaviour of Oscinella frit, 446. (See also Infra-red Light, Photoperiod Ultraviolet Light.)

Light-traps, for Hemiptera, 316, 372; for Lachnosterna consanguinea, 171; for Lepidoptera, 84, 165, 219, 220, 248, 292, 306, 334, 335, 338, 477, 630; predators of Pectinophora gossypiella attracted to, 338; machine for changing position of, 551; types of light for, 165, 334, 338, 476, 477, 630; types of, 219, 220; windbreaks used with, 220. ligustici, Otiorhynchus (Brachyrhinus).

Lily, as reservoir of plant viruses, 271. (See also *Lilium* and *Zantedeschia*.) Lily, Arum (see *Zantedeschia aethiopica*). Lily, Easter (see *Lilium longiflorum*).

Lilium longiflorum, aphids on, in Oregon, 278; treatment of bulbs of, 278.

Lima Bean Pod Borer (see Etiella zinckenella).

Lima Beans, pests of, in Maryland, 325; susceptibility of strains of, to Epilachna varivestis and Empoasca fabae, 414; use of, in experiments with E. varivestis and Tetranychids, 272, 278, 516, 519; systemic activity of insecticides in, 325.

limacina, Caliroa.

limbata, Gynandrophthalma.

Lime (Citrus), tests with aphids and tristeza disease of, in India, 80.

Lime, against ants, 589; against Lepidoptera, 48, 80, 181; effects of, on populations of orchard pests and their natural enemies, 36; effects of, on persistence of insecticides, 83, 641; in sprays, 36, 48, 80; seed treatment with, 589; trunk application of, 181; and calcium arsenate, 80; and lead arsenate. 48.

Lime-sulphur, against Cydia pomonella. 20; against Psylla pyricola, 411; against Quadraspidiotus perniciosus, 167, 168; against Rhagoletis pomonella, 273; against Thysanoptera, 71; against Eriophyids, 287, 298, 432; against Tetranychids, 183, 189, 288, 289, 544, 678; against other mites, 28, 626; effects of, on beneficial arthropods, 35, 36, 672; toxicity of, to black currant, 287; hazards of use of, 28; in dusts, 71; in dormant sprays, 411; in sprays, 20, 28, 35, 36, 167, 168, 183, 189, 273, 287, 288, 298, 432, 544, 626, 670, 672, 678; comparison of highand low-volume applications of, 670; soil treatment with, 71; factors affecting evolution of hydrogen sulphide from 672; and chlorobenzilate, 36; and DDT, 20, 36, 544; and lead arsenate, 36; and nicotine sulphate, 35; and oil emulsion, 411; and sulphur, 287; and tar distillate, 189.

Limnerium (see Campoplex). limonicus, Typhlodromus.

Limonius agonus, in Connecticut, 582; destroyed by bacterial toxins, 582.

Limothrips, in Alberta, 156; relation of, to silver top of grasses, 156.

Limothrips cerealium, 156. Limothrips denticornis, 156.

Linaria vulgaris, question of control of, by Coleoptera in Canada, 386.

Lindingaspis rossi, on Leptospermum in New Zealand, 588.

linearis, Atomaria; Spintherus. lineata, Celerio.

lineatella, Anarsia.

lineaticollis, Antestiopsis.

lineatum, Trypodendron (Xyloterus).

lineatus, Sitona.

lineola, Thymelicus.

lineolaris, Lygus (Liocoris).

lineolatus, Adelphocoris.

lingnanensis, Aphytis.

lini, Thrips.

Linoleic Acid, in diets for insects, 5, 627.

Linseed Oil, tests of methyl esters of

fatty acids of, as attractants for barkbeetles, 44.

Liocoris (see Lygus).

Liosomaphis (see Rhopalosiphum).

liparidis, Apanteles.

Lipase, rôle of, in nutrition of Agrotis orthogonia, 69.

Liposcelis (see Troctes).

Liriomyza, species of, on tomato in California, 508; Zectran against, 508. Liriomyza cicerina, 351.

Lissonota biguttata, parasitising larvae of Operophtera brumata in Europe, 378; introduced into Canada, 378.

Lissonota buolianae, parasitising Lepidoptera in Czechoslovakia, 602, 603.

Lissonota nigra, parasitising Eucosma nigricana in Czechoslovakia, 602.

Lissorhoptrus oryzophilus, on rice in U.S.A., 329, 421; insecticides against, 329, 421.

Listroderes, parasites of, in S. America, 353.

Listronotus oregonensis, on carrots in Ontario, 529.

Litchi (Litchi chinensis), pests of, in Hawaii, 463; Aceria sp. on, in Pakistan, 88; tests of wetting agents in relation to toxicity to, 463.

litchii, Aceria.

Lithocolletis, on apple in Italy, 309.

Lithocolletis blancardella, in Italy, 533-535; in Switzerland, 86; bionomics of, on fruit trees, 86, 534; parasites of, 86; insecticides against, 86, 534, 535.

Lithocolletis corylifoliella, on fruit trees and other plants in Switzerland, 86; bionomics and parasites of, 86; effects of sprays on populations of, 86.

Lithocolletis platani, bionomics and parasites of, on plane in Yugoslavia, 31; insecticides against, 31.

Lithuania, Leptinotarsa decemlineata on potato in, 15.

litigiosus, Agriotes.

Litomastix truncatella, attacking Trichoplusia ni in New York, 401.

littoralis, Scaphoideus.

litura, Prodenia.

liturata, Hylemya (see H. trichodactyla). livornica, Celerio lineata.

Lixophaga diatraeae, parasitising Dia-traea saccharalis in Jamaica, 289; effects of sprays on populations of,

Lixus juncii, characteristics and world distribution of, 507.

Lizards, destroying natural enemies of Pseudaulacaspis pentagona in Bermuda 229; destroying noxious insects, 166

638: bird introduced into Bermuda

against, 229.

Lobesia botrana (on vines), in Bulgaria, 60; in France, 8, 9, 494, 667, 668; in Germany, 108; characteristics and world distribution of, 507; Bacillus thuringiensis ineffective against, 9; insecticides against, 9, 60, 108, 494, 667, 668; effects on mites of sprays against, 8.

lobosus, Tetranychus.

Locust, Desert (see Schistocerca gregaria). Locust, Moraine (see Gleditsia).

Locusta migratoria, foci and forms of, in Sinking, 13: migration of, to U.S.S.R., 13: nutritional requirements of, 5, 6, 172; cannibalism among hoppers of, 172; toxicity of insecticides to, 172; method of rearing, 606.

Locusta migratoria capito, bionomics of,

in Madagascar, 297.

Locusta migratoria manilensis, 13.

Locusta migratoria migratoria, U.S.S.R., 15; characters of breeding areas of, 13, 15.

Locusta migratoria migratorioides, in Mali, 98, 620; factors affecting development and dispersal of, 98, 285, 286; phases of, 620; estimation of mortalities in successive instars of, 590; sensory equipment of, 423; effects of Nosema locustae on pigmentation and flight of, 582; factors affecting resistance to insecticides in, 201.

locustae, Nosema.

Locusts, in Argentina, 356; in Mali, 620; forecasting abundance of, in U.S.S.R., 15; factors affecting populations of, 98, 285, 286; sensory equipment of, 423; flight habits and dispersal of, 199, 423, 500, 620; studies on nutritional requirements of, 5, 6; factors affecting pigmentation of, 5; cannibalism among hoppers of, 172; natural enemies of, 313, 356; infected by Nosema locustae, 582; pick-up of spray droplets by, 291; factors affecting toxicity of sprays to, 657; factors affecting resistance to insecticides in, 201; metabolism and excretion of insecticides by, 321, 322; methods of rearing and breeding, 40; cages for, (See also Anacridium, Locusta, Nomadacris and Schistocerca.)

Lodgepole Pine (see Pinus contorta).

loeflingiana, Tortrix.

Loganberry, Byturus urbanus on, Britain, 671; Cicadellids on. in British Columbia, 395; insecticide residues on, 395, 671.

Lolium, Eurygaster spp. on, in California, 520; stem-boring Diptera on, in England, 487, 488; Oscinella frit on, in Germany, 45: Hyperodes bonariensis on, in New Zealand, 506; survival of Chloriona orizicola on, 132.

Lolium multiflorum, Abacarus hystrix transmitting mosaic virus of, in Europe,

Lolium perenne, Abacarus hystrix transmitting mosaic virus of, in Europe, 71; Oscinella frit on, in Britain, 430; pests of, in Canada, 162, 585; Rhopalosiphum padi on, in Wisconsin, 95.

Lombardy Poplar (see *Populus italica*). Long Ashton Research Station, review of

work of, 301.

longicauda, Apanteles. longicornis, Diabrotica. longinoda, Oecophylla. longior, Tyrophagus.

longipes, Anoplolepis.

longipilis, Typhlodromus.

longissima, Brontispa.

Longitarsus waterhousei, on Mentha spp. Washington, 324; insecticides against, 324.

longulus, Pnigalio (Eulophus).

loreyi, Leucania.

Lotus corniculatus, Eurytoma kolobovae infesting seeds of, in Hungary, 306; pests of, in U.S.A., 213, 564.

Louisiana, cotton pests in, 407, 466, 467; Chloriona orizicola and diseases of rice in, 405, 508; pests of sugar-cane in, 216, 219, 280, 469; pests and virus disease of sweet potato in, 131; Acarapis spp. infesting honey bees in, 295; Coleoptera in stored cereals in, 402, 403, 571; other insects in, 563.

lounsburyi, Aspidiotiphagus.

Lounsburyia, gen.n., Coccophagus trifasciatus as type species of, 490, 491. Lounsburyia affinis, sp.n., parasitising Saissetia in Zanzibar, 491.

Lounsburyia trifasciatus, introduced from S. Africa into California against Saissetia oleae, 491.

Louse, Body (see *Pediculus humanus* humanus).

loxonephes, Eulia.

Loxostege commixtalis, sprays against, on beet in Nebraska, 335.

Loxostege similalis, sprays against, on cotton in Texas, 16.

Loxostege sticticalis, sprays against, on beet in Nebraska, 335.

Lozotaeniodes cupressana, bionomics and parasites of, on cypress in Italy, 137, 138; survey of data on, 137; insecticides against, 138.

lubricipeda, Diacrisia (Spilosoma).

Lucerne, pests and pollinators of, in S. Australia, 233, 234; pests of, in Bulgaria, 59, 60, 537, 538, 539; pests of, in Canada, 72, 210; Contarinia medicaginis on, in Czechoslovakia, 187; Sitona spp. on, in Finland, 41, 102; pests of, in Hungary, 306, 307; Hypera variabilis on, in India, 119, 317; survey of insects on, in New Zealand, 75, 506; pests of, in S. Rhodesia, 48, 88; pests of, in U.S.A., 66, 67, 92, 134, 280, 331, 400, 462, 468, 521, 564, 576, 581, 631, 636, 639, 642; effects of, on development of Agrotis ypsilon, 346; populations of Otiorhynchus sulcatus on vines unaffected by presence of, 255; comparison of injury to, by Therioaphis maculata and Macrosiphum pisum, 322; varietal susceptibility of, to aphids, 629, 642; effect of irrigation on response of, to infestation by M. pisum, 210; detection of growth inhibitors in, 22; pollination of, 234, 411, 565; suction machine for sampling insect fauna of, 224; biological control of pests of, 199; seed treatment of, 468; residues of insecticides on, 92, 323, 324, 330, 331; influence of, on distribution and persistence of insecticides in soil, 573, 578; use of, in rearing Colias eurytheme, 281.

Lucerne Gall-midge (see Contarinia medi-

caginis).

Lucerne Meal, in baits for *Drosophila*, 327.

lucifugus, Reticulitermes.

Lucite, use of, in rearing *Ips*, 390.

ludens, Anastrepha. lugubris, Formica.

lugubrosa, Lambdina fiscellaria.

Lumbricus terrestris, insecticide residues in, 641.

lupulinus, Hepialus (Korscheltellus).

lurida, Scotinophara.

luteola, Galerucella.

luteolus, Aphycus (Metaphycus); Scaphoideus.

lutescens, Amblypelta. lybica, Empoasca.

Lychee (see Litchi).

lycimnia, Coccophagus.

lycopersici, Aculus (Vasates).

Lyctus, survey of data on treatments against, infesting building timber in Europe, 646; infesting oak timber, 622; starch content of sapwood related to susceptibility of timber to, 622.

Lyctus brunneus, in Eucalyptus and Tristania spp. in Queensland, 589; starch content of trees in relation to

infestation by, 589; treatments of timber against, 489, 648; method of rearing, 489.

Lydella stabulans grisescens (parasitising Ostrinia nubilalis), introduced into Delaware, 509; in France, 372; bionomics of, in U.S.S.R., 245, 246.

Lygaeus pandurus, bionomics of, on fruit trees, etc., in India, 315.

Lygus, in Arizona, 638, 639; in Finland, 103; in Uganda, 652; on beet, 103; on cotton 638, 652; natural enemies of, 638, 639; insecticides against, 639, 652.

Lygus borealis, sprays against, on lucerne in Canada, 72.

Lygus elisus, soil treatments against, on cotton in New Mexico, 409.

Lygus hesperus (on cotton in U.S.A.), insecticides against, 409, 478, 552.

Lygus kalmii, on cotton in Turkey, 352. Lygus lineolaris, in Canada, 72; in U.S.A., 213, 409, 467, 515, 570; on cotton, 409, 467, 515; on Lotus corniculatus, 213; on lucerne, 72; on strawberry, 570; insecticides against, 72, 213, 409, 467, 515, 570.

Lygus pratensis, on cotton in Turkey, 82, 352.

Lygus rufidorsus, sprays against, on lucerne in Canada, 72.

Lygus rugulipennis, on beet in Finland, 103; damage to beet by, in relation to seedling age, 103.

Lygus unctuosus, sprays against, on lucerne in Canada, 72.

Lymaenon, species of, parasitising leafhoppers in California, 630.

Lymantria dispar, in Czechoslovakia, 683; in Germany, 253, 376; in Japan, 450; in Sardinia, 660; in Spain, 381, 659; in U.S.S.R., 248, 368, 598; in U.S.A., 92, 220, 276, 399, 400, 513; in Yugoslavia, 54, 55, 367-371; on forest trees, 54, 55, 253, 376, 513, 514, 598, 659, 660, 683; on fruit trees, 54, 55, 514; on herbaceous plants, 55; symposium on bionomics and ecology of, 54, 55; factors affecting stages of, 368, 369, 450; pupal weight related to number of eggs produced by, 598; sex ratio of, 369; effects of fertilisers on populations of, 253, 376; fluctuations in populations of, 367, 369, 370, 371; natural enemies of, 54, 55, 367, 381, 399, 400, 660, 683; diseases of, 54, 55, 367, 368, 369, 400, 690; effects of insecticides on susceptibility of, to viruses, 368; attractants for, 55; female sex attractant of, 55, 368, 369,

370; traps for, 248, 368, 369; insecticides against, 55, 92, 220, 276, 399, 513, 514, 659; strains and races of, 54,

369, 370,

Lymantria monacha (on forest trees), in Germany, 253; in Spain, 381, 659; in U.S.S.R., 243; characteristics and world distribution of, 507; factors affecting population fluctuations in, 243, 253; natural enemies of, 381; DDT dust against, 659.

Lyonetia clerkella, in Switzerland, 86, 663; on apple, 663; on cherry, 663;

sprays against, 86, 663.

parasitising larvae of Lypha dubia, Operophtera brumata in Europe, 378; introduced into Canada, 378.

Lypha fumipennis, parasitising larvae of Choristoneura fumiferana in British

Columbia, 457.

M

mcdanieli, Tetranychus. machaon, Papilio. mckenziei, Vasates.

Maclura, Eulecanium corni on, in U.S.S.R., 13.

Maconellicoccus hirsutus, characteristics

and world distribution of, 507. Macquartia, parasitising Ambrostoma

quadriimpressum in China, 593. Macrocentrus ancylivorus, parasitising Cydia molesta, 665; introduced into

France, 665. gifuensis, Macrocentrus parasitising Ostrinia nubilalis in Delaware, 509.

Macrocentrus homonae, introduced into Ceylon for control of Homona coffearia, 553; effects of dieldrin sprays on populations of, 554.

Macrocoma brunnipes, on cotton

Turkey, 177.

Macrocoma brunnipes obscuricolor. on cotton in Turkey, 177.

Macroplectra nararia (see Natada).

Macropsis fuscula, on loganberry in British Columbia, 395; characteristics and world distribution of, 507; sprays against, 395.

Macrosiphum, on wheat in China, 625. Macrosiphum avenae, in Belgium, 375; in China, 625; in New Brunswick, 585; in U.S.A., 131, 160, 264, 401, 570, 639; on cereals, 132, 160, 375, 401, 585, 625, 639; virus diseases of cereals transmitted by, 132, 160, 375, 401; on grasses, 585; on beans, 264; on potato, 570; transmitting virus disease of beet, lettuce and radish, 131; injection of, with plant viruses, 133; infested by Entomophthora ignobilis, 570; trap ineffective for, 639; seed treatment against, 401.

Macrosiphum dirhodum, in New Brunswick, 585; in U.S.A., 131, 160, 401; on cereals, 160, 401, 585; relation of, to yellow-dwarf virus, 401; on grasses, 160; transmitting virus disease of beet, lettuce and radish, 131; comparison of Canadian and European strains of, 585; seed treatment against, 401.

Macrosiphum euphorbiae, in Australia, 34; in Brazil, 556; in Britain, 431; in New Brunswick, 633; in Poland, 544; in U.S.A., 131, 133, 278, 461, 510, 569, 573, 633; on beet, 17; on broad bean, 34; on carnation, 431; on Easter lily, 278; on potato, 461, 510, 544, 556, 569, 633; on sweet potato, 131, 133; on roses, 461, 633; on tomato, 573; transmitting virus diseases of plants, 17, 35, 131, 133, 544; forecasting of outbreaks of, 460; infested by *Ento*mophthora spp., 569, 570; insecticides against, 278, 431, 510, 556, 573, 633.

Macrosiphum fragariae, on Rubus procerus in British Columbia, 529; on strawberry in Germany, 106; question of transmission of plant viruses by, 107,

529.

Macrosiphum granarium (see M. avenae). Macrosiphum malvae rogersi (on strawberry), in Germany, 106; in Holland, 192; transmission of strawberry virus to Fragaria vesca by, 107.

Macrosiphum martorelli, sp.n., on cacao in Dominican Republic, 144; ants associated with, 144.

Macrosiphum pisum (pisi), in Alberta, 210; in U.S.A., 66, 211, 264; on beet, 17; on beans, 264, 603, 605, 629; on broad beans, 525; on lucerne, 66. 210, 629; comparison of damage to lucerne by Therioaphis maculata and, 322; effects of pea varieties on weight and reproduction of, 453; on hairy vetch, 211; and virus diseases of plants, 17, 264; varietal susceptibility of plants to, 551, 629; plant-growth regulators in relation to, 525, 551, 629; parasites of, 66; insecticides against, 210, 603, 605; diverse effects of sesamex on toxicity of insecticides to, 64; effects of insecticides against cereal pests on populations of, 211; method of rearing, 606.

Macrosiphum rosae, in British Columbia, 692; on roses in Maryland, 633; transmission of strawberry viruses by,

692; sprays against, 633.

Macrosiphum scoliopi, on Easter lily in Oregon, 278; systemic treatments

against, 278.

Macrosiphum solani, in British Columbia, 692; in Poland, 544; in U.S.A., 278, 461; on Easter lily, 278; on potato, 461, 544, 633; and virus diseases of potato, 544; transmission of strawberry viruses by, 692; oviposition habits of, 461; insecticides against, 278, 633.

Macrosiphum solanifolii (see M. euphor-

biae).

Macrosteles fascifrons, in Canada, 70, 524, 551; in U.S.A., 17, 470; on beans, 17; on other vegetables, 70, 470, 524; on cereals, 70, 524; transmission of plant viruses by, 70, 470, 524; relation of, to virus transmission by Myzus persicae, 689; bionomics and adult habits of, 70, 524; migration of, 70; colour phases of, 70; parasites of, 70, 551; insecticides against, 470.

Macrotermes natalensis, infesting Eucalyptus in Nigeria, 294; insecticides

against, 294.

mactata, Triecphora (see Cercopis sanguinolenta).

maculata, Therioaphis; Vespula.

maculator, Itoplectis.

maculatus, Callosobruchus; Dermestes; Elasmus.

maculicornis, Aphytis; Phaeogenes.

maculipennis, Plutella. maculipes, Pnigalio.

Madagascar, new Mirid on cacao in, 355; breeding of *Locusta migratoria capito* in, 297; list of insects damaging crop plants in, 250; natural enemies of insects in, 653.

madagascariensis, Boxiopsis.

madens, Tribolium.

maderae, Calosoma; Leucophaea (Rhyparobia).

Magdalis armicollis, bionomics of, on elm in Quebec, 455.

Magdalis barbita, bionomics of, on elm in Quebec, 455.

Magdalis frontalis, on pines in U.S.S.R., 116.

Magdalis inconspicua, bionomics of, on elm in Quebec, 455.

Magnesium Carbonate, effect of, on toxicity of insecticide dusts, 118.

Magnesium Fluosilicate, treatment of timber with, against *Hylotrupes bajulus*, **253.**

Magnetic 70, 20.

magnifica, Eacles.

Mahogany (see Swietenia).

maidiradicis, Anuraphis.

maidis, Aphis (Rhopalosiphum).

Maine, Choristoneura fumiferana on balsam fir in, 121, 122; aphids and Phytophthora infestans on potato in, 460, 569, 570; primary food-plants of aphids in, 461; fungi infesting aphids in, 570.

Maize, Cercopids on, in Britain, 321; development of Agrotis ypsilon on, in Bulgaria, 346; pests of, in Canada, 27, 162, 163, 164, 258, 276, 394, 514; Lepidoptera on, in China, 502; pests of, in France, 371, 372, 433, 666; pests of, in Hungary, 306; Calligypona striatella on, in Israel, 349, 350; pests of, in Italy, 349, 350; pests of, in Kenya, 179; Sesamia nonagrioides on, in Morocco, 383; survey of insects on, in New Zealand, 75, 506; Lepidopterous stem borers of, in Nigeria, 292, 293; Cicadulina bipunctella on, in Philippines 96; Ostrinia nubilalis on, in Poland, 365; Busseola fusca on, in S. Rhodesia, 48; O. nubilalis on, in Rumania, 56; virus disease of, in Spain, 350; pests of, in U.S.S.R., 12, 14, 245, 345, 446, 595; pests of, in U.S.A., 21, 65, 95, 152, 215, 225, 227, 275, 276, 277, 281, 325, 327, 328, 329, 333, 396, 398, 420, 461, 508, 509, 511, 580, 639; Blissus leucopterus on, in Venezuela, 98; systematic list of pests of, in South-East Asia and Pacific area, 143; varieties of, in relation to pests, 12, 56, 152, 153, 215, 327, 333, 398, 476, 595; extraction, separation and estimation of factors in varieties of, resistant to O. nubilalis, 22, 68; experiments with strains of O. nubilalis and, in relation to foodplant selection, 14; Nitidulid in relation to O. nubilalis and, 329; relation of O. nubilalis to stalk rot of, 398; physiological effect of Oscinella frit on tissues of, 495; study of, in relation to feeding by O. frit, 595; survival of Chloriona orizicola on, 132; sowing date in relation to infestation of, by S. nonagrioides, 666; virus diseases of, 349, 350; in diets for Diatraea saccharalis, 214, 216; factors affecting yields of, 281, 293; treatment of selfing bags against pests of, 277; destruction of débris of, in control of stem borers, 293; systemic activity of insecticides in, 325, 433; question of endrin residues in, 225; effects of insecticide residues on growth and respiration of, 562.

Maize (Stored), insects in, 51, 205, 206, 269, 403, 554, 560; systematic list of pests of, in South-East Asia and Pacific area, 143; sorption of methyl bromide

by, 554: effects of fumigants on germination of, 331, 332, 579; other measures against pests in, 51, 269, 402, 403, 560, 561.

Maize Meal, in baits for Lepidoptera, 465; in diet for Drosophila, 339; determina-

tion of DDT in, 694.

Maize Oil, in diet for Anthonomus grandis, 404.

Maize Protein Hydrolysate, in baits for Anastrepha ludens, 280.

Maize Steep Liquor, in bait-sprays for Anastrepha ludens, 280.

Maize Syrup, in diet for Drosophila, 339.

major, Penthaleus.

malachitica, Perotis (Aurigena).

Malacosoma americanum (on Prunus spp.), in Canada, 343, 391; in New York, 19; Ephialtes buolianae reared on, 391; bacterial sprays against, 343; systemic insecticides ineffective against, 19.

Malacosoma disstria, in Ontario, 140; key to puparia of Dipterous parasites

of, 140.

Malacosoma neustria (on forest trees), in Holland, 42; in Sardinia, 660; in Spain, 381, 659; in U.S.S.R., 359; characteristics and world distribution of, 507; fluctuations of, 42; fungus infesting, 44; natural enemies of, 379, 381, 660; insecticides against, 659: study of virus disease in relation to DDT-resistance in, 359.

Malacosoma pluviale, infected by new species of Clostridium in British Col-

umbia, 342.

Maladera brunnea, characteristics and world distribution of, 507.

Malaoxon (see O,O-Dimethyl S-1,2-Di-(ethoxycarbonyl)ethyl Phosphorothi-

oate).

Malathion, against Aleurodids, 80, 498; against aphids, 16, 59, 468, 479, 498, 516, 537, 559, 608, 616, 617, 633; ineffective against Myzus persicae, 637; against Coccids, 167, 257, 265, 298, 557, 612, 616; against other Hemiptera, 230, 235, 289, 395, 470, 554; ineffective against Eurygaster, 382; against Curculionids, 16, 51, 270, 317, 331, 402, 403, 500, 516, 560, 571, 685; against other Coleoptera, 51, 179, 211, 225, 265, 270, 471, 497, 500, 560, 561, 571, 600, 671, 685, 687; influence of oxygen uptake on sus-ceptibility of Alphitobius laevigatus to, 201; against Trypetids, 98, 99, 273, 274, 279, 280, 533, 590, 668; against other Diptera, 516, 623, 644, 687; against Noctuids, 161, 211, 222, 516,

559; against Pyralids, 51, 289; unsatisfactory against Zeadiatraea grandiosella, 21; against Tortricids, 20, 26, 248, 466, 538, 555, 637, 671; against other Lepidoptera, 31, 85, 99, 154, 161, 222, 269, 273, 374, 421, 448, 480, 514; against eggs of Operophtera brumata, 669: against Neurotoma nemoralis, 541; toxicity of, to Blattella germanica, 516; against locusts, 172, 383; against Thysanoptera, 193, 212, 569; against Sminthurus viridis, 34; effects of, on populations of Eriophyes vitis, 298; against Tetranychids, 16, 217, 242, 264, 288, 289, 419, 478, 616, 628, 678; not ovicidal to Tetranychus telarius, 278; T. telarius favoured by, 633; cross-resistance to, in strain of T. pacificus, 512; effects of, on beneficial arthropods, 36, 159, 222, 289, 627; repellent action of, 270; vapour toxicity of, 416, 471; factors affecting toxicity and persistence of, 471, 633; in aerosols, 85; in atomised sprays, 85; in baits and bait-sprays, 98, 99, 279, 280, 533, 590; in dusts, 51, 99, 161, 217, 227, 269, 402, 448, 560, 561, 571, 637; in granules, 21; in sprays, 16, 20, 26, 31, 34, 36, 59, 80, 85, 98, 154, 159, 167, 179, 193, 195, 211, 212, 217, 222, 225, 230, 235, 242, 248, 257, 264, 269, 270, 273, 288, 289, 298, 383, 395, 419, 421, 448, 466, 468, 470, 498, 514, 537, 538, 541, 554, 555, 557, 569, 571, 608, 616, 617, 633, 637, 644, 668, 669, 671, 678, 685; applied from aircraft, 211, 280; as deposit, 195, 273; distribution of spray deposits of, 222; trunk applications of, 600; as soil drench, 265; uses of, against pests of stored products, 51, 179, 195, 212, 213, 225, 269, 270, 374, 402, 403, 448, 500, 560, 561, 571, 685; persistence of, in soil, 567; tainting of vegetables and fruit by, 229; residues of, in plants, 270, 571, 644, 671; residues of, in milk and tissues of cattle, 146, 147, 483; metabolism of, 64, 146, 147, 213, 687; labelled with radioactive isotopes, 146, 147; deterioration of packaged preparation of, 227; effects of spray components on toxicity of, 20; and azinphos-methyl, 211; and BHC, 264; and chlorbenside, 242, 678; and DDT, 264, 466; and ethyl-DDD, 161, 222; and fungicides, 26, 36, 264; and methoxy-DDT, 633; and oil emulsions, 167, 557, 616; and parathion, 154, 612; and Tedion, 678; experiments with synergists and, 687: stabilising agents for, 620.

Malaya, Chilo suppressalis on rice in, 78; rubber attacked by termite in, 76, 77; book on Scolytid and Platypodid fauna of forests in, 296; Actia spp. in, 293; Nacoleia octasema in, 353.

malella, Stigmella.

Mali, locusts in, 98, 620.

mali. Aphelinus; Atractotomus; Dasyneura; Eriophyes pyri; Psylla.

Maliarpha separatella, on cereals in Nigeria, 292.

malinellus, Hyponomeuta padellus.

malinus, Pseudaphycus.

malisuctus, Myzus. Malix (see Thiodan).

malleifera, Micrutalis.

Malt, Trogoderma granarium infesting stores of, 295.

Malt Extract, in bait for Drosophila, 327.

Malta, Sitophilus granarius in stored wheat in, 178; use of fossae for storage of cereals in, 177, 178.

Malus (see Pyrus).

Malva parviflora, aphids transmitting virus disease of, in California, 480.

malvae, Macrosiphum (Acyrthosiphon); Podagrica.

Malvastrum tricuspidatum, Earias spp. on, in India, 37.

malvella, Pexicopia (Pectinophora).

Mamestra brassicae, on poppy in Germany, 11; in U.S.S.R., 115, 244; on cabbage, 244; transmission of tobacco mosaic by larvae of, 443; characteristics and world distribution of, 507; parasites of, 115, 244, 245.

Mamestra oleracea (see Diataraxia). mamillata, Agria (Pseudosarcophaga).

mancus, Agriotes.

Maneb, toxicity of, to predator of Leptinotarsa decemlineata, sprays with acaricides, 264, 324.

Manga, characters of pupae of, 292. Manga basilinea, on cereals in Nigeria,

Manganese Sulphate, effects of, fecundity of Aptesis basizona, 155.

mangiferae, Coccus.

Mango, pests of, in India, 170, 315; Coccus mangiferae on, in Israel, 613; Cricula trifenestrata on, in Pakistan, 88; sooty mould on, 613.

Manihot esculenta (utilissima), Amblypelta lutescens papuensis on, in Papua, 235. (See also Cassava Flour.)

manilensis, Locusta migratoria.

Manitoba, Macrosteles fascifrons cereals, carrot and lettuce in, 524; insects on *Melilotus* in, 93, 94, 414, 529; Pristiphora erichsonii on conifers in, 124, 125, 456; Kakimia robinsoni on *Delphinium* in, **529**; virus diseases of plants in, 524; pests of stored products in, 154, 155, 225, 529; beneficial insects in, 125, 414, 529, 551.

Manorina melanophrys, attracted to excretions of Psyllid in New South

Wales, 233.

manto, Platygaster.

Maple, Norway (see Acer platanoides). Maps, of distribution of pests, 1, 357; question of accuracy of, for grasshopper forecasts, 631.

Marasmia trapezalis, DDT against, on

maize in Kenya, 179.

Margaronia indica, in Florida, 507; characteristics and world distribution of, 507.

marginata, Biston; Calligypona (see C. pellucida); Nepiera.

marginatum, Mesogramma.

marginella, Dichomeris. marianae, Tetranychus.

maritimus, Pseudoccocus.

marmorata, Histochaeta.

Marrow-stem Kale (see Kale).

marsupialis, Putionella. martorelli, Macrosiphum.

Maryland, maize pests in, 325; pests of pines in, 20, 336; pests of vegetables in, 222, 325; vectors of virus complex of sweet potato in, 133; Drosophila spp. in tomato fields in, 640; pests of roses in, 212, 264, 324, 477, 633; Amphorophora rubi on wineberry in, 132; Tetranychus telarius overwintering on weeds in, 477; virus diseases of insects in, 336, 404.

mashona, Eulepida.

Masicera zimini, parasitising Dendrolimus

superans in U.S.S.R., 344.

Massachusetts, Sparganothis sulfureana on cranberry in, 337; Ostrinia nubilalis on maize in, 276; Rhyacionia rigidana on pine in, 19; Acarapis sp. infesting honey bees in, 295; parasites and disease of Lymantria dispar in, 399, 400.

Matsucoccus resinosae, on red pine in New York, 19; tests of systemic insecticides against, 19.

Mattesia dispora, infecting Cryptolestes capensis, 658.

maura, Eurygaster.

mauritanicus, Tenebroides.

mauritia, Spodoptera.

Mayetiola, high-altitude catches of, in Britain, 199.

Mayetiola destructor, 199; in Kansas, 65; resistance to, in varieties of wheat, 22, 65, 420.

Meadow Pea (see *Lathyrus pratensis*). Mealworms, larvae of *Tenebrio molitor* (a,v,), 105.

Meat Meal, in baits for *Gryllotalpa*, **442**. Mecarbam, ineffective against *Phylloxera yitifoliae* in galls, **668**.

medicaginis, auct., Aphis (see A. cracci-vora).

medicaginis Koch, Aphis.

medicaginis, Contarinia.

Medicago falcata, Eurytoma roddi infesting seeds of, in Hungary, 306.

Medicago minima, new species of Eurytoma infesting seeds of, in Hungary, 306.

Medicago sativa (see Lucerne).

medicagus, Plagiognathus.

mediterraneus, Pnigalio.

Medlar (Mespilus germanica), Lithocolletis spp. on, in Switzerland, 86; Rhopalosiphum insertum on, 585.

Medlure, attraction of *Ceratitis capitata* to, 383; isomers and analogues of, 383, 384.

megacephala, Pheidole.

Megachile frontalis (in New Guinea), on cacao, 76, 234; on coffee and guava, 234.

Megachile quinquelineata, bionomics of, pollinating lucerne in S. Australia, 234.

Megachile rotundata, pollinating lucerne in U.S.A., 411; artificial nesting sites for, 411.

megalocola, Cataglyphis.

Megaselia, species of, parasitising stages of Caligo eurilochus in Ecuador, 274.

Megaselia pygmaeoides, on mushrooms in Bulgaria, 59.

Megastigmus spermotrophus, on Douglas fir in California, 355.

Megatoma pici, egg-predator of Lymantria dispar in Yugoslavia, 54.

megatoma, Attagenus.

Megatox, containing sodium fluoroaceta-

mide (q.v.), 74.

Megoura viciae, effects of phosphorus compounds on respiration of, 443, 444. Melanagromyza, key to species of, in

Queensland, 295.

Melanagromyza indigoferae, sp.n., on Indigofera suffruticosa in Queensland, 295.

Melanagromyza obtusa, new parasite of, in India, 497.

Melanagromyza phaseoli, 295; on beans in Ceylon, 484; bionomics of, on peas in India, 173; world distribution of, 1, 507; varietal susceptibility of peas to, 173; insecticides against, 173, 484; survey of data on, 507.

Melanagromyza pisi, sp.n., on peas in Queensland, 295.

melanaria, Feronia.

Melanaspis corticosus, new parasite of, in S. Africa, 491.

melanderi, Nomia.

melanocerus, Anthocoris.

melanogaster, Drosophila.

Melanolophia imitata, on Tsuga heterophylla, 395; DDT spray against, 395. melanopa, Lema.

Melanoplus bilituratus, in British Columbia, 155; in U.S.A., 224, 521; bionomics of, 521; adult activity of, 155; infected by Nosema locustae, 582; not reproducing with M. confusus, 694.

Melanoplus bivittatus, bionomics of, in Missouri, 521; infected by Nosema

locustae, 582.

Melanoplus confusus, not reproducing with M. bilituratus, 694.

Melanoplus cuneatus, on rangelands in Arizona, 224.

Melanoplus differentialis, bionomics of, in Missouri, 521.

Melanoplus femurrubrum, bionomics of, in Missouri, 521; survival of, in nitrogen atmospheres, 416.

Melanoplus mexicanus (see M. bilituratus).

melanoscelus, Apanteles.

Melanotus, destroyed by bacterial toxins, 582.

Melanotus caudex, on wheat in China, 625.

melanotus, Coeloides.

Melasoma saliceti, on willows in Poland, 40.

Melcha nursei (see Goryphus).

meles, Hypera.

Meligethes, on cruciferous oil-seed crops in France, 373; natural enemies of, 373.

Meligethes aeneus, in Britain, 318; in Poland, 61; on mustard, 318, 373; on rape, 61, 318, 599; insecticides against, 61, 373, 374; use of, in evaluation of insecticides, 599.

Melilotus, Sitona cylindricollis on, in Canada, 93, 94, 390, 414; Sitona spp. on, in Finland, 102; Therioaphis riehmi on, in U.S.A., 469; tests of resistance to T. riehmi in varieties and species of, 469.

Melilotus alba, Therioaphis riehmi on, in Canada, 529; resistance to T. riehmi in varieties of, 469.

Melilotus dentata, development of Sitona cylindricollis on, 414.

Melilotus officinalis, development of Sitona cylindricollis on, 414; resistance to Therioaphis riehmi in varieties of, 469.

melinus, Strymon.

Melittobia chalybii, parasitising Dioryctria zimmermanni in Illinois, 153.

mella, Tachina (Exorista).

mellonella, Galleria.

Meloboris (see Nythobia).

Melolontha, bionomics of, in Bulgaria, 355; on conifers in Germany, 682, 683; in Poland, 678; measures against, 355, 678, 682, 683.

Melolontha hippocastani, in Czechoslovakia, 186; in Germany, 140; in U.S.S.R., 111, 541, 542; pines injured by, 111; natural enemies and diseases of, 140, 187; other factors limiting larval populations of, 140; effects of insecticides on proportions of strains

of, 541, 542.

Melolontha melolontha (vulgaris), in Bulgaria, 349; in Czechoslovakia, 186; in France, 84; in Germany, 140, 439; in U.S.S.R., 191; on beet, 439; on cabbage, 349; in woods and grassland, 191; characteristics and world distribution of, 506; natural enemies and diseases of, 140, 186, 187; other factors limiting larval populations of, 140, 191; sampling methods for larvae of, 84; effects of electro-magnetic fields on orientation of, 99; measures against, 191, 439.

melolonthae, Neoaplectana.

Melon (Cucumis melo), pests of, in India, 595; Tetranychus desertorum on, in Japan, 452; Dacus cucurbitae on, in W. Pakistan, 175; Leptoglossus australis on, in Papua, 235; Myiopardalis pardalina on, in Turkey, 352; Hemiptera and virus diseases of, in U.S.A., 279, 408; Quadraspidiotus perniciosus reared on, 379; effect of virus diseases on yields of, 408, 409; translocation of insecticides in, 279.

Menadione, in diet for Anthonomus

grandis, 404.

Menazon (PP 175), against aphids, 505, 633, 670; seed treatment with, 670; in sprays, 505, 633.

menciana, Dicranura erminea.

mendicus, Cleonus (Temnorhinus).

Mentha, Noctuids on, in Indiana, 284.

Mentha piperita, Heliothis spp. on, in
Bulgaria, 59; pests of, in Washington,

324. *Mentha spicata*, pests of, in Washington, 324.

Mentha viridis, new pest on, in India, 424. menthae, Phorodon.

menthastri, Sphaerophoria; Spilosoma (see Diacrisia lubricipeda).

mercator, Oryzaephilus.

Mercuric Chloride, soil treatment with, against termite, 77.

Merkaptofos (see Demeton).

merkeli, Neodiprion.

merkeri, Chermes (Dreyfusia).

Mermithids, parasitising insects, 576, 581.

Merodon equestris (on Narcissus), in England, 288; in Holland, 42; measures against, 42, 288.

Meromyza, on cereals and grasses in U.S.S.R., 344, 345; key to Palaearctic species of, 96.

Meromyza nigriventris, on wheat in U.S.S.R., 345.

Meromyza saltatrix, 344; bionomics of, on cereals and grasses in Japan, 96, 97; natural enemies and factors affecting development of, 97; synonymy of, 96.

Meromyza variegata, on grasses in U.S.S.R., 345.

merwei, Aceria (Eriophyes).

Mesites pallidipennis, infesting marine timbers in U.S.S.R., 112.

Mesogramma marginatum, aphids destroyed by, in New York, 401.

Mesoleius aulicus (tenthredinis), parasitising Pristiphora erichsonii in U.S.A., 210.

Mesopolobus, hyperparasite of Evagora milleri in California, 222; effects of sprays on populations of, 222.

Mespilus germanica (see Medlar).

Messor, relation of, to fungus disease of aphids in Congo, 610.

Meta-Isosystox (see Methyl-demeton-S). *Metaphycus* (see *Aphycus*).

Metasyrphus wiedemanni, predacious on

Eriosoma lanigerum in Nova Scotia, 388.

Metasystox (see Methyl-demeton).

Metatetranychus ulmi (see Panonychus). Metcalfa pruinosa, on Citrus in U.S.A., 465; experiments on transmission of tristeza virus by, 465; parasite of, 465.

Meteorus, parasitising Choristoneura in Europe and N. America, 169.

Meteorus bakeri (parasitising Hyphantria cunea), introduced into Yugoslavia, 682; competition between other parasites and, 682.

Meteorus hyphantriae (parasitising Lepidoptera), in Ontario, 162; introduced into Yugoslavia, 682; competition between other parasites and, 682.

Meteorus versicolor, parasitising Stilpnotia salicis, 126; established in British Columbia, 126.

Methionine, action of, on larvae of *Corcyra cephalonica*, 41.

I-Methionine, use of, labelled with 14C in experiments on oviposition by Gryllus assimilis, 686.

Methocel MC-4000 cps., 272.

6-Methoxybenzoxazolinone, separation and activity of, as factor in resistance of maize varieties to Ostrinia nubilalis, 22, 68; inhibition of Penicillium chry-

sogenum by, 22. Methoxy-DDT (Methoxychlor), against Coleoptera, 211, 225, 247, 331, 560, 561, 641; ineffective against Sitophilus oryzae and Tribolium confusum, 270; T. confusum repelled by, 270; against Hemiptera, 230, 395, 633; against Lepidoptera, 26, 222, 398, 411, 503, 644; against Rhagoletis cerasi, 680; against Chaetanaphothrips orchidii, 503; against termites, 577; against Vespula germanica, 422; Tetranychus telarius favoured by, 633; in aerosols, 247, 680; in dusts, 560, 561, 644; in sprays, 26, 211, 222, 225, 230, 270, 395, 398, 411, 503, 633, 641, 680; applied from aircraft, 211, 680; soil treatment with, 577; treatment of nests with, 422; uses of, against pests of stored products, 225, 270, 560, 561; effects of, on growth and respiration of peas, 562; residues of, on plants and plant products, 270, 398, 644, 680; banana fruit damaged by, 503; and malathion, 633.

Methylamine, as metabolite of Sevin,

263.

Methyl Bromide (as fumigant), against Dysmicoccus brevipes, 557; toxicity of, to Coleoptera, 267, 274; against pests of stored products, 296, 331, 332, 499, 554, 661, 685, 694; factors affecting toxicity of, 274, 331, 332; toxicity of, to pineapple, 557; effects of, on seed germination, 579; sorption of, by cereals, 554; determination of, in air, 496.

O-Methyl O-4-tert.-Butyl-2-chlorophenyl Methylphosphoramidate, residues and stability of, in milk, 266.

Methyl Cellulose, as adhesive for seed

treatments, 218, 401.

Methyl Cellulose Polymer, tests and uses of, against Epilachna varivestis and Tetranychus telarius, 272; phytotoxicity of, 272; efficacy of mevinphos increased by, 272; formulation of, 272.

Methyl Chloroform (see 1,1,1-Trichloro-

ethane).

Methyl-demeton, against aphids, 59, 74, 84, 192, 249, 312, 431, 505, 537, 549, 550, 556, 559, 604, 605, 608, 668, 680; against other Hemiptera, 558, 559; against Coleoptera, 139, 176, 180; against Diptera, 139, 549; against Hoplocampa testudinea, 666; against Lepidoptera, 536, 559; against mites, 8, 431, 439, 559, 603, 604; Panonychus ulmi favoured by, 666; toxicity of, to honey bees, 550; effects of, on beneficial arthropods, 312, 313, 439, 605; in sprays, 59, 74, 84, 176, 192, 249, 312, 313, 431, 505, 536, 537, 549, 550, 556, 558, 559, 604, 605, 608, 666, 680; evaluation of spray programmes of, 376; in drenches, 558; soil treatment with, 180; tests of systemic activity and other methods of application of, 176, 249, 431, 559; and ethion, 604, 605; and lead arsenate, 559; and trichlorphon. 549.

Methyl-demeton-S (in sprays), against aphids, 505, 617; against Paratetranychus ilicis, 19; tests of systemic activity

of, 19.

O-2,4-Dichlorophenyl Iso-O-Methyl propylphosphoramidothioate, effects on Macrosiphum pisum of broad beans treated with, 525.

O-Methyl O-2,4-Dichlorophenyl Methylphosphonothioate, against aphids, 216, 402; ineffective against Lepidoptera, 413; in dusts, 216; in sprays, 402, 413.

Methyl 1,1-Ethoxycarbonylmethyl-2,2-dichloro-1,2-propylene Phosphate, in sprays against Dacus oleae, 309.

Methyl 1,1-Ethoxycarbonylmethyl-2,2dichloro-1,3-propylene Phosphate. against Aphis pomi, 616; against Dacus oleae, 135, 438; against Coccids, 616; against Panonychus ulmi, 616; in sprays, 135, 438, 616; and oil emulsion, 616.

Methyl 3-(Ethoxy(p-dimethylaminophenyl) phosphinyloxy) crotonate, against Macrosiphum pisum, 64; against Musca domestica, 63, 64; against Tetranychus telarius, 64; with synergists, 63, 64.

Methyl Ethyl Ketone, as carrier for solu-

tions of pyrethrins, 3.

2-(Methyl-n-heptyl)-4,6-dinitrophenylcrotonate, effects on beneficial Mirids of sprays containing, 672.

β-Methylindole, in baits for *Drosophila*,

Methyl p-Nitrophenyl Ethylphosphonate. anticholinesterase activity and toxicity of, to Musca domestica, 407.

Methyl-paraoxon (see Dimethyl p-Nitro-

phenyl Phosphate).

Methyl-parathion, against aphids, 16, 59, 84, 312, 537, 559, 668; against other Hemiptera, 115, 382, 560; against

Curculionids, 16, 317, 322, 330, 515, 541; resistance to, in strains of Anthonomus grandis, 147; against other Coleoptera, 58, 61, 647; against Diptera, 45, 63, 64; against Lepidoptera, 9, 413, 515, 538, 558, 559; Heliothis zea favoured by, 323; against Frankliniella occidentalis, 409; toxicity of, to Locusta migratoria, 172; against mites, 16, 191, 515, 548; Tetranychids favoured by, 256; influence of foodplant on susceptibility of Tetranychus telarius to, 548; effects of, on beneficial insects, 312, 560; question of influence of, on susceptibility of insects to diseases, 115, 538; in dusts, 9, 45, 58, 59, 61, 115, 256, 537, 541; in sprays, 9, 16, 84, 115, 191, 213, 312, 322, 409, 413, 515, 537, 538, 558, 559, 560, 632; treatment of timber with, 647; tests of, against pests of stored products, 212, 213; effects of, on growth of cotton, 632; and Beauveria bassiana, 541; and DDT, 322, 515; and parathion, 191, 647; and Shell SD-4402, 413; and toxaphene, 409; antagonistic action of synergists with, 63, 64.

dl-α-Methylphenethylamine (as sulphate), effects of, in mixture with rotenone against *Epilachna varivestis*, 279.

Methylphosphoric Acid, as metabolite of dimethoate, 563.

3-Methyl-5-pyrazolyl Dimethylcarbamate, treatment of rubber strips with, against *Piophila casei*, 100; colorimetric determination of, 227.

Methylthiomethylpyrimidinyl Dimethylcarbamate, in sprays against Dacus

oleae, 438.

4-Methylthio-3,5-xylyl Methylcarbamate, toxicity of, to *Aphis spiraecola*, **516**; against Lepidoptera, **509**; toxicity of, to *Musca domestica* and honey bees, **516**; in dusts, **509**; in sprays, **509**; and piperonyl butoxide, **516**.

O-Methyl O-2,4,5-Trichlorophenyl N-Ethyl Phosphoramidothioate, toxiciyt of, to *Rhagoletis pomonella*, **268**.

O-Methyl O-2,4,5-Trichlorophenyl Phosphoramidothioate, toxicity of, to *Rhagoletis pomonella*, 268; against *Tetranychus telarius*, 217; in dusts, 217; in sprays, 217.

Methyl Trithion (see O,O-Dimethyl S-p-Chlorophenylthiomethyl Phosphoro-

dithioate).

Metopolophium dirhodum (see Macrosiphum).

Metrazol (see 6,7,8,9-Tetrahydro-5-H Tetrazoleazepine).

Mevinphos, against aphids, 16, 19, 312, 324, 412, 468, 498, 637, 667, 668; against other Hemiptera, 395, 434, 667; against Coleoptera, 16, 271, 497; against Fenusa pusilla, 19; against Trypetids, 268, 668, 669; against Lepidoptera, 9, 19, 161, 256, 267, 434, 480, 667, 668; against mites, 16, 19, 324, 628; toxicity of, to beneficial arthropods, 312, 313, 434, 627; toxicity of, to bees, 412; anticholinesterase activity and mammalian toxicity of, 384, 385; in dusts, 161, 637; in sprays, 9, 16, 213, 256, 267, 312, 313, 324, 395, 434, 468, 667, 668, 669; tests of, against pests of stored products, 212, 213; systemic activity of, 19, 434; metabolism of, in pea, 88; residues of, on oranges, 267; bioassay of residues of, 397; adjuvants increasing residual activity of, 271; and oil emulsion, 256, 667.

mexicanus, Melanoplus (see M. bilituratus). Mexico, Copturus aguacatae on avocado in, 356; Citrus pests in, 22, 279, 280, 325; pests of cotton in, 407, 473, 511; Pectinophora gossypiella on other species of Gossypium and other plants in, 511; Chloriona spp. and hoja blanca virus of grasses and rice in, 518; Dasiops alveofrons in, 128; beneficial insects in, 22, 518, 529.

meyricki, Leucoptera. micado, Scapsipedus. micans, Rhadinoceraea.

Mice, wild bee destroyed by, 565; Bacillus thuringiensis not pathogenic to, 282; survival of B. thuringiensis in faeces of, 282.

Michigan, pests and mosaic virus of beans in, 16, 17; forest pests in, 210, 327, 410, 413, 481, 495; beneficial insects in, 210, 327; fungi infesting *Pristiphora erichsonii* in, 210.

Microbracon (see Bracon).

Micro-capillary Tubes, topical application of insecticides with, 3.

Microcells, for rearing mites, 518.

Microcryptus galactina (see Aptesis).

Microctonus aethiops (parasitic on Curculionids), introduced into Canada, 390, 414; in France, 390, 414; in Sweden, 390; bionomics of, 390.

Microdiprion pallipes (on pines), bionomics of, in Sweden, 426; in U.S.S.R., 116.

Microgaster polita, parasitising Argyres-

thia conjugella in Germany, 10.

Micronematus monogyniae (see Pristi-

phora).

Micronesia, parts of a work on insects of, 496.

Microphanurus (see Asolcus).

Microphanurus anitus (see Asolcus rufiventris).

Microphanurus choaspes (see Asolcus

vassilievi).

Microplitis alaskensis, bionomics of. parasitising *Pseudaletia unipuncta* in Ontario, 129.

Microplitis plutellae, Plutella maculipennis attacked by, in New York, 401.

Microtermes obesi, attacking sugar-cane in India, 119, 553; measures against,

119, 553.

Microterys flavus (parasitic on Coccids), in California, 150, 597; in Pakistan, 150; introduced into U.S.S.R., 597; host selection by, 150.

Microterys frontatus, parasitising Coccus mangiferae in Israel, 613; in U.S.S.R.,

597.

Microtrombidium demeijerei, parasitic on

Oscinella frit in Germany, 45.

Micrutalis malleifera (in Florida), bionomics of, on egg-plant, 636; transmitting curly-top virus of tomato, tobacco and other plants, 635, 636; infested by Beauveria, 636.

middendorffi, Dendrosoter.

Middle East, Pentatomids on cereals in, 484, 485, 654, 655, 656; natural enemies of Pentatomids in, 484, 485, 654, 655. (See also individual countries.)

Midol 556, 674.

migrator, Tetrastichus. migratoria, Locusta.

migratorioides, Locusta migratoria.

Milk, insecticide residues in, 92, 225, 266, 384, 483; recovery of insecticide residues from, 266.

Milk Powder, in baits for Gryllotalpa, 442.

milleri, Evagora (Recurvaria).

Millet (see Panicum, Pennisetum and Sorghum).

mindara, Proxenus.

minidoka, Eurygaster.

minima, Carulaspis.

minimus, Pachygonatopus.

minki, Anthocoris.

Minnesota, cabbage pests in, 172; Empoasca fabae on potato in, 627; Ostrinia nubilalis on maize in, 276, 398, 461; forest pests in, 149, 212, 411, 419, 467, 474, 552, 619; parasites of Choristoneura pinus in, 212.

(Pentatrichopus); minor, Capitophorus Formica rufa rufopratensis (see F. polyctena); Sitophilus oryzae (see F.

oryzae).

minor pratensoides, Formica (see nigricans).

Mint (see Mentha).

minuta, Hoplocampa. minutum, Trichogramma.

miranda, Abraxas.

Miscanthus saccharifer (sacchariflorus). Aphis sacchari overwintering on, in China, 624; destruction of, for control of A. sacchari, 624.

Mississippi, fire ants in, 464, 639; cotton pests in, 410, 473, 515, 573, 574; Rhyacionia rigidana on pine in, 20; Acarapis infesting honey bees in, 295.

Missouri, forest pests in, 19, 88; Ostrinia nubilalis on maize in, 276; grasshoppers in, 520, 521, 576; nematode infesting Chortophaga viridifasciata in, 576.

Mitaf, in sprays, against Eotetranychus carpini, 183; constituents of, 183.

Mite, European Red (see Panonychus

ulmi). **Tetranychus** Mite, (see

telarius).

Mites, in S. Africa, 83; part of work of British species of, 41; on vines in France, 445; book including section on, in Israel, 37; on olives, vines and fruit trees in Italy, 40; on wheat in Jordan, 684; on fruit trees in Switzerland, 682; surveys of data on, not occurring in U.S.A., 506-508; morphological study of injury to bean leaves by, 107, 108; influence of, on growth of plum trees, 671; comparison of damage to vines by virus and, 663; and virus diseases of plants, 70, 71, 302, 303, 523, 684; relation of, to leprosis of Citrus, 556; and silver top of grasses, 156; associated with stored products, 41, 149, 154, 155, 364, 424, 518, 529; infesting houses, 419, 519; infesting honey bees, 295, 296, 342, 504, 505, 617; flight of Dendroctonus pseudotsugae not influenced by, 127; habits and ecology of, 41, 90, 477, 518, 528; regulation of water loss by, 564; factors affecting development of, 256, 259, 444, 583; factors affecting oviposition and diapause in, 361, 516, 547; natural enemies of, 122, 125, 183, 259, 302, 319, 378, 379, 439, 454, 522, 528, 549, 627, 677, 682; virus disease of 581; predacious species of, 30, 34, 45, 122, 125, 149, 159, 183, 259, 268, 302, 319, 364, 376, 378, 379, 386, 387, 414, 439, 454, 459, 500, 522, 523, 528, 549, 640, 641, 677, 682; infesting cultures of Conoderus falli, 575; effects of insecticides on populations of, 36, 111, 298, 503, 523; resistance to acaricides in strains of, 278, 284, 602, 688; relation of resistance to TEPP to diapause in, 428; influence of food-plant on

resistance to acaricides in, 548; highaltitude collection of, 578; microcells for study of, 518; method for transport of, 640; method of rearing, 641; colour forms of, 548; classification and new species of, 41, 452, 521.

Mitin FF, treatment of textiles with, against Hofmannophila pseudospretella,

290.

mitis, Zenillia (Platymyia).

mitterbacheriana, Ancylis.

Mocis repanda, toxicity of insecticides to.

Molasses, in diet for Aphelinus mali, 591; as adhesive with Bacillus thuringiensis, 107.

molesta, Cydia (Grapholitha, Laspeyresia). molitor, Tenebrio.

mombinpraeoptans, Anastrepha.

monacha, Corythaica; Lymantria (Ocneria).

monachus, Apate.

Monarthropalpus buxi, bionomics of, on box in Italy, 694; measures against, 694.

Monellia costalis, on pecan in California, 151: Trioxys pallidus reared from, 151.

mongeneti, Pimelia.

Mongolian Oak (see Quercus mongolica). Mongolian Republic, Dendrolimus superans and secondary pests on conifers in, 116; parasites of D. superans in, 116.

Monochaeta albicans, parasitising Operophtera brumata in Europe, 257, 378; introduced into Canada, 378; factors affecting survival of, 257.

Monochamus, species of, on timber in

Finland, 42.

Monochamus oregonensis, damaging firekilled spruce in British Columbia, 457. Monochamus sutor, bionomics of, on timber in Finland, 42; sprays against,

Monoctenus juniperi, damaging juniper in Holland, 302, 303; parasites of, 302.

Monoctonus cerasi, parasitising Anuraphis plantaginea in Holland, 673.

Monoctonus paulensis, parasitising Macrosiphum pisum in California, 66.

Monodontomerus aereus, survey of data on, parasitising Choristoneura murinana in France and Germany, 168.

Monoglyceride, in baits for fire ants, 464.

monographus, Xyleborus.

monogyniae, Pristiphora (Micronematus). Monosteira unicostata, on almond in Italy, 38; characteristics and world distribution of, 507; insecticides against,

Monsanto 16226 (see Octachlorodipropyl Ether).

Monsanto CP-7769 (see Hexaethyl Ethylthiomethylidynetriphosphonate).

Monsanto CP-10502 (see Dimethyl 1-(Dimethoxyphosphinyl)vinyl Phosphate).

monspeliensis, Gryon (Hadronotus),

Montan Wax, in stable formulations of dichlorvos, 562.

Montana, Tortricids on Douglas fir in, 24, 460, 461; Chorizagrotis auxiliaris on mustard in, 640; grasshoppers on rangelands in, 223; parasite of Choristoneura fumiferana in, 24.

montana, Omphale.

monticolae, Dendroctonus (see D. ponderosae).

Moraine Locust (see Gleditsia).

Mordwilkoja vagabunda, bionomics of, on Populus deltoides in Minnesota, 552, 619.

moreonella, Evagora (Recurvaria).

mori, Bombyx.

morigerus, Xyleborus.

morio, Eurytoma.

Morocco, pests and diseases of avocado in, 383; Coccids on Citrus in, 382; pests of date in, 54, 382; Pentatomids on cereals and grasses in, 382, 653, 654, 655; Sesamia nonagrioides on maize in, 383; pests and diseases of cotton in, 490; Aelia spp. in, 355; Bruchidius algiricus in, 249; Schistocerca gregaria in, 383; natural enemies of pests in, 53, 54, 382, 490, 653, 654, 655; diseases of Coccids in, 53, 54.

morosa, Lachnosterna (Holotrichia). Morphothion, against Dacus oleae, 237, 238; effects of, on parasite of Eriosoma lanigerum, 312; in dusts, 237; in sprays, 237, 238, 312; residues of, in olives, 238; decomposition of residues of, 238.

morstatti, Xyleborus.

Morus, pests of, in India, 626; Pseudaulacaspis pentagona on, in Ontario, 529. (See also Mulberry.)

mosellana, Sitodiplosis.

Mosquitos, review of behaviour of, in relation to control, 90; metabolism of DDT in, 64.

Mould, relation of, to development of Ahasverus advena, 657. (See also Sooty Mould.)

Mountain Ash (see Sorbus aucuparia).

mucronifera, Ceracia.

Mulberry, Hyphantria cunea on, in Austria, 312; Pseudaulacaspis pentagona on, in Italy, 437. (See also Morus.)

multisetis, Tetranychus.

multistriatus, Scolytus.

munda, Agrotis (see Euxoa radians). Mung Bean (see Phaseolus aureus). murinana, Choristoneura.

murinanae, Apanteles; Cephaloglypta.

Musca domestica, experiments on parasitisation of, by Nasonia vitripennis, 128: survival of, in nitrogen atmospheres, 416; toxicity to, of extract and fractions from Bacillus larvae, 332, 333; used in tests of insecticides, 2, 3, 63, 202, 221, 263, 407, 416, 512, 516, 562, 573, 603, 634, 635, 687, 688; use of, in bioassay of insecticide residues, 397; toxicity of haemolymph from DDTtreated cockroach to, 63; resistance to insecticides in strains of, 63, 148, 263, 573; metabolism of insecticides in, 64, 148, 263, 476; effects of insecticides on enzymes of, 407, 634, 635; enzyme activity of homogenates of, on phosphorus compounds, 213.

Musca domestica nebulo, used in tests of

insecticides, 171.

Muscatox (see Coumaphos).

musculus, Anthocoris.

Musgraveia sulciventris, characteristics and world distribution of, 507.

Mushroom, Japanese Pine (see Tricholoma matsutake).

Mushrooms, Sciara infesting, in Brazil, 644; Megaselia pygmaeoides on, in Bulgaria, 59; pests and diseases of, in Germany, 425; Tyrophagus putrescentiae on, in Ohio, 573; insecticide residues in, 644.

Mustard, Chorizagrotis auxiliaris on, in Alberta and Montana, 640. (See also Brassica juncea, B. nigra and

Sinapis alba.)

Mustard, Black (see Brassica nigra).

Mustard Oil (see Allyl Isothiocyanate). Myelois ceratoniae (see Ectomyelois).

myersi, Carabunia.

Myiopardalis pardalina, bionomics and natural enemies of, on cucurbits in Turkey, 351, 352; characteristics and world distribution of, 507.

Myllocerus dorsatus, on grapefruit in India, 173.

Myllocerus evasus, on lemon in India, 173; characters of, 173.

Mymar behmani, parasitising leafhoppers in California, 630.

Myochrous armatus, on jute in British Guiana, 27; BHC against, 27.

myopaeformis, Tetanops.

Myriangium thwaitesii, Eriococcus orariensis infested by, in New Zealand, 587,

Myrmarachne, destroying Oecophylla sp. in Zanzibar, 29.

Myrmecomimesis, parasitising Phasmatids in Australia, 32, 231; effect of bush fires on larvae of, 231.

Myrothecium roridum, toxicity of culture filtrates of, to Epilachna varivestis,

Mystox B, containing pentachlorophenol (q.v.), 290.

Mystox LPL, containing pentachlorophenyl laurate (q.v.), 290.

mytilaspidis, Aphytis.

Mytilococcus beckii (see Lepidosaphes).

Myxexoristops hertingi, parasitic on Acantholyda erythrocephala in Germany, 10. Myzocallidium riehmi (see Therioaphis). Myzocallis alhambra, honeydew of, in

diets for Ephialtes buolianae, 391.

Myzodes persicae (see Myzus).

Myzotoxoptera staphyleae tulipaellus (see

Rhopalosiphoninus).

Myzus ascalonicus, in British Columbia, 692; on beet in France, 605; on strawberry and stored onions in Germany, 106, 492; on Easter lily in Oregon, 278; plant viruses in relation to, 107, 605, 611, 692; feeding habits of, 493; mechanism of saliva secretion by, 45; effects on plant cells of feeding by 92; overwintering of, 605; respiration of, 443; labelled with 32P, 45; systemic use of insecticides against, 278.

Myzus brevipilosus (see Capitophorus minor).

Myzus cerasi, insecticides against, on cherry in Poland, 188.

Myzus circumflexus, relation of, to virus transmission by M. persicae, 689.

Myzus malisuctus, on apple in Japan, 36; effects of insecticide sprays on populations of, 36.

Myzus ornatus, in British Columbia, 692; on vegetables in California, 131, 480; transmission of plant viruses by, 17, 131, 480, 692; relation of, to virus trans-

mission by M. persicae, 689.

Myzus persicae, in Australia, 34, 35; in Belgium, 249; in Brazil, 556; in Britain, 199, 200, 320, 431; in Canada, 633, 692; in Ceylon, 81; in France, 605, 669, 670; in Gaza Strip, 97; in Germany 303, 546, 547, 680; in Israel, 130, 614; in Poland, 544; in U.S.A., 68, 131, 133, 216, 264, 278, 331, 333, 339, 401, 406, 408, 412, 461, 463, 474, 480, 508, 566, 569, 570, 573, 630, 636, 637; on broad beans, 34, 35; and virus diseases of beans, 34, 35, 264; on beet, 331, 463, 545, 546, 605, 669, 670; transmitting virus yellows of beet, 17, 131, 249, 331, 463, 482, 545, 605, 669; on

Capsicum, 216, 630, 636, 637; transmitting virus diseases of Capsicum, 636; on crucifers, 401, 508, 691; relation of populations of, to planting patterns in crucifers, 68, 69; transmitting virus diseases of crucifers, 131, 197; on lettuce, 569; transmitting virus diseases of lettuce, 131, 467; on potato, 320, 333, 412, 461, 474, 544, 546, 547, 556, 570, 614, 633, 680; transmitting virus diseases of potato, 320, 544, 547, 641, 680, 688, 689, 690, 691; transmitting virus disease of spinach, 480; transmitting virus disease of sweet potato, 130, 131, 133; on Citrus, 97; transmitting virus disease of melons, 408; on peach, 547; transmission of strawberry viruses by, 692; transmitting vine viruses, 303, 304; experiments with virus of groundnuts and, 611; on tobacco, 566; feeding habits of, in relation to nicotine content of tobacco, 566; susceptibility tobacco varieties and Nicotiana spp. to, 406; transmission of tobacco viruses by, 81, 303, 690; on carnation, 431; on Easter lily, 278; on greenhouse plants, 339; preference of, for excised or virus-infected leaves of Physalis floridana, 584; other food-plants of, 461, 480, 573, 689, 691; rôle of wavelength in food-plant selection by, 200, 201; stomach size of, in relation to food-plants, 423; feeding habits of, 482; electron-microscopic study of salivary glands of, 300; experiments on transmission of non-persistent viruses by, 250; respiration of, 443; influence of electro-magnetic fields on moulting of, 444; flight activity of, 547, 605; forecasting outbreaks of, 460; natural enemies of, 401; fungi infesting, 401, 570; traps for, 547, 614; fluctuations in trap catches of, 320; high-altitude catches of, 199; sampling methods for, 463; insecticides against, 216, 249, 278, 320, 331, 333, 412, 431, 474, 508, 556, 569, 573, 605, 631, 633, 637, 670, 680; favoured by DDT, 631; resistance to parathion in strains of, 412.

Myzus poae, on grass in Wisconsin, 160.

Myzus ribisnigri, on Ribes alpinum in Ontario, 529.

Myzus solani (see Macrosiphum).

N

Nabam, toxicity of, to predator of Leptinotarsa decemlineata, 378.

Nabis, predacious on cotton pests in Arizona, 639.

Nacoleia octasema (on banana), in Fiji, 353; in Indonesia, Malaya, New Britain and New Guinea, 353, 354; in Queensland, 503; in Singapore, 354; other food-plants of, 354; parasites of, 354; insecticides against, 503.

nanella, Recurvaria.

Nanus ipomoeae, 133.

1-Naphthol, as metabolite of Sevin, 263.1-Naphthyl Dimethylcarbamate, hydrolysis and anticholinesterase activity of, 635.

1-Naphthyl Methylcarbamate (see Sevin). Napomyza lateralis (see Phytomyza).

nararia, Natada (Macroplectra).

Narcissus, Merodon equestris on bulbs of: in England, 288; in Holland, 42; treatments of, against M. equestris, 42, 288.

Nasonia vitripennis, experiments on parasitism of Musca domestica by, 128.

Nasonovia ribisnigri (see Myzus).

nasturtii, Aphis; Contarinia. Nasturtium (see Tropaeolum).

Nasutitermes, in India, 627; treatments

of nests of, with insecticides, 627.

Nasutitermes columbicus, toxicity of treated concrete blocks to, 416.

Nasutitermes corniger, infesting building timber in British Honduras, 76.

Nasutitermes ephratae, resistance of timbers to infestation by, 649.

Nasutitermes exitiosus, in building timbers in New South Wales, 33; Reduviid predacious on, 33.

Nasutitermes nigriceps, infesting building timber in British Honduras, 76.

Natada nararia, on tea in Céylon, 448; granulosis virus of, 448; measures against, 448, 449.

natalensis, Macrotermes.

Neave, Dr. S. A., death of, 1.

Nebraska, Loxostege spp. on beet in, 335; pests of cereals in, 214; Therioaphis spp. on leguminous forage crops in, 469, 631.

nebulo, Musca domestica.

nebulosus, Anthribus.

Necrobia rufipes, intercepted on oil-seed cake in Germany, 11; infesting copra imported into Italy from India, 436; bionomics and food preferences of, 436; insecticides against, 436.

Nematocerus, species of, on cereals in Kenya, 180; parasite of, 180; tests of insecticides against eggs and larvae of, 180.

Nematocerus brevicornis, 180.

Nematocide 18133 (see O,O-Diethyl O-2-Pyrazinyl Phosphorothioate).

Nematodes, transmitting virus disease of vines in Germany, 71; weevil as vector of pathogenic, 575; infesting insects, 127, 162, 163, 187, 207, 208, 210, 496, 576, 580, 581, 619; bacteria associated with, 162, 207.

Nematus ribesii, pathogenicity of Bacillus

cereus to, 282.

Nemeritis canescens, study on multiparasitism of Anagasta kuehniella by Nythobia chrysostictos and, 618, 619.

nemoralis, Anthocoris; Neurotoma. Nemorilla floralis, parasitising Ostrinia

nubilalis in U.S.S.R., 245.

nemorum, Anthocoris; Phyllotreta.

nenuphar, Conotrachelus.

Neoaplectana melolonthae, infesting Melolontha melolontha in Czechoslovakia, 187.

Neocnemodon elongatus (see Cnemodon).

neocynodonis, Aceria.

Neocypholaelaps, species of, infesting honey bees in Queensland, 296.

Neoderostenus (see Achrysocharoides). Neodiprion, on pines in California, 23, 24, 125; bionomics and parasites of, 23, 24;

sampling method for, 125.

Neodiprion abietis, pathogenicity of Bacillus cereus to, 282.

Neodiprion lecontei, in Canada, 341; effects and transmission of virus diseases of, 341, 342; pathogenicity of Bacillus spp. to, 283.

Neodiprion merkeli, sp.n., on Pinus

elliottii densa in Florida, 462.

Neodiprion pratti banksianae, in Canada, 342; virus disease of, 342; pathogenicity of Bacillus spp. to, 283.

Neodiprion pratti pratti, on pines in U.S.A., 336; polyhedrosis virus of, 336; aerial application of virus against, 336.

Neodiprion sertifer (on pines), in Canada, 163, 209, 341, 387; in Sweden, 426; parasite of, 209, 387; effects and transmission of virus diseases of, 341, 342.

Neodiprion swainei (on pines in Canada), 163; extraction of cocoons of, from soil samples, 258, 259; natural enemies of, 73, 459; pathogenicity of Bacillus spp. to, 283; trans-ovarial transmission of polyhedrosis virus of, 580.

Neodiprion warreni, sp.n., on Pinus echinata in Arkansas, 462.

Neoleucopis obscura (see Leucopis).

Neophilaenus, on grasses and cereals in Britain, 321; experiments on decomposition of froth masses of, 321; insecticides against, 321.

Neosciara amoena (see Sciara).

Neotran, development of resistance to, in strains of *Panonychus citri*, 510.

Nephantis serinopa, on coconut in Ceylon, Cochin and India, 169; parasites of, 169.

nephopyropa, Lechriolepis.

Nephotettix cincticeps, on rice in China, 625; seed treatments against, 625.

Nepiera marginata, bionomics of, parasitising Pseudaletia unipuncta in Ontario, 129.

nervosa, Depressaria (see D. apiella).

Netelia latungula (see Parabates).

Netelia sayi, bionomics of, parasitising Pseudaletia unipuncta in Ontario, 129. Neurotoma nemoralis, on cherry, peach and other stone-fruits in Bulgaria, 540, 541; bionomics and parasite of, 540;

insecticides against, 540, 541.

neustria, Malacosoma.

Nevada, *Dendroctonus* spp. on pines in, 340.

New Britain, coconut pests in, 166, 167, 293; Leptoglossus australis in, 235; Nacoleia octasema in, 353; natural enemies of coconut pests in, 166, 293.

New Brunswick, Rhagoletis pomonella on apple in, 273; Thysanoptera on blueberry in, 71; aphids on cereals and grasses in, 584–586; aphids on potato in, 633; forest pests in, 25, 121, 122, 124, 394, 455, 529; winter food-plants of aphids in, 585; Panonychus ulmi in, 528; Scymnus impexus introduced into, 394.

New Caledonia, pests of coconut in, 81. New Guinea, pests of cacao in, 76, 196, 234; coffee pests in, 76, 234; Megachile frontalis on guava in, 234; new Coccid on yams in, 234; Nacoleia octasema on Heliconia in, 353, 354; Xyleborus morigerus introduced into Britain in orchids from, 625; new parasite reared from Agonoxena pyrogramma in, 52. (See also New Britain, New Ireland, and Papua and New Guinea, Territory of.)

New Ireland, Leptoglossus australis in, 235.

New Jersey, Sparganothis sulfureana on cranberry in, 337; Hypera variabilis on lucerne in, 400; Bryobia praetiosa infesting houses in, 519; parasite of H. variabilis introduced into, 400.

New Mexico, cotton pests in, 409; pests of pines in, 458, 460; natural enemies of

Dendroctonus barberi in, 460.

New South Wales, Anodontonyx noxius attacking cereals in, 493; Ceroplastes destructor on Citrus in, 35; forest pests

in, 32, 33, 231, 232, 233; Nasutitermes exitiosus damaging building timber in, 33; host-plant range of wilt virus in, 34; beneficial insects in, 32, 33, 231, 233; bird attracted to Psyllid excre-

tions in, 233.

New York, apple pests in, 18, 23, 268; Hendecaneura shawiana on blueberry in, 567; pests of peach in, 23, 326, 631; pests of other fruit trees in, 23, 216, 470; virus disease of stone-fruits in, 23; Psila rosae on carrot in, 226; Trichoplusia ni on cauliflower in, 328; pests of cole in, 401; ecological study of invertebrate fauna of crucifers in, 68; Macrosteles fascifrons as vector of yellows virus of lettuce in, 470; Hylemya antiqua on onion in, 226; forest pests in, 18, 19, 220, 276, 337; pests of leguminous forage plants in, 213, 280, 323, 576, 581; pests of ornamental plants in, 19; other insects in, 164, 399; natural enemies of pests in, 337, 401, 402, 581; diseases of insects in, 328, 401; DDT and copper residues in vineyard soils in, 228.

New Zealand, Quadraspidiotus perniciosus on apple in, 167; Vespula germanica damaging grapes and other fruits in, 422; *Plusia chalcites* on beans in, 75; pests of crucifers in, 74, 75, 167, 505, 506; cereal pests in, 75, 506, 684; pests of grasses in, 231, 505, 506, 684; virus diseases of cereals and grasses in, 684; survey of insects associated with lucerne and other fodder crops in, **75, 506;** Agriophara coricopa on Olearia fosteri in, 75; natural enemies of honey bees in, 295, 422; history of establishment of *V. germanica* in, 421-423; other insects in, 232, 424; beneficial insects in, 75; introduced and indigenous Coccids on Leptospermum in, 587, 588; fungus infesting Eriococcus

orariensis in, 587, 588.

Newfoundland, Chermes piceae balsam fir in, 456; predators of C. piceae introduced into, 394, 456.

Nexion (see Methyl 1,1-Ethoxycarbonylmethyl-2,2-dichloro-1,3-propylene Phosphate).

Nezara viridula smaragdula, parasitised by Asolcus basalis in West Indies, 589. ni, Trichoplusia (Phytometra, Plusia).

nicolasi, Campylomma.

Nicotiana, experiments on susceptibility or resistance of species and hybrids of, to pests of tobacco, 406.

Nicotiana debneyi, grasshoppers attracted

to, 406.

Nicotiana exigua, gossei, nudicaulis, repanda, tabacum (see also Tobacco), tomentosiformis, trigonophylla, velutina, 406.

Nicotine, against aphids, 148, 217; in relation to feeding by Myzus persicae, 566; rôle of, in resistance of solanaceous plants to Coleoptera, effects of, on symbiont of Lasioderma serricorne, 650; against Lyonetia clerkella, 663; susceptibility of sexes of Periplaneta americana to, 117; occurrence of, in tobacco, 566; in dusts, 217; in sprays, 217, 663.

Nicotine Sulphate (in sprays), against aphids, 188; against *Psylla mali*, 159; toxicity of, to *Rhagoletis pomonella*, 274; effects of, on beneficial arthropods, 35, 36; and calcium arsenate,

274; and lime-sulphur, 35.

niger, Coptotermes.

Nigeria, pests of cacao in, 293, 652; diseases of cacao in, 652; Lepidopterous stem borers of cereals in, 292, 293; first record of Contarinia sorghicola on sorghum in, 684, 685; Coccus hesperidum on Citrus in, 531; Oecophylla longinoda associated with C. hesperidum in, 531; Coleoptera infesting field and stored cowpeas in, 49, 50; Diparopsis watersi on cotton in, 53; termites attacking seedlings Eucalyptus in, 293, 294; termites infesting timber and furniture in, 485, 650; beneficial insects in, 28, 53, 292; fungus infesting Busseola fusca in, 292.

nigericus, Gastrimargus.

nigerrimus, Ooencyrtus.

nigra, Lissonota.

nigricana, Cydia; Eucosma (Epinotia).

nigricans, Formica.

nigriceps, Nasutitermes.

nigricornis, Trioza. nigrifemur, Perisierola.

nigripes, Blondelia.

nigrirostris, Hypera.

nigrisigna, Autographa (Plusia).

nigriventris, Meromyza.

nigrofasciatus, Dysdercus; Phonoctonus.

nigrorepletus, Hieroglyphus.

nilgiriensis, Thrips.

Ninhydrin, use of, in detection of infestation of stored cereals, 571.

Nipa fruticans, Nacoleia octasema on, in Indonesia, 354.

niponensis, Carposina (see C. sasakii).

Niptus hololeucus, damaging textiles in Germany, 377.

nitidalis, Diaphania. nitidulator, Opius.

Nitric Acid, experiments with, in decomposition of froth masses of Cercopids, 321.

Nitrogen, influence of, on infestation of pines by insects, 598; other effects of, on insects, 324, 416; question of use of, as anaesthetic, 416.

p-Nitrophenyl Ethyl-tert.-butylphosphinate, anticholinesterase activity and toxicity of, to *Musca domestica*, **407**.

Nitrosan, effects of sprays containing, on Aphelinus mali, 678.

nivaliensis, Dioryctria.

nivella, Scirpophaga. nobilellus, Horcias.

Noctuelia rufofascialis, reared from capsules of Cienfuegosia sulphurea in Texas, 518.

noctuella, Nomophila.

Nomadacris septemfasciata, in Tanganyika, 98; factors affecting populations of, 98, 285, 286; estimation of mortality in successive instars of, 426; sensory equipment of, 423; history of preventive control of, 87.

Nomadopsis, parasitised by Heterostylum

robustum in U.S.A., 565.

Nomia australiaca, pollinating lucerne in S. Australia, 234.

Nomia melanderi (in U.S.A.), as pollinator of lucerne, 565; bionomics and natural enemies of, 565; nesting sites of, 140, 299.

Nomophila noctuella, on Douglas-fir seedlings in Washington, 478; hep-tachlor spray against, 478.

nonagrioides, Sesamia.

Norbanus scabriculus (attacking parasitised Cephus spp.), in U.S.S.R., 161; introduced into Canada, 161.

nordmannianae, Chermes.

norvegicoides, Vespula norwegica.

norvegicus, Calocoris.

Norway, outbreak of *Plutella maculi*pennis in, 430; handbook on diseases and pests of field crops in, 622.

Norway Maple (see Acer platanoides).

norwegica, Vespula.

Nosema, infecting Oiketicus kirbyi in Costa Rica, 638; Tyria jacobaeae infected by undescribed species of, 260.

Nosema curvidentis, sp.n., infecting Ips curvidens in Slovakia, 283.

Nosema locustae, infectivity of, to locusts, 582; effects of, on pigmentation and flight of Locusta migratoria, 582.

Nosema tortricis, infecting Tortrix viridana in Czechoslovakia and Germany, 105. notatus, Pissodes.

Nothroholaspis tridentinus, in stored grain in Canada, 154.

Nova Scotia, apple pests in, 159, 259, 387, 388, 523; ecological effects of chemicals on fauna of apple in, 262; Lepidoptera on cabbage in, 525; wireworms attacking grasses, potatoes and turnip in, 73; forest pests in, 121, 394, 458, 529; Panonychus ulmi in, 528; natural enemies of pests in, 159, 259, 388, 394, 523.

novemnotata, Coccinella.

noxius, Anodontonyx.

nubilalis, Ostrinia (Pyrausta).

nubilosus, Hypolixus.

Nudaurelia cytherea capensis, on pines in S. Africa, 425; aircraft application of sprays against, 425.

nuesslini, Dreyfusia (see Chermes nord-

mannianae).

numismatica, Toumeyella. nursei, Goryphus (Melcha).

Nutgrass (see Cyperus).

Nyasaland, survey of data on Hemiptera of cotton in, 352; first record of *Contarinia sorghicola* on sorghum in, 684. nymphaea, Ephesia (Catocala).

nymphagoga, Catocala.

Nysius graminicola, on cotton in Turkey, 495.

Nysius vinitor, characteristics and world distribution of, 506.

Nythobia (Angitia) (parasitising Lepidoptera), in New York, 401; in Ontario, 162; in Yugoslavia, 138; synonymy and type species of, 493.

Nythobia (Angitia) armillata, parasitising Hyponomeuta padellus malinellus in

Finland, 195.

Nythobia chrysostictos, study on multiparasitism of Anagasta kuehniella by Nemeritis canescens and, 618, 619.

Nythobia exareolata, parasitising Tortricids in Czechoslovakia, 683.

Nythobia (Angitia) fenestralis, parasitising Plutella maculipennis in U.S.S.R., 87, 245; bionomics of, 245; experiments with Galleria mellonella and, 87.

Nythobia (Meloboris) pusio, 493.

Nythobia (Angitia) punctoria, parasitising Ostrinia nubilalis in France, 372.

0

Oak, Operophtera brumata on, in Belgium, 85; Lepidoptera on, in Czechoslovakia, 683; pests of, in Germany, 106, 140, 248, 250, 376; Pelatea festivana on, in Italy, 535; pests of, in

Spain, 353, 355, 659; pests of, in U.S.S.R., 190, 597, 598, 599, 620; pests of, in U.S.A., 88, 271, 336, 337, 407, 418; pests of, in Yugoslavia, 31, 55, 181; insects infesting acorns of, 88; development of Aegeria vespiformis in cankers on, 539; Antheraea pernyi feeding on dried leaves of, 192; Apechthis rufata parasitising pupae of Choristoneura murinana transferred to, 168; influence of birds on insect fauna of, 494; (timber), Coleoptera in, 31, 489, 622; factors affecting infestation of, by Lyctus, 622; treatment of, 489. (See also Quercus spp.).

Oak, Cork (see Quercus suber).
Oak, Evergreen (see Quercus ilex).

Oak, Mongolian (see *Quercus mongolica*).
Oak, Red (see *Quercus falcata* and *Q. rubra*).

Oak, White (see Quercus alba).

Oats, pests of, in Britain, 319, 320, 321; pests of, in Bulgaria, 348, 537; pests of, in Canada, 70, 162, 394, 527, 529, 585; Calligypona pellucida on, in Finland, 102; Oscinella frit on, in Germany, 255; Anodontonyx noxius on, in New South Wales, 493; pests of, in U.S.A., 95, 401; Eurygaster integriceps on, in U.S.S.R., 14; Camarota curvipennis on, in Yugoslavia, 249; Hemiptera and virus diseases of, 132, 133, 134, 401; factors affecting resistance of varieties of, to Cephus cinctus, 72, 527; stage or variety of, in relation to infestation by Calligypona pellucida, 103; survival of Chloriona orizicola on, 132; question of Meromyza saltatrix on, 96, 97; oviposition sites of O. frit on, 255; as food-plant for Toxoptera graminum, 585; seed treatment of, 401; effects of insecticides and residues on, 401, 562, 578; effects on vetch fauna of sprays against pests of, **211.** (See also Avena).

Oats (Stored), insects in, 154, 155, 554; mites in, 154, 155; sorption of methyl bromide by, 554; effects of fumigants

on germination of, 579.

Oats, Rolled, in diets for Coleoptera, 656, 657.

Oats, Wild, Eurygaster spp. on, in California, 520.

obesa, Carcelia.

obesi, Microtermes.

obesus, Dendroctonus; Odontotermes.

obliquus, Eumerus.

obovatus, Brevipalpus.

obscura, Leucopis (Neoleucopis); Phorocera.

obscurator, Orgilus.

obscuricolor, Macrosoma brunnipes. obscurus, Habrolepis; Hylastes (Hylastinus); Phytocoris; Tenebrio.

obsoletus, Hyalesthes.

obtectus, Acanthoscelides.

obtusa, Melanagromyza (Agromyza). occidentalis, Casca; Frankliniella; Praon; Typhlodromus.

ocellana, Spilonota (Tmetocera).

ocellatella, Scrobipalpa (Gnorimoschema, Phthorimaea).

ochraceus, Anacanthotermes.

Ochsenheimeria vacculella, habits and bionomics of, on cereals and grasses in U.S.S.R., 596; natural enemies of, 596; insecticides against, 596.

Ocneria monacha (see Lymantria).

Octachlorodipropyl Ether, toxicity of, to Lepidoptera, 476; as synergist: with carbamates, 476; with pyrethrum, 2, 3.

1,3,4,5,6,7,8,8-Octachloro-3a,4,7,7a-tetrahydro-4,7-methanophthalan, against Coleoptera, 396, 515; against Lygus lineolaris, 515; against Noctuids, 269, 413, 465, 515; against Pyralids, 21, 469, 503; against other Lepidoptera, 269, 425, 465, 513; against Thysanoptera, 477, 503; banana fruit damaged by, 503; in granules, 396, 469; in sprays, 21, 269, 413, 425, 465, 477, 503, 515; applied from aircraft, 425; soil treatment with, 396; and methyl-parathion, 413.

octasema, Nacoleia.

Octosporea viridanae, infecting Tortrix viridana in Czechoslovakia and Germany, 105.

Odontotermes, in India, 627; treatment of nests of, with insecticides, 627.

Odontotermes assumthi, sett treatments of sugar-cane against, in India, 119.

Odontotermes gurdaspurensis, in India, 627; treatment of nests of, with insecticides, 627.

Odontotermes obesus, in India, 627; treatment of nests of, with insecticides, 627.

Oecophylla, in Congo, British Cameroons, Nigeria, N. Rhodesia and Tanganyika, 28; damage to coconut by Pseudotheraptus wayi reduced by, 28; ineffective against Diocalandra frumenti, 29; other prey of, 29; bionomics and natural enemies of, 28, 29; effects of insecticides on populations of, 29; labelled with ³²P, 29.

Oecophylla longinoda, association of, with Coccus hesperidum in Nigeria, 531; question of identity of, 28.

Oecophylla smaragdina, rôle of, in control of Promecotheca papuana in New

Britain, 166; occurrence of Amblypelta lutescens papuensis in absence of, in Papua, 235; question of O. longinoda

in relation to, 28.

Ohio, pests of beans in, 279, 328, 397, 413, 463; Ostrinia nubilalis on maize in, 152, 215; Rhyacionia spp. on pines in, 20, 481; pests of tomato in, 573; Myzus persicae on vegetables and annual flowering plants in, 573; Tyrophagus putrescentiae on mushrooms in, 573; beneficial insects in, 20.

Oiketicus kirbyi, bionomics of, on banana in Costa Rica, 637, 638; natural enemies and diseases of, 638; insecti-

cides against, 638.

Oil, Mustard (see Allyl Isothiocyanate).

Oil, Pine (see Pine Oil). Oil, Tar (see Tar Oil).

Oil Emulsions, against aphids, 616, 617, 631, 667, 668, 674; ineffective against Eriosoma lanigerum, 312; against Coccids, 120, 167, 221, 227, 434, 470, 557, 616, 617, 666, 667; against other Hemiptera, 158, 411, 412, 436; against Lepidoptera, 15, 20, 256, 534, 663, 668, 669; against mites, 8, 510, 604, 616, 617; effects of, on beneficial arthropods, 35, 36, 158, 312, 434; against fungus disease of banana, 462; study on distribution of deposits of, on Citrus, 221; fluorescent dye as tracer with, 221; and DDT, 434, 631; toxicity of DDT decreased by, 20; and dieldrin, 463, 667; and DNC, 412, 434, 534; and lime-sulphur, 411; and phosphorus insecticides, 8, 9, 120, 167, 256, 312, 434, 510, 534, 557, 604, 616, 617, 663, 667, 668, 669; and tar distillate, 434.

Oil Palm (see Elaeis guineensis).

Oil-seed Cake, insects intercepted in importations of, into Germany, 10, 11. Oils, in baits for fire ants, 464; concentrations of, in sprays against Coccids and overwintering stages of Panonychus ulmi and Aphis pomi, 616, 617; viscosity of, in relation to use against Eulecanium corni, 470; in dormant sprays against Phylloxera vitifoliae, 84; against Lepidoptera, 534; toxicity of, to beet, 670; aircraft application of insecticides in, 55, 211, 659; treatment of timber with preparations containing, **181**, **647**, **648**; other uses of, as solvent or carriers for insecticides, 2, 3, 4, 31, 39, 117, 118, 399, 413, 425, 651, 669, **670, 686, 687;** types of, in relation to insecticide treatment, 104; effects of, on insecticides, 145, 287, 472; and creosote, 181; and parathion, 534;

and tar distillate, 534. (See Kerosene)

Oils, Essential, relation of content of, in pine needles to infestation by *Dendrolimus pini*, **599**.

Oils, Vegetable, stability of heptachlor not influenced by, 145. (See Castor, Cottonseed, Groundnut, Linseed, Maize, Olive and Soy-bean Oils.)

Oklahoma, *Toxoptera graminum* on wheat in, 219, 470; pests of greenhouse plants in, 339.

Okra (see Hibiscus esculentus).

oleae, Dacus; Parlatoria; Saissetia.

Oleander, *Pseudaulacaspis pentagona* on, in Bermuda, 229; *Saissetia oleae* on, in Pakistan, 491.

Olearia fosteri, Agriophara coricopa on, in New Zealand, 75.

olearia, Chrysocraspeda.

oleellus, Prays.

oleivora, Phyllocoptruta (Phyllocoptes).

Oleoethion, 604, 667.

Oleoparathion, 8, 9, 84, 256, 312, 434, 534, 663, 666, 667, 668.

Oleophosdrin, 667.

Oleoresin, infestation by bark-beetles in relation to exudation of, by pines, 340. oleracea, Diataraxia (Mamestra, Polia); Haltica; Tipula.

oleraceum, Eurydema.

Oligonychus, species of, on Hisbiscus esculentus in India, 317; on date in Morocco, 54; insecticides against, 317. Oligonychus coffeae, acaricides against.

on tea in Assam, 288; effects of Tedion on developing eggs of, 288, 289.

Oligonychus ilicis (see Paratetranychus).
Oligonychus peruvianus, acaricides against, on cotton in Peru, 559.

Oligonychus (Paratetranychus) simplex, species referred to as, on date in Morocco, 54.

olivaceus, Phytodecta.

Olive, Parlatoria oleae on, in California, 227; Dacus oleae on, in France, 435; D. oleae on, in Greece, 237, 238; pests of, in Italy, 40, 46-48, 135, 308, 310, 438, 532, 533, 694; pests of, in Spain, 99; pests of, in Turkey, 81, 177; Prays oleellus on, in Yugoslavia, 609; effects of ripening or skin characters on infestation of, by D. oleae, 47, 238; review of pests and diseases of, 693; residues of insecticides in (see also next entry), 237, 238; metabolism of dimethoate in fruit of, 562.

Olive Oil, residues of insecticides in, 47, 48, 135, 238, 309, 439, 563; removal of parathion residues from, 309; as

solvent for insecticides, 321.

olivina, Euphyllura.

Omophlus, species of, in Bulgaria, 348. Omophlus lepturoides, bionomics of, on plants in Bulgaria, 348.

Omophlus picipes, in Bulgaria, 348.

Omophlus proteus, bionomics of, on plants in Bulgaria, 348.

Omorga (see Campoplex).

Omphale montana, parasitic on Leucoptera scitella in Italy, 136.

Omphisa anastomosalis, characteristics and world distribution of, 507.

Oncopeltus fasciatus, light wave-length preferences of, 518; toxicity of extract from Bacillus larvae to, 332; absorption and metabolism of insecticides in, 263, 512, 513; enzyme activity of homogenates of, on phosphorus compounds, 213; used in tests of carbamate insecticides, 516.

Oncopera, larval taxonomy of, 424.

Oncopera fasciculata, influence of, on pastures in S. Australia, 504.

Oncopera intricata, in pastures in Tasmania, 424.

Oncopera rufobrunnea, in pastures in Tasmania, 424.

Onion, new virus disease of, in Bulgaria, 345; Hylemya antiqua on, in Denmark, 608; Trioza nigricornis on, in Italy, 436; H. antiqua on, in Quebec, 455; Exora collaris on, in Turkey, 82; pests of, in U.S.A., 226, 477; use of, in experiments on feeding by Myzus ascalonicus, 92; seed treatment of, 226, 608; residues of heptachlor in, **479**; (stored), *M. ascalonicus* on, **492**.

onobrychidis, Eurytoma. Onobrychis viciifolia (sativa), Eurytoma onobrychidis infesting seeds of,

Hungary, 305.

Ononis spinosa, Eurytoma ononis infesting seeds of, in Hungary, 306.

ononis, Eurytoma.

Ontario, pests of cereals and grasses in, 70, 71, 161, 162, 163, 164, 276, 389, 390, 523, 524, 529; virus diseases of cereals and grasses in, 71, 523, 524; pests of leguminous forage plants in, 71, 529; pests of crucifers in, 161, 162, 163, 211, 385, 529; Leptinotarsa decemlineata on potato in, 207, 377; pests of other vegetables in, 125, 529; virus diseases of lettuce, carrot and celery in, 70; forest pests in, 24, 25, 73, 74, 122, 123, 124, 158, 163, 206, 208, 261, 387, 392, 393, 457, 458, 529; Ceratocystis ulmi on elm in, 124; orchard pests in, 154, 163, 164, 404, 522, 529; Myzus ribisnigri on Ribes alpinum in, 529; Malacosoma americanum on Prunus virginiana in 343; Aphrodes bicinctus on strawberry in, 686; tobacco pests in, 388; Drosophila spp. in, 327; M. disstria in, 140; Trogoderma parabile recorded from, 529; Typhlodromus caudiglans misidentified as T. rhenanus in, 522; Acarapis externus infesting honey bees in, 295; diseases of insects in, 389, 390, 459; beneficial insects and biological control in, 24, 25, 70, 74, 129, 140, 155, 157, 158, 161, 162, 164, 209, 377, 387, 393, 459, 522, 529, 664; beneficial nematode introduced into, 162, 207; Coleoptera controlling Linaria vulgaris in, 386.

Ooencyrtus, parasitising **Ambrostoma** quadriimpressum in China, 593; eggparasite of Amblypelta lutescens papuen-

sis in Papua, 235.

Ooencyrtus fecundus, sp.n., egg-parasite of Pentatomids in Morocco, 654, 655; parasitising eggs of Lepidoptera, 654.

oencyrtus kuvanae (egg-parasite of Lymantria dispar), in Bulgaria, 367; Ooencyrtus in U.S.A., 399, 400; in Yugoslavia, 54, 367; bionomics of, 400; persistence of, in sprayed areas, 399.

Ooencyrtus nigerrimus, sp.n., egg-parasite of Pentatomids in Morocco, 654, 655; parasitising eggs of Lepidoptera, 654. Ooencyrtus telenomicidus, egg-parasite of Pentatomids in Morocco, 654, 655;

parasitising eggs of Lepidoptera, 654. opacipennis, Circulifer.

Phthorimaea operculella, (Gnorimoschema).

Operophtera brumata, in Belgium, 85; in Canada, 257, 378; in France, 257, 669; in Germany, 250, 257; in U.S.S.R., 190, 359; characteristics and world distribution of, 507; on forest trees, 85, 190, 250; on orchard trees, 85, 359, 669; bionomics of, 190; natural enemies of, 190, 251, 257, 258, 378; experiments with Thelohania weiseri and, 105; other factors reducing populations of, 190; tests and uses of insecticides against, 85, 669.

Opius, use of species of, parasitising

Dacus oleae in Italy, 532.

Opius concolor (parasitising Trypetids), introduced into France, 435; in Morocco, 435; oviposition preferences

Opius fulvicollis, parasitising Pegomyia betae in Germany, 442.

Opius nitidulator, parasitising Pegomyia betae in Germany, 442.

Opomyza germinationis, on grasses in England, 488.

opsiphanis, Apanteles.

Ora picta, association of, with rice in India, 315.

Orange, pests of: in Brazil, 556, 644, 645; in Israel, 611, 614, 684; in Italy, 181, 299; in U.S.A., 267, 409; new Tortricid feeding on, 230; mites in relation to leprosis of, 556; effects of acaricides on yields of, 645; insecticide residues in fruits of, 267, 561, 574.

orariensis, Eriococcus. orbonalis, Leucinodes. orchestis, Chrysocharis. orchidii, Chaetanaphothrips.

Orchids, *Xyleborus morigerus* on, in Java and New Guinea, **625**; damage by cellulose polymer to, **272**. (See also *Cattleya*.)

Orconectes, experiments on physiological activity of body fluids from insecticide-treated, 63.

Orconectes propinquus, 63. Orconectes virilis, 63.

ordinarius, Apanteles.

Oregon, yellows virus of beet in, 17; Sitona lineatus on peas in, 134; pests of cereals in, 211; forest pests in, 185, 209, 385, 386, 532; pests of leguminous forage crops in, 134, 211, 275, 417; aphids on Easter lily in, 278; mites in stored grain in, 149; mites attacking Ips spp. in, 386.

oregonensis, Contarinia; Listronotus;

Monochamus. oregonis, Ips.

Orgilus obscurator, established in Ontario as parasite of Rhyacionia buoliana, 73. Orgyia antiqua, on elm in Holland, 42. Orgyia basalis, on pines in S. Rhodesia, 81.

Orgyia gonostigma, bionomics of, on apple, apricot and pear in Switzerland, 46; insecticides against, 46.

orientalis, Achrysocharella; Aonidiella; Eutetranychus.

Orius, predacious on cotton pests in Arizona, 639; predacious on Pieris rapae in Japan, 78.

orizicola, Chloriona (Sogata).

Ormenis pruinosa (see Metcalfa).

ornatum, Eurydema.

ornatus, Ischnurgops; Myzus.

ornigis, Apanteles.

ornithogalli, Prodenia.

Oroscopa hacupha, damaging Heliconia in Costa Rica and Honduras, 520.

Orotic Acid, larval growth of Corcyra cephalonica inhibited by, 41.

orris, Hydropsyche. orthogonia, Agrotis.

Orthophosphoric Acid, use of, for labelling Anthonomus grandis, 471;

absorption of, by cotton seed, 224; labelled with ³²P, 224, 471.

Orthosia incerta, bionomics of, on apple, pear and forest trees in Switzerland, 46; sprays against, 46.

Orthosia stabilis, bionomics of, on apple, pear and forest trees in Switzerland, 46; sprays against, 46.

Orthotran (see Chlorfenson).

Oryza ridleyi, tests on susceptibility to Chilo suppressalis of rice and, 78.

Orvza sativa (see Rice).

oryzae, Agromyza; Chironomus; Chlorops; Latheticus; Pachydiplosis; Sitophilus (Calandra).

Oryzaephilus mercator, in stored groundnuts in Gambia, 178; intercepted on oil-seed cake in Germany, 11.

Oryzaephilus surinamensis, in stored products: in Canada, 155; in Germany, 11; in Kenya, 51; tests and uses of insecticides against, 51, 202.

oryzophilus, Lissorhoptrus.

OS-1836 (see Diethyl 2-Chlorovinyl

Phosphate).

Oscinella (Oscinis, Oscinosoma) frit, in Britain, 199, 319, 320, 430; in Bulgaria, 537; in Germany, 45, 255; in Hungary, 306; in E. Pakistan, 174; in U.S.S.R., 12, 244, 446, 595; on barley, 45, 537; on maize, 12, 306, 446, 495; varietal susceptibility of maize to, 595; on oats, 255, 319, 320, 537; on rice, 174; on wheat, 244, 595; on grasses, 45, 430; study of food-plants in relation to. 595; bionomics and adult habits of, 12, 174, 255, 319, 320, 446; vertical distribution of, 30, 199; fluctuations in populations of, 45; natural enemies of, 12, 45; trap catches of, 319; detection of, by X-rays, 495; insecticides against, 45, 46, 430, 537.

Oscinella frit pusilla, on maize in U.S.S.R., 12; parasites of, 12.

Oscinella pusilla (see O. frit pusilla).
Ostrinia nubilalis, in Britain, 488; in Canada, 27, 162, 163, 164, 258, 276, 514; in China, 502; in France, 372; in Hungary, 306; in Poland, 365; in Rumania, 56; in U.S.S.R., 12, 14, 245; in U.S.A., 4, 152, 164, 215, 225, 275, 276, 306, 325, 329, 333, 398, 461, 509, 511, 630; on Capsicum, 630; on eggplant, 12; on grasses, 488; on hemp, 14, 245, 306; on maize, 12, 27, 56, 152, 162, 163, 164, 215, 225, 245, 258, 275,

162, 163, 164, 215, 225, 245, 258, 275, 276, 306, 325, 329, 333, 365, 372, 398, 461, 502, 509, 511, 514; varietal susceptibility of maize to, 12, 56, 152, 153,

215, 333, 398, 476; on other cereals, 14; factors for resistance to, in cereals,

22, 68; on potato, 333; on wormwood, 14; food-plant preferences of strains of, 14; relation of, to stalk rot, 398; detection of growth inhibitors in other plants by, 22; bionomics and adult habits of, 12, 27, 56, 306, 365, 372, 502; factors affecting diapause and development of, 4, 164, 276; cold-hardiness of, 4, 502; sampling for construction of life tables for, 258; mortality factors of, 258; characters of fringe populations of, 461; relation of Nitidulid to, 329; light-trap catches of, 306, 630; grassland management in relation to, 488; fungus infesting, 164; natural enemies of, 27, 162, 164, 245, 246, 372, 509, 511; hot-water treatment of eggs of, against infection by Perezia pyraustae, 343; insecticides against, 12, 164, 225, 325, 333, 372, 514, 630; soil treatment with phorate ineffective against, 333; other measures against, 12, 56, 372, 514.

Otiorhynchus cribricollis, bionomics of, on globe artichoke and blackberry in California, 509; fungus infesting, 509; tests of insecticides against, 509.

Otiorhynchus ligustici (on lucerne), in Bulgaria, 60; in New York, 280, 399; soil treatments against, 280; statistical analysis of experiments on control of, 399.

Otiorhynchus ovatus, on Mentha spp. in Washington, 324; soil treatments against larvae of, 324.

Otiorhynchus sabulosus, on olive

Turkey, 177.

Otiorhynchus sulcatus (on vines), in Germany, 108, 109, 255; transmission of mottling virus by, 304; bionomics and ecology of, 109; measures against, 108, 109, 255.

ou, Rachiplusia.

ovatus, Otiorhynchus (Brachyrhinus).

Ovex (see Chlorfenson).

Ovotran (see Chlorfenson).

Oxycarenus hyalinipennis, characteristics and world distribution of, 507.

Oxygen, treatment of Narcissus bulbs with, against Merodon equestris, 42.

oxylus, Campoletis.

Oxythyrea funesta, bionomics of, on cabbage and other plants in Bulgaria, 349; tests of insecticides against, 349.

P

Pachyceras xylophagorum (see Roptrocerus).

Pachydiplosis orvzae, characteristics and and world distribution of, 507.

Pachygonatopus minimus, parasitising Macrosteles fascifrons in Manitoba, 551.

Pachynematus alaskensis, pathogenicity of Bacillus cereus to, 283.

Pachynematus clitellatus, on wheat and grasses in U.S.S.R., 345.

Pachynematus scutellatus, on spruce in Czechoslovakia, 246; parasites of, 246. Pachyscapus, classification and new

species of, 493.

Pacific Islands, pests of coconut in, 81; Leptoglossus australis in, 235. (See also Fiji, New Britain and Solomon Islands.)

Pacific Silver Fir (see Abies amabilis).

pacificus, Tetranychus.

padellus, Hyponomeuta. padi, Eriophyes; Rhopalosiphum.

paenulata, Epilachna.

Paillotellavirus pieris, 401.

painei, Actia.

Paint, protection of timber with, against termites, 486.

Pakistan, Sphenoptera gossypii on cotton in, 176; mite on litchi in, 88; Cricula trifenestrata on mango in, 88; Dacus cucurbitae on melon in, 175; pests of other fruits in, 169, 170; rice pests in, 174; pests and virus diseases of other cereals in, 684; Saissetia coffeae on Rauwolfia serpentina in, 176; review of insect pests of pulses in, 196; Epilachna varivestis on vegetables in, 88; Coccids on ornamental shrubs in, 170, 491; Larvacarus transitans on Ziziphus jujuba in, 176, 626; pests of stored products in, 236, 295; virus disease of Pseudaletia unipuncta in, 175, 693; beneficial insects in, 150, 170, 175, 491; parasites of Quadraspidiotus perniciosus introduced into, 170.

palaestinensis, Calliptamus.

palestinensis, Tropidophryne.

pallida, Gilpinia (Diprion); Podagrica.

pallidicornis, Chirothrips. pallidipennis, Mesites.

pallidula, Pheidole.

pallidus, Creontiades; Dalopius; Steneotarsonemus (Tarsonemus); Trioxys.

pallipes, Aphaereta; Microdiprion (Dip-

Palm, Oil (see Elaeis guineënsis). palmarum, Rhynchophorus.

Palorus shikhae, sp. n., predacious on Latheticus oryzae in India, 626.

paludosa, Tipula.

Pamphilius balteatus, on roses in U.S.S.R.,

Pandanus, Nacoleia octasema on species of, in Indonesia, 354.

Pandanus tectorius, 354.

Pandemis ribeana, bionomics of, on fruit trees in U.S.S.R., 360.

pandurus, Lygaeus.

Pangola Grass (see Digitaria decumbens).

experiments with Panicum, Ostrinia nubilalis and, 14.

Panicum crusgalli (see Echinochloa).

Panicum maximum, Laelia exclamationis on, in India, 79; Blissus leucopterus on, in Venezuela, 98.

Panicum purpurascens, Blissus leucopterus

on, in Venezuela, 98.

Panolis flammea (on pines), in Germany, 376; parasite of, in Poland, 379; in U.S.S.R., 598; characteristics and world distribution of, 507; effects of soil fertilisers on populations of, 376. Panonychus, on soy bean in Delaware, 414.

Panonychus citri (on Citrus), in California, 147, 509; in U.S.S.R., 242; inclusion bodies associated with virus disease of, 581; toxicity of methylcarbamates to, 516; resistance to acaricides in strains of, 147, 509, 510; use of, in rearing media for predactious mites, 641.

Panonychus ulmi, in Austria, 302; in Britain, 379, 671, 672; in Canada, 159, 259, 404, 454, 523, 528; in France, 8, 255, 256, 604, 666; in Germany, 108, 439, 440; in Holland, 444; in Italy, 616, 663; in Japan, 35, 36; in Poland, 189, 544, 677, 678; in U.S.S.R., 242, 360, 361; in U.S.A., 266; in Yugoslavia, 241, 242; on fruit trees, 8, 35, 36, 159, 189, 241, 242, 255, 256, 259, 266, 302, 360, 361, 379, 404, 410, 439, 440, 444, 454, 523, 544, 604, 616, 666, 671, 672, 677, 678; pear leaf scorch caused by, 404; effects of infestation by, on growth of plum trees, 671; resistance of apple varieties to, 242; on vines, 8, 108, 663; bionomics of, 242, 361; factors affecting oviposition and diapause in, 361; cold-hardiness of forms of, 528; natural enemies of, 302, 379, 439, 454, 522, 672; experiments with, as prey for Typhlodromus pyri, 122; effects of plant treatment and nutrition on populations of, 256, 259, 410; effects of insecticides and cultural practices on populations of, 8, 35, 36, 108, 444, 523, 544, 666; measures against, 8, 159, 189, 242, 266, 439, 440, 444, 454, 544, 604, 616, 672, 677, 678, 685; resistance to acaricides in strains of, 266, 672.

Pansepta teleturga, on cacao in Dutch New Guinea, 76.

Papaver somniferum, Mamestra brassicae

on, in Germany, 11.

Papaya. Hemitarsonemus latus in Brazil, 120; Aonidiella orientalis on, in China, 491; toxicity of insecticides

Paper Chromatography, of amino acids and auxins in secretions of aphids, 95, 629; determination of insecticides by, 228, 512, 552.

Papilio demoleus, characteristics

world distribution of, 507.

Papilio machaon, on Citrus in Turkey, 82. Papua and New Guinea, Territory of, Amblypelta lutescens papuensis on coconut, rubber, cassava and Ipomoea in, 235; ants in, 235; food-plants and distribution of Leptoglossus australis in, 235; Pericyma cruegeri on Poinciana regia in, 300; parasites of A. l. papuensis in, 235.

papuana, Promecotheca.

papuensis, Amblypelta lutescens.

Parabates latungula, parasitising larvae of Operophtera brumata in Europe, 378; introduced into Canada, 378.

parabile, Trogoderma.

Paraffin, use of, in rearing Ips, 390.

Parahormius leucopterae, parasitising Leucoptera in Tanganyika, 652.

Paralobesia viteana, sprays against, on

vines in Pennsylvania, 398.

Paramyelois transitella, on almonds and walnuts in California, 276, 482; bionomics of, 276; Cydia pomonella and Bacterium juglandis in relation to, 276; measures against, 276.

Paranthrene tabaniformis, characteristics

and world distribution of, 507.

Paraoxon, enzymatic hydrolysis of, by arthropod and mammalian homogenates, 213.

Parasetigena silvestris (see Phorocera).

parasitellae, Apanteles. parasitica, Sturmiopsis.

Paratetranychus ilicis, in Japan, 452; in New York, 19; on azalea, 19, 452; on Ilex crenata, 19; on tea, 452; characters of, 452; tests and uses of systemic insecticides against, 19.

Paratetranychus peruvianus (see Oligonychus).

Paratetranychus pilosus (see Panonychus ulmi).

Parathion, against aphids, 59, 64, 84, 188, 192, 279, 324, 468, 498, 532, 537, 549, 556, 569, 604, 608, 616, 617, 624, 633, 637, 643, 667, 668, 674; effects of, on respiration of aphids, 444; resistance

to, in strains of aphids, 148, 412; against Coccids, 167, 257, 298, 434, 612, 616, 666, 667; against other Hemiptera, 38, 39, 57, 213, 437, 498, 265, 314, 471, 497, 603, 647, 674; against Agromyzids, 139, 484, 674; against Trypetids, 47, 48, 98, 135, 137, 237, 238, 309, 311, 438, 533, 535, 603, 668; against other Diptera, 45, 101, 226, 254, 516, 644; resistance to, in strain of Musca domestica, 263; against Noctuids, 179, 347, 398, 512; against Pyralids, 61, 80, 145, 234, 328, 329; ineffective against Pyralids, 297; against Tineids, 31, 111, 347, 441, 448, 533, 536, 663; against Tortricids (sens. lat.), 9, 20, 57, 108, 248, 256, 337, 434, 466, 538, 558, 637; against other Lepidoptera, 7, 35, 46, 85, 154, 226, 267, 309, 326, 335, 480, 534, 535, 641, 668, 669; acceleration of embryonic development of Bombyx mori by, 437; toxicity of, to Locusta migratoria, 172; against Tenthredinids, 56, 139, 541, against other Hymenoptera, 110, 422; against Thysanoptera, 193, 345, 346, 477, 499, 569; against Tetranychids, 8, 183, 317, 324, 517, 604, 616, 628; Tetranychids favoured by, 256, 666; resistance to, in strains of Tetranychus spp., 278, 511, 512, 614; against other mites, 191, 298, 675; development of Steneotarsonemus fragariae stimulated by, 545; ineffective against Larvacarus transitans, 626; against Scutigerella immaculata, 433; effects of, on natural enemies of pests, 35, 36, 159, 227, 312, 434, 469, 627; toxicity of, to bees, 412; factors affecting toxicity of, 20, 47, 328, 329, 471; humidity in relation to fumigant action of, 471; metabolism 45, 47, 56, 57, 58, 61, 80, 84, 85, 108, 110, 111, 135, 137, 138, 139, 154, 159, 167, 183, 188, 191, 192, 193, 212, 213, 226, 234, 237, 248, 256, 257, 267, 279, 298, 309, 311, 312, 317, 324, 326, 328, 335, 336, 337, 345, 346, 347, 398, 412, 434, 437, 438, 441, 466, 468, 477, 484, 498, 499, 512, 517, 532, 533, 534, 535, 536, 537, 538, 541, 545, 549, 554, 556, 558, 604, 608, 616, 617, 624, 626, 632, 632, 641, 608, 616, 617, 624, 626, 633, 637, 641, 643, 644, 663, 666, 667, 668, 669, 674, 675; in bait-sprays, 98; in dormant sprays, 84; applied from aircraft, 569;

as deposit, 278; persistence of deposits of, 356; in transplanting dips, 673, 674; in drench, 254; treatment of sugar-cane setts with, 297; seed treatment with, 226, 433, 674; soil treatment with, 7, 171, 226, 468, 578; persistence and toxicity of, in soils, 83, 416, 578; treatment of nests with, 422; treatment of timber with, 647; tests and uses of, against pests of stored products, 212, 213, 448; translocation of, 83; residues of, in plants and plant products, 47, 135, 238, 267, 309, 398, 439, 644; tolerance limits for, 267; removal of residues of, 309; effects of, on germination and growth of plants, 562, 675; toxicity of, to strawberry, 675; and BHC, 532; and copper sulphate, 47, 309, 438; effects of copper salts and bordeaux mixture on efficiency of, 135, 439; and DDT, 39, 137, 248, 311, 437, 438, 466, 534, 535, 637; and dieldrin, 477; and ferrous sulphate, 36; and Kelthane, 532; and malathion, 154, 612; and methyl-parathion, 191, 647; and oil emulsions, 8, 9, 84, 167, 256, 312, 434, 534, 616, 663, 666, 667, **668**, **669**; and zinc sulphate, **36**; antagonistic action of synergists with, 63, 64.

Paratoil, containing parathion (q.v.), 183. Paratrioza cockerelli, on potato in U.S.A., 473, 510; sex determination of nymphs of, 496; insecticides against, 474, 510. Pardalaspis cyanescens, map of distribution of, 357.

pardalina, Myiopardalis.

Paremydica divareti, sp.n., on Cola in Cameroun Republic, 355.

Paremydica pujoli, sp.n., on Ivory Coast and Guinea, 355. Cola in

pariana, Anthophila (Simaethis). parkeri, Tersilochus (Porizon).

Parlatoria blanchardii, on date Morocco, 54; natural enemies of, 54. Parlatoria oleae, sprays against, on olive in California, 227; map of distribution of, 357; parasite of, 227.

Parlatoria ziziphus, characteristics and world distribution of, 507.

Parthenolecanium corni (see Eulecanium)... parumpunctatus, Rhopalus (Corizus).

parviflori, Amphorophora.

parvula, Epitrix.

parvulus, Pediobius (Pleurotropis).

Paspalum dilatatum, not infested by Hyperodes bonariensis in New Zealand,

Passerinia fragaefolii (see Capitophorus). Passiflora edulis, Leptoglossus australis on, in New Guinea, 235.

Passiflora quadrangularis, Calacarus citrifolii on, in S. Africa, 83.

passiflorae, Corythaica.

Passion Fruit (see Passiflora edulis). pastranai, Apanteles; Hemiteles.

Pasture (see Grasses).

Patanga succincta, cannibalism among

hoppers of, 172. patruelis, Pentodon. paulensis, Monoctonus. pauperculus, Stethorus. PCPBS (see Fenson).

Pea, pests of, in Britain, 287, 431; Bruchus pisorum on, in Bulgaria, 58, 537; Sitona spp. on, in Finland, 102; pests of, in Holland, 193; Melanagromyza phaseoli on, in India, 173; new Agromyzid on, in Queensland, 295; S. lineatus on, in U.S.A., 134; Bruchidius incarnatus on, 249; varietal susceptibility of, to M. phaseoli, 173; effects on Macrosiphum pisum of feeding on resistant varieties of, 453; crop losses of, in relation to infestation, 193, 431; systemic activity of chlorinated hydrocarbons in, 146; metabolism of mevinphos in, 88; effects of insecticide residues on growth and respiration of, 562.

Pea, Meadow (see Lathyrus pratensis). Pea Moth (see Cydia nigricana).

Peach, Aculus cornutus on, in S. Africa, 83; Cydia molesta on, in Brazil, 555, 646; Neurotoma nemoralis on, Bulgaria, 540; pests of, in France, 8, 84, 255, 256, 664, 665, 666, 667, 668, 669; as winter food-plant of Myzus persicae in Germany, 547; Tetranychus telarius on, in Holland, 682; pests of, in Italy, 181, 310, 437, 616, 662; pests of, in Ontario, 154, 163, 164, 522; Quadraspidiotus perniciosus on, in Pakistan, 170; Hyalopterus amygdali on, in Portugal, 617; Eulecanium corni on, in U.S.S.R., 13; pests of, in U.S.A., 23, 71, 147, 226, 326, 335, 336, 631, 641, 642; pests of, in Yugoslavia, 239, 240; Pseudaulacaspis pentagona on, 529; vectors of virus diseases of, 23, 71, 336; effects of nutrition of, on Tetranychid populations, 256; feeding habits of predacious mites on, 522; Thiodan treatments of planting material of, 631; toxicity of insecticides to, 8, 642; penetration of dimethoate into fruits of, 669.

Pear, Eriophyes pyri on, in S. Africa, 83; Cydia pyrivora on, in Austria, 247; Lepidoptera on, in Bulgaria, 59, 110; pests of, in Canada, 158, 163, 395, 404, 411, 529; C. pomonella on, in China, 592; pests of, in France, 8, 379, 550, 668, 669; Quadraspidiotus perniciosus on, in Pakistan, 170; Psylla spp. on, in Poland, 365; C. pyrivora on, in Rumania, 57; Agrilus sinuatus on, in Ricily, 299; pests of, in Switzerland, 46, 86, 664; Tetranychus viennensis on, in Turkey, 177; pests of, in U.S.S.R., 15, 112, 191, 242; pests of, in U.S.A., 147, leaf scorch of, caused by Panonychus ulmi, 404; treatment of cuttings of, against E. pyri, 191; question of systemic activity of dimethoate in, 395, 669; toxicity of DNC to, 8; use of, in rearing wireworms, 575.

rearing wireworms, 575.

Peat, in baits for *Drosophila*, 327; in rearing medium for *Operophtera bru-*

mata, 258. (See also Soil.)

Pecan, insects and diseases of, in U.S.A., 151, 424.

pectinatae, Chermes (Aphrastasia); Diplosis abietis (see Agevillea abietis).

Pectinophora gossypiella (on cotton), in Brazil, 557, 558, 645; in China, 502; in India, 140, 317; in Mexico, 473, 511; in Morocco, 490; measures against introduction of, into U.S.S.R., 113; in U.S.A., 219, 220, 269, 334, 338, 403, 409, 417, 473, 477, 511, 567, 568, 577, 629; other food-plants of, 511; resistance to, in races and varieties of Gossypium hirsutum, 269, 270; bionomics of, 502; factors affecting oviposition, development and diapause in, 334, 403, 465, 474, 502; airborne movement of, 577; weight and lipid and water contents of stages of, 629; sex attractant of females of, 642; plant age in relation to infestation by, 568; effects of flooding on immature stages of, 569; effects of mechanical processes on larvae of, 334, 417, 479; diseases of, 490; factors affecting susceptibility of larvae of, to Bacillus thuringiensis, 426, 427; natural enemies of, 317, 338, 490; light-traps for, 334, 477; light-trap catches of, 219, 220, 338; insecticides against, 409, 502, 557, 558, 645; resistance to DDT in strains of, 473; labelled with 32P, 338.

Pectinophora malvella (see Pexicopia). pectinicornis, Cladius.

pedalis, Pimpla.

Pediculus humanus humanus, metabolism of DDT in, 64.

Pediobius coffeicola, hyperparasite of Leucoptera spp. in Tanganyika, 652.

Pediobius furvus, parasitic on Lepidoptera in Nigeria 292.

in Nigeria, 292.

Pediobius parvulus, parasitising Promecotheca papuana in New Britain, 166.

Pegomyia, description of larvae of species of, 662.

Pegomyia betae (on beet), in Britain, 549; in Germany, 253, 442; in U.S.S.R., 542; virus-infected plants less heavily infested by, 253; factors affecting diapause and populations of, 443, 542; characters of puparium of, 442; parasites of, 442; insecticides against, 253, 443, 549; status of, 442.

Pegomyia bicolor, bionomics of, on begonia in Italy, 661; measures against,

662.

Pegomyia hyoscyami, 442.

Pegomyia hyoscyami betae (see P. betae). Pegomyia ventralis, species doubtfully identified as, on Rumex obtusifolius in Italy, 662.

pegomyiae, Phygadeuon.

pelargonii, Acyrthosiphon (see Macrosiphum malvae).

Pelatea festivana, bionomics and parasites of, on oak in Italy, 535.

pellio, Attagenus.

pellucida, Calligypona; Camnula. peltigera, Heliothis (Chloridea).

Pelts (see Skins).

Pempherulus affinis, characteristics and world distribution of, 507.

Pemphigus betae, insecticides against, on beet in Alberta, 468; predator of, 468. Pemphigus bursarius, method of rearing, 496.

Penicillium brevicompactum, infection of larvae of Malacosoma neustria by, 44. Penicillium chrysogenum, measurement of activity of resistance factors to Ostrinia nubilalis by inhibition of, 22, 68.

penniseti, Sesamia.

Pennisetum typhoides, Lepidopterous stem borers of, in Nigeria, 292, 293; destruction of débris of, in control of stem borers, 293; effects of infestation by stem borers on yields of, 292.

Pennsylvania, Panonychus ulmi on apple in, 266; Paralobesia viteana on vines in, 398; Amphorophora rubi on wineberry in, 132; aphids on lucerne in, 639; pests of cereals in, 276, 639; Pristiphora erichsonii on larch in, 210; natural enemies and disease of P. erichsonii in, 210.

Pent (chlorinated cyclopentadiene), tests of, in sprays against Coleoptera, 310, 533.

Pentachlorophenol, treatment of textiles with, against *Hofmannophila pseudospretella*, 290; soil treatment with, against termites, 180, 577.

Pentachlorophenyl Laurate, treatment of

textiles with, against Hofmannophila pseudospretella, 290.

pentagona, Pseudaulacaspis (Diaspis). n-Pentane, presence of, after fumigation of stored wheat, 215.

Pentatrichopus (see Capitophorus). Penthaleus, on wheat in China, 625.

Penthaleus major (haematopus), on lettuce in Italy, 531; survey of data on, 531; acaricides against, 532.

Pentodon patruelis, on cereals and other crops in China, 593.

Pentylenetetrazol (see 6,7,8,9-Tetrahydro-5-H Tetrazoleazepine).

Peppermint (see Mentha piperita).

Peppers (see Capsicum).

Peptone, in diets for locusts, 5, 6.

Perezia pyraustae (infecting Ostrinia nubilalis), in Delaware, 511; effects of, on stages and populations of O. nubilalis, 511; hot-water treatment against, 343.

Perfumery Chemicals, question of effect of, on efficiency of pyrethrum, 496.

Perga affinis affinis, bionomics of, on Eucalyptus in Australia, 504.

pergandei, Phenacoccus (see P. aceris). Pericyma cruegeri, on Poinciana regia in Papua, 300.

Peridroma saucia, parasitised by Hyposoter exiguae in California, 66.

Perilampus fulvicornis, hyperparasite of Rhyacionia rigidana in Ohio, 20.

Perilitus bicolor, bionomics of, parasitising Chaetocnema spp. in U.S.S.R., 244.

Perilitus rutilis (parasitising Curculionids), introduced into Canada, 391, 414; in France, 391; in Italy, 391; in Sweden, 391, 414; bionomics of, 391.

Perilloides (Perillus) bioculatus, predacious on Leptinotarsa decemlineata in Canada and U.S.A., 377; introduced into Germany, 377; bionomics and adult habits of, 377, 378; toxicity of insecticides and fungicides to, 378; massrearing of, 378.

Periplaneta americana, toxicity of extract from Bacillus larvae to, 332; mode of action of insecticides on, 63, 213; metabolism of DDT in, 64; used in tests of insecticides, 2, 117.

Perisierola, parasitising Oiketicus kirbyi in Costa Rica, 638.

Perisierola nigrifemur, parasitising Rhyacionia buoliana in Argentina, 144; redescription of, 144.

Perissopneumon tectonae, bionomics of, on guava and other trees in India, 174. peritana, Ptycholoma (Clepsis).

perla, Chrysopa.

perniciosi, Prospaltella.

perniciosus, Quadraspidiotus (Aspidiotus, Diaspidiotus).

pernyi, Antheraea.

Perotis malachitica, on apricot in Turkey, 495.

perpallidus, Aphelinus.

perparvulus, Bruchidius.

perrisi, Asolcus (Microphanurus).

Persea gratissima (see Avocado).

perseae, Saissetia (see S. nigra).

Persia (Iran), cereal pests in, 428, 429, 485, 653, 655, 656, 684; virus diseases of wheat in, 684; hibernation of Asolcus tumidus in, 654; strain of Trioxys pallidus introduced into California from, 151; survey of pests of stored products in, 236; beneficial insects in, 485, 653.

persicae, Myzus (Myzodes).

persimilis, Phytoseiulus.

Persimmon, wood-boring beetles damaging, in Japan, 450; Eulecanium corni on, in U.S.S.R., 13; Ceratitis capitata on, in Yugoslavia, 239.

Perspiration, as basis of attractiveness of textile stains to insects, 615.

Perthane (see Ethyl-DDD).

Peru, Phyllocoptruta oleivora on Citrus in, 560; cotton pests in, 559, 560; Blissus leucopterus in, 98; beneficial insects in, 560.

peruvianus, Dysdercus; Oligonychus (Paratetranychus).

Pestox 3 (see Schradan).

Petrobia latens, on wheat in China, 625. Petrol, trunk injection with, against larvae of Saperda carcharias, 601.

Petroleum Distillate, in aerosol mixture against *Bryobia praetiosa*, 419.

Petunia nana compacta, experiments with

groundnut virus and, 611.

Pexicopia malvella, on cotton and other plants in U.S.S.R., 113; effects of X-rays on fecundity and development of, 113; quarantine measures against, 113.

Phaedon cochleariae, effects of leaf texture on larval feeding by, 495.

Phaenobremia, predacious on Aphis pomi in Nova Scotia, 388.

Phaeogenes, parasitising Evagora starki in Canada, 586.

Phaeogenes hariolus, parasitic on Choristoneura pinus in Minnesota, 212; effect of tree location on degree of parasitism by, 212.

Phaeogenes invisor (parasitising Tortricids), in Czechoslovakia, 683; in U.S.S.R., 598.

Phaeogenes maculicornis, survey of data on, parasitising Choristoneura murinana in France and Germany, 168.

Phaeophytin, metabolism of chlorophyll to, in *Tetranychus telarius*, 108.

phaeosema, Chilo.

INDEX

Phanerotoma dentata, parasitising Cydia funebrana in U.S.S.R., 112.

phaselus, Tetranychus.

phaseoli, Melanagromyza (Agromyza).
Phaseolus, relation of maturity to injury
by Epilachna varivestis in species of,
414. (See Beans.)

Phaseolus acutifolius, susceptibility of strains of, to Epilachna varivestis, 414.

Phaseolus aureus, resistance to Epilachna varivestis in strains of, 413; experiments with Rivellia damaging root nodules of, 609; (stored), Callosobruchus maculatus infesting, 317, 500.

Phaseolus calcaratus, resistance to Epilachna varivestis in strains of, 413.

Phaseolus lathyroides, susceptibility to Epilachna varivestis of strains and hybrids of, 413, 414.

Phaseolus lunatus (see Lima Beans).

Phaseolus mungo, susceptibility to Epilachna varivestis of strains and hybrids of, 413, 414.

Phaseolus vulgaris (see Beans).

Pheidole megacephala, antagonistic to Oecophylla sp. in Zanzibar, 29.

Pheidole pallidula, predacious on Myiopardalis pardalina in Turkey, 352.

Pheidole punctulata, in S. Africa, 87; in Tanganyika, 87; antagonistic to Oecophylla sp. in Zanzibar, 29; effects of temperature on activity of, 87.

Phenacoccus aceris, new parasite of, in

Japan, **491**.

Phenacoccus hirsutus (see Maconellicoccus).

Phenacoccus pergandei (see P. aceris).

Phenkapton, in sprays against mites, 183, 439, 672; effects of, on beneficial mites, 439; persistence of, on apple leaves, 672; and Isolan, 183.

Phenobarbital (sodium salt), effects of, in mixture with rotenone against

Epilachna varivestis, 279. 2-Phenylethylisothiocyanate (in turnips),

toxicity of, to insects, 484.

Philadelphus coronarius, as primary foodplant of Aphis fabae in Bulgaria, 536.

Philaenus leucophthalmus (spumarius), on grasses and cereals in Britain, 321; on strawberry in Kentucky, 570; experiments with acids on decomposition of froth masses of, 321; insecticides against, 321, 570.

Philippines, Cicadulina bipunctella maize and rice in, 96.

Philopedon plagiatus (see Cneorrhinus).

phlei, Toxoptera graminum.

Phleum pratense, Oscinella frit on, in Britain, 430; pests of, in Canada, 162, 389, 390, 393, 585; pests of, in Finland, 41, 102; Rhopalosiphum padi on, in Wisconsin, 95; as food-plant of stages of Calligypona pellucida, 102.

phloeocoptes, Aceria (Eriophyes).

phloxiphaga, Heliothis.

Phlyctinus callosus, bionomics of, on vines in S. Africa, 650.

Phobetes, parasitising Oiketicus kirbyi in

Costa Rica, 638.

crassiuscula (parasitising **Phobocampe** Operophtera brumata), in Europe, 378; introduced into Canada, 378: in U.S.S.R., 190.

phoenicis, Brevipalpus.

Phoma, association of Sitona hispidulus with species of, on alsike clover in Oregon, 275.

Phonoctonus nigrofasciatus, number of Dysdercus fasciatus required by, for

development, 429.

Phorate, against aphids, 263, 278, 279, 333, 357, 401, 412, 467, 468, 474, 508, 510, 511, 556, 633, 645, 670; ineffective against Aphis pomi, 402; against other Hemiptera, 325, 395, 405, 409, 470, 474, 510, 628; against Coleoptera, 16, 219, 325, 396, 467, 511; ineffective against Fenusa pusilla, 19; toxicity of, to Diptera, 330, 512; against Lepidoptera, 19, 21, 273, 632, 642; ineffective against Pyralids, 333, 469; toxicity of, to Locusta migratoria, 172; against Thysanoptera, 151, 152, 212, 257, 352, **409**, **474**, **515**, **645**; against *Sminthurus* viridis, 34; against Tetranychids, 19, 264, 324, 515, 570, 628; against other mites, 645; toxicity of, to predator of Tetranychids, 627; toxicity of, to bees, 412; toxicity of, to man, 670; metabolism of, in insects, 64, 512, 513; anticholinesterase activity of, 395, 396, 409, 468, 470, 474, 510, 511, 515, 642, 645; in sprays, 34, 212, 213, 264, 273, 279, 324, 395, 412, 511, 570, 633, 642, 645; seed treatment with, 218, 224, 263, 277, 352, 357, 398, 401, 405, 467, 468, 470, 670; soil treatment with, 151, 219, 263, 273, 277, 325, 330, 333, 396, 405, 409, 468, 474, 508, 510, 511, 628, 642, 645; volatilisation of, from soils,

416; tests of systemic action and other

methods of application of, 16, 17, 19, 212, 257, 273, 278, 279, 395, 405, 467, 470, 556, 632, 633; factors affecting systemic activity of, 277, 508; tests of, against pests of stored products, 212, 213; phytotoxicity of, 152, 263, 277, 330, 352, 396, 401, 556; other effects of, on plants, 218, 409; effect of, on incidence of Pythium, 398; residues of, in chrysanthemum, 395; determination of residues and metabolites of, 424; labelled with radioactive isotopes, 224, 263; toxicity of DDT decreased by, 21; with other insecticides, 264; and fungicides, 264, 352; question of use of fertilisers with, 645; oxygen analogue of (see O,O-Diethyl S-Ethylthiomethyl Phosphorothioate).

Phormia regina, experiments on cuticular

penetration of BHC in, 43.

Phorocera agilis, parasitising Lymantria dispar in Yugoslavia, 370.

Phorocera obscura, parasitising larvae of Operophtera brumata in Europe, 378; introduced into Canada, 378.

Phorocera silvestris, parasitising Lymantria dispar in Massachusetts, 400; persistence of, in sprayed areas, 400.

Phorodon menthae, on Mentha spp. in Washington, 324; phosphorus com-

pounds against, 324.

Phosdrin (see Mevinphos). Phosphamidon, against aphids, 216, 249, 279, 333, 415, 668; ineffective against Myzus persicae, 637; against Empoasca, 325; against Coleoptera, 325, 511; against Diprion similis, 418, 419; against Lepidoptera, 9, 222, 333, 351, 421, 434, 466, 513, 637, 663; against Trypetids, 237, 268, 668; toxicity of, to Tetranychus telarius, 614; effects of, on parasite of Eriosoma lanigerum, 313; toxicity of, to bees, 412; in dusts, 637; in sprays, 9, 216, 222, 237, 249, 268, 279, 313, 325, 333, 351, 415, 421, 434, 466, 511, 637, 663, systemic activity and other 668; methods of application of, 249, 279, 418, 419, 434, 663; residues of, in olive, 238; activity of endrin enhanced by, 351; and schradan, 511.

Phosphine, as fumigant (see Aluminium

Phosphide).

Phosphoric Acid, use of, in labelling Pectinophora gossypiella with 32P, 338; as metabolite of dimethoate, 563.

Phosphorothioic Acid, tests of esters of, against Tetranychus telarius, 478.

Phosphorus (as plant nutrient), effect of, on translocation of dimethoate, 473. Phosphorus, Radioactive, used in studies of insect behaviour, 29, 45, 265, 338, 478; methods for labelling Anthonomus grandis with, 471; effects of, on stages of A. grandis, 472; insecticides labelled with, 224, 467, 472, 475, 497, 562.

Phostex (mixture of bis(dialkoxyphosphinothioyl) disulphides), toxicity of: to bees, 412; to Rhagoletis pomonella,

268.

Phostoxin, 483.

Photoperiod, effects of: on egg development and feeding activity in Coleoptera, 205; on pupation by Anthrenus verbasci, 249; on oviposition by Panonychus ulmi, 361; on diapause in insects, 4, 6, 94, 204, 276, 319, 451, 465, 488, 489, 522, 542, 634, 665, 679; on diapause in strains of Tetranychus telarius, 688; on development of parasites in Lepidoptera, 379; on transmission of virus disease by aphids, 610; reviews of, in relation to insects and mites, 90, 423.

Photosynthesis, influence of feeding by aphids on mechanism of, 44, 45.

Phryxe (see Zenillia).

Phthorimaea, type and status of, 235.

Phthorimaea ocellatella (see Scrobipalpa). Phthorimaea operculella, in British Columbia, 529; in Bulgaria, 236; in Czechoslovakia, 236; in India, 448, 499; in Sweden, 503; on potato, 236, 529; infesting stored potatoes, 448, 499; on tobacco, 236, 503; other foodplants of, 236; bionomics of, 236; measures against, 448, 499, 503; geographical origin of, 236; characters distinguishing Scrobipalpa spp. from, 236.

Phygadeuon pegomyiae, parasitising Pego-

myia betae in Germany, 442.

Phyllocnistis, parasite of, on Citrus in Aden, 80.

Phyllocnistis citrella, characteristics and

world distribution of, 507.

Phyllocoptruta (Phyllocoptes) oleivora (on Citrus), in S. Africa, 83; in Brazil, 556, 644; in Florida, 409; in Gaza Strip, 97; in Israel, 614, 684; in Peru, 560; sex ratio of, 614; relation of, to leprosis of Citrus, 556; measures against, 409, 410, 556, 560, 644, 645; di(p-chlorophenyl)-trifluoromethylcarbinol ineffective against, 684; favoured by DDT, 560; question of resistance to zineb in strains of, 409.

Phyllodecta tibialis, dusts against, on

willows in Poland, 40.

Phyllopertha horticola (see Anomala). Phyllophaga (see Lachnosterna).

Phyllospilum ipomoeae, 133.

Phyllotreta (on crucifers), in Finland, 103, 674; in France, 373; natural enemies of, 373; insecticides against, 104, 674.

Phyllotreta atra (on crucifers), in Austria, 184; in Yugoslavia, 181; Bacillus thuringiensis ineffective against, 184.

Phyllotreta cruciferae (on crucifers), in Canada, 529; in India, 495; in New York, 69; in Yugoslavia, 181; bionomics of, 495; relation of diversity of plant species to populations of, 69.

Phyllotreta nemorum, on radish in Austria, 184; characteristics and world distribution of, 507; Bacillus thuringiensis

ineffective against, 184.

Phyllotreta pusilla, systemic insecticides against, on potato in Texas, 510.

Phyllotreta vittata, relation of diversity of plant species to populations of, in New York, 69.

Phyllotreta vittula, on cereals in Yugo-

slavia, 181.

Phylloxera vitifoliae (vastatrix) (on vines), in France, 84, 667, 668; transmitting mottling virus, 303; insecticides against, 84, 572, 667, 668.

Physalis floridana, preference of Myzus persicae for excised or virus-infected leaves of, 584; experimental transmission of potato viruses to, by M.

persicae, 689, 691.

Physalis minima, Heliothis assulta on, in Queensland, 589.

Physalis peruviana, Heliothis assulta on, in Queensland, 589.

Physcus, parasitising Aonidiella aurantii in Burma, 266; introduced into California, 266; identity of species of, from Formosa, 491; bionomics of, 266; oviposition of, on other Coccids, 266; method of rearing, 267.

Physcus confusus, sp.n., parasitising Aoni-

diella in Burma, 491.

Physcus debachi, sp.n., parasitising Comperiella bifasciata associated with Aonidiella aurantii in Burma and Siam, 491.

Physcus flaviceps, new species of Physcus misidentified as, 491.

Physcus flavicornis, sp.n., parasitising Pseudaonidia duplex in China, 491.

Physcus fulvus, sp.n., in China and Formosa, 491; question of, parasitising Unaspis yanonensis, 491; previously known as Physcus B, 491; misidentified as P. flaviceps, 491.

Physcus reticulatus, sp.n., parasitising Aonidiella aurantii in India and Paki-

stan, 491.

Physostigmine Salicylate, effects of, in mixture with rotenone against *Epilachna* varivestis, 279.

Phytagromyza populicola, on poplar in Quebec, 529.

Phytocoris obscurus, on cotton in Turkey, 352.

Phytodecta fornicata, on lucerne Bulgaria, 60.

Phytodecta | olivaceus, on broom England, 30; precipitin-test identification of predators of, 30.

Phytodietus segmentator, parasitising Tortricids in Czechoslovakia, 683.

Phytometra ni (see Trichoplusia).

Phytomyza, species of, on chick-pea in Turkey, 351.

Phytomyza atricornis, question of occurrence of, on chick-pea in Turkey, 351.

Phytomyza lateralis, on chicory, carrot and other plants in Holland, 673; comparison of injury by Psila rosae and, 674; insecticides against, 674.

Phytomyza rufipes, bionomics of, crucifers in Denmark, 139; characters distinguishing damage by, 139; sprays against, 139.

Phytonomus (see Hypera).

Phytophaga destructor (see Mayetiola).

Phytophthora infestans, 569.

Phytoptus avellanae, on hazel in Turkey, 82, 83; effects of sulphur dusts against, on populations of Gypsonoma dealbana, 83.

Phytoptus ribis (see Cecidophyopsis). Phytoseiulus persimilis, in Chile, 125; introduced into Ontario, 125; predacious on Tetranychus telarius, 125.

Pianos, fumigation of, against termites, 296.

Picea abies, Ips typographus on, in France, 606; pests of, in Italy, 136, 615; Formica spp. in forests of, 615.

Picea alba, Ips curvidens on, in Czechoslovakia, 283.

Picea engelmanni, Dendroctonus engelmanni on, in Colorado, 150.

Picea excelsa (see P. abies).

Picea glauca, pests of, in Canada, 25, 157, 457; Choristoneura pinus on, in Minnesota, 474.

Picea sitchensis, Rhopalosiphum abietinum on, in Germany, 105; relation of nitrogen content of, to feeding habits of R, abjetinum, 105.

piceae, Chermes (Adelges, Dreyfusia). piceus, Attagenus (see A. megatoma).

pici, Megatoma.

picipes, Aphidius (see A. avenae); Omophlus.

picta, Ora.

pictipes, Aegeria (Synanthedon). pieridis, Apanteles.

Pieris brassicae, in Austria, 184; in Poland, 545; in U.S.S.R., 114; on cabbage, 114, 184, 545; bionomics and adult habits of, 51, 290, 291, 488, 489; factors affecting induction of diapause in, 678; colour preferences of, 52; parasites of, 87, 114, 115, 381, 678; characteristics and world distribution of, 507; Bacillus thuringiensis unsatisfactory against larvae of, 184; hemp guard-rows ineffective against, 545.

Pieris rapae (on crucifers), in Canada, 161, 163, 385, 525; in Japan, 78, 449, 450; in New Zealand, 75; in U.S.S.R., 114; in U.S.A., 172, 222, 269, 401, 404; infestation patterns of, 385, 449; relation of populations of, to other cabbage pests, 172; effects of crowding on larvae of, 450; tests of Bacillus thuringiensis against, 75, 269, 525; diseases of, 78, 401; natural enemies of, 78, 114, 115, 401; tests and uses of nematode against, 163; other mortality factors of, 78; insecticides against, 161, 222, 269, 525.

Piesma quadratum, characteristics and world distribution of, 507.

Piezotrachelus varium (see Apion).

Pigs, insecticide residues in tissues of, 384.

Pike, toxicity of insecticides to, 304.

Pikonema alaskensis (see Pachynematus). pilleriana, Sparganothis.

Philophorus uhleri, predacious on larvae of Neodiprion swainei in Quebec, 73. pilosus, Paratetranychus (see Panonychus

ulmi).

Pimelia mongeneti, insecticides against, on cotton in Greece, 352.

Pimento, Acroleucus coxalis on, in Brazil, 356; Zonosemata electa on, in Ontario, 529.

Pimpla pedalis, parasitising Thymelicus lineola in Ontario, 162.

Pimpla spuria, parasitising Operophtera brumata in U.S.S.R., 190.

Pimpla tomyris, parasitising Rhyacionia buoliana in Argentina, 230.

Pimpla turionellae (parasitising Lepidoptera), in France and Germany, 168;

established in Ontario, 73, 74; survey

of data on, 168.

Pine, Nudaurelia cytherea capensis on, in S. Africa, 425; Rhyacionia buoliana on, in Argentina, 144, 230; R. buoliana on, in Belgium, 606; Bupalus piniarius on, in Britain, 550; attacked by Coptotermes niger in British Honduras, 76; pests of, in Canda, 24, 25, 73, 74, 121, 122, 123, 158, 163, 206, 260, 261, 387,

391, 392, 457, 458, 459, 474, 529, 586; Dioryctria nivaliensis on, in Canary Islands, 659; Dendrolimus punctatus on, in China, 501, 623, 624; Exoteleia dodecella on, in Czechoslovakia, 246; Hylobius abietis on, in Finland, 194; pests of, in Germany, 10, 376, 683; pests of, in Italy, 615; D. superans and secondary pests on, in Mongolia, 116; Icerya purchasi infesting seedlings of, in Puerto Rico, 257; pests of, in S. Rhodesia, 81; pests of, in Spain, 659, 660; sawflies on, in Sweden, 426; pests of, in U.S.S.R., 111, 112, 115, 116, 344, 598; pests of, in U.S.A., 19, 20, 23, 24, 73, 74, 121, 124, 125, 129, 153, 212, 221, 336, 340, 386, 410, 413, 418, 458, 460, 462, 467, 474, 481, 495, 632, 642; pests of, in Yugoslavia, 31, 536; systematics of needle-mining Lepidoptera on, in N. America and Europe, 121; Galleria mellonella on, 130; factors affecting susceptibility of, to insects, 340, 598, 599, 642; vigour of, related to damage by R. buoliana, 481; attraction of Trypodendron lineatum to logs of, 459; difficulty of establishment of Formica spp. in forests of, 615; fumigant toxicity of resins of, to Dendroctonus spp., 223; effects of BHC on insect fauna of, 683; DDT treatment of seedlings of, 194; systemic activity of insecticides in, 19, 418, 419, 633; Coleoptera infesting timber of, 112, 647, 648; penetration of timber of, by wood preservatives, 647, 648; use of timber of, in tests of treatments against termites, 649.

Pine, Black (see Pinus nigra).

Pine, Corsican (see Pinus nigra poiretiana).

Pine, Jack, (see Pinus banksiana).

Pine, Japanese Red (see Pinus densiflora).

Pine, Jeffrey (see Pinus jeffreyi).

Pine, Lodgepole (see Pinus contorta).

Pine, Pitch (see Pinus rigida).

Pine, Ponderosa (see Pinus ponderosa).

Pine, Red (see Pinus resinosa).

Pine, Scots (see Pinus sylvestris).

Pine, Shortleaf (see *Pinus echinata*).

Pine, Sugar (see Pinus lambertiana).

Pine, Western White (see *Pinus monti-cola*).

Pine, Weymouth (see Pinus strobus).

Pine, White (see Pinus strobus).

Pine Beetle, Southern (see *Dendroctonus* frontalis).

Pine Caterpillar (see *Dendrolimus puncta-*

Pine Oil, in emulsion sprays with parathion against Hemiptera, 667, 668.

Pine Shoot Moth, European (see Rhy-acionia buoliana).

Pineapple, *Dysmicoccus brevipes* on, in Brazil, 557; list of pests of, in South-East Asia and Pacific area, 423; toxicity of insecticides to, 557.

Pinene, Chlorinated, in sprays against

Eriophyes pyri, 191. Pineus (see Chermes).

pini, Chermes (Pineus); Dendrolimus; Diprion; Elachertus; Eurytoma; Ips; Trisetacus.

piniarius, Bupalus.

pinicolana, Rhyacionia (Evetria).

pinifolii, Aethecerus.

piniphilus, Gnophothrips.

piniradiatae, Schizolachnus.

Pinus banksiana, pests of, in Canada, 73, 122, 123, 163, 260, 261, 459, 474, 529; Choristoneura pinus on, in U.S.A., 212, 467, 474.

Pinus caribaea, Lechriolepis nephopyropa

on, in S. Rhodesia, 81.

Pinus cembra, Dendrolimus superans and secondary pests on, in Mongolia, 116.

Pinus contorta, Scolytus spp. on, in British Columbia, 458; Evagora milleri on, in California, 124, 221.

Pinus contorta latifolia, Rhyacionia buoliana on, in Belgium, 606.

Pinus densiflora, Dioryctria zimmermanni on, in Illinois, 153.

Pinus echinata, pests of, in U.S.A., 20, 336, 462.

Pinus elliottii densa, new sawfly on, in Florida, 462.

Pinus insignis (see P. radiata).

Pinus jeffreyi, pests of, in U.S.A., 23, 24, 386; fumigant toxicity of resins of, to Dendroctonus spp., 223.

Pinus lambertiana, Neodiprion spp. on, in

California, 125.

Pinus massoniana, Dendrolimus punctatus on, in China, 623, 624.

Pinus monticola, pests of, in British Columbia, 458, 459.

Pinus nigra, Thaumetopoea pityocampa on, in Yugoslavia, 536.

Pinus nigra poiretiana, Dioryctria zimmermanni on, in Illinois, 153.

Pinus patula, pests of, in S. Rhodesia, 81. Pinus ponderosa, pests of, in British Columbia, 458, 529; pests of, in U.S.A., 23, 24, 340, 386, 460; fumigant toxicity of resins of, to Dendroctonus spp., 223; relation of water supply and resin exudation to infestation of, by bark-beetles, 340.

Pinus radiata, Nudaurelia cytherea capensis on, in S. Africa, 425; Ips spp.

infesting, in California, 386; Dioryctria nivaliensis on, in Canary Islands, 659. Pinus resinosa, pests of, in Ontario, 24, 122, 261, 392, 458; pests of, in U.S.A., 19, 20, 129, 410, 413, 474, 481, 632; vigour of, related to damage by Rhyacionia buoliana, 481; activity of systemic insecticides in, 19, 633.

Pinus rigida, pests of, in U.S.A., 20, 336. Pinus strobus, pests of, in Canada, 122, 123, 206, 529; Acantholyda erythrocephala on, in Germany, 10; pests of, in U.S.A., 418, 474; Conophthorus coniperda on, 261; effects of growth of, on populations of Pissodes strobi, 206, 207; systemic activity of insecticides in, 418, 419.

Pinus sylvestris, Rhyacionia buoliana on, in Belgium, 606; Hylobius abietis on, in Finland, 194; pests of, in Ontario, 24, 261, 457, 458; Lepidoptera on, in

U.S.A., 153, 474; DDT treatment of seedlings of, 194.

Pinus taeda, Lechriolepis nephopyropa on,

in S. Rhodesia, 81.

Pinus thunbergi, Rhyacionia frustrana on, in New York, 19; activity of systemic insecticides in, 19.

Pinus virginiana, Neodiprion pratti pratti on, in U.S.A., 336.

pinus, Choristoneura.

Pionea ferrugalis (see Udea).

Piophila casei, infesting cheese-storage premises in Switzerland, 100; Necrobia rufipes predacious on, 436; Dimetilan-

treated strips against, 100.

Piperonyl Butoxide, as synergist: with carbamates, 148, 476, 508, 516; action of, on inhibition of cholinesterase by carbamate compounds, 635; with chlorinated hydrocarbons, 587, 687; with dinoseb, 687; toxicity of phosphorus compounds reduced by, 63; with pyrethrins, 2, 3, 4, 63, 82, 128, 225, 270, 402, 500, 560, 587, 591, 687, 688; mode of action of pyrethrins with, 688; chromatographic separation of pyrethrum and, in mixtures, 562.

pipiens, Culex; Syritta.

Pirazinon, toxicity of, to Rhagoletis

pomonella, 268.

pisi, Bruchus (see B. pisorum); Contarinia; Macrosiphum (Acyrthosiphon, Aphis) (see M. pisum); Melanagromyza.

pisivorus, Aphidius; Kakothrips.

pisorum, Bruchus.

Pissodes notatus, on pine in Spain, 660; characteristics and world distribution of, 507.

Pissodes strobi, on white pine in Ontario, **206**; factors affecting populations of, **206**.

Pissodes validirostris, on pines in U.S.S.R., 116.

Pistacia vera (Pistachio), pests of, in Turkey, 81, 609; (stored nuts), Plodia interpunctella in, 337, 609; use of, in rearing P. interpunctella, 338.

pistaciae, Eumegastigmus; Lepidosaphes. pisum, Macrosiphum (Acyrthosiphon).

Pitangus sulphuratus, introduced into Bermuda from Trinidad against harmful lizards, 229.

Pitch Pine (see Pinus rigida).
pityocampa, Thaumetopoea.
Pityokteines curvidens (see Ips).
plagiatus, Cneorrhinus (Philopedon).

Plagiognathus medicagus, sprays against, on lucerne in Canada, 72.

planaris, Corythaica (see C. passiflorae).

Plane (see Platanus acerifolia).

Planococcus citri, in S. Africa, 298; in Israel, 612, 614; on greenhouse plants in Oklahoma, 339; on Citrus, 612, 614; on pomegranate, 614; on vines, 298, 614; natural enemies and hyperparasites of, 614; insecticides against, 298, 612. Planococcus dioscoreae, sp.n., on yams in

New Guinea, 234.

plantaginea, Anuraphis (Dysaphis). Plantago (Plantain), migration of Anura-

Plantago (Plantain), migration of Anuraphis plantaginea to, in Holland, 673.

Plant Protection, bibliography of, for 1954, 248, 424.

Plasmopara viticola, on vines in Bulgaria, 538; combination of insecticides with measures against, 538.

Plaster of Paris, in substrates for mites, 518; use of, in artificial nesting sites for bees, 411; use of, in determination of insecticide residues in soils, 339.

Plastic Wood, protection of timber with,

against termites, 486.

plastographus, Ips; Tarsonemoides.

Plastophora rufa, predacious on Eulecanium spp. in France, 683.

platani, Lithocolletis.

Platanus acerifolia, Lithocolletis platani on, in Yugoslavia, 31.

platensis, Sitophilus oryzae (See S. zeamais).

Platyedra gossypiella (see Pectinophora). Platygaster, parasitising Contarinia oregonensis in British Columbia, 386.

Platygaster americana, 386.

Platygaster manto, biological control of Agevillea abietis with, in Germany, 375; effects of insecticides on populations of, 375.

Platymyia mitis (see Zenillia).

Platyparea poeciloptera, characteristics and world distribution of, 507.

platyptera, Eurytoma.

Platypus cylindrus, infesting oak logs in Yugoslavia, 31; treatment of timber against, 31.

plebeja, Euscelis.

Pleonomus canaliculatus, on wheat in China, 625.

Plesiocoris rugicollis, characteristics and world distribution of, 507.

plesuis, Epigonatopus.

pleurostigma, Ceutorhynchus. Pleurotropis (see Pediobius).

Plodia interpunctella, in Japan, 451: in Turkey, 609; infesting pistachio nuts, 337, 609; factors affecting larval diapause in, 451; method of rearing, 337; toxicity of Zectran to, 516.

plotnikovi, Eurytoma.

Plum, Eriophyes padi on, in Austria, 85; Panonychus ulmi on, in Britain, 671; Hyponomeuta padellus padellus on, in Bulgaria, 110; Pristiphora monogyniae on, in Finland, 103; Cydia molesta on, in Ontario, 163; Quadraspidiotus perniciosus on, in Pakistan, 170; pests of, in Poland, 61, 188, 189, 544, 677, 678; Hoplocampa minuta on, in Rumania, 56; Tetranychus viennensis on, in Turkey, 177; pests of, in U.S.S.R., 13, 15, 112, 242, 360; pests of, in U.S.A., 23, 147, 470; pests of, in Yugoslavia, 32, 138, 240, 241; influence of Panonychus ulmi on growth of, 671; fruit of, tainted by BHC, 544. Plum, Cherry, Hyponomeuta padellus on,

in Bulgaria, 110.

Plum, Japanese, Pseudococcus comstocki on, in Ontario, 164.

Plum Curculio (see Conotrachelus nenuphar).

Plusia argentifera, sprays against, on tobacco in Queensland, 503.

Plusia californica (in U.S.A.), on Mentha spp., 324; parasitised by Hyposoter exiguae, 66; DDT against, 324.

Plusia chalcites, on beans in New Zealand, 75; tests of Bacillus thuringiensis against, 75.

Plusia ni (see Trichoplusia).

Plusia nigrisigna (see Autographa).

Plutella maculipennis (on crucifers), in Belgium, 430; in Britain, 197, 198, 430; in Canada, 161, 211, 525; in Denmark, 198, 430; in Estonia, 430; in Finland, 198, 430; in France, 430; in Germany, 430; in Holland, 430; in New Zealand, 75; in Norway, 430; in Sweden, 198, 430; in U.S.S.R., 87, 198, 244, 245, 430;

in U.S.A., 172, 401; relation of populations of, to other cabbage pests, 172; bionomics and migration of, 198; sampling method for populations of, 211; fungus infesting, 198; natural enemies of, 87, 245, 401; tests of Bacillus thuringiensis against, 75, 525; insecticides against, 161, 198, 430, 525.

plutella, Aphelinoidea. plutellae, Microplitis.

plutus, Crepidodera (Chalcoides).

pluviale, Malacosoma.

Pnigalio longulus, parasitising Dacus oleae

in Italy, 308, 438, 532.

Pnigalio maculipes, parasitising stages of Bucculatrix ainsliella in Michigan, 327. Pnigalio mediterraneus, parasitising Dacus oleae in France, 435.

Poa, mites and silver top of, in Alberta, 156; stem-boring Diptera on, in Eng-

land, 487, 488.

Poa pratensis, relation of mites and thrips to silver top of, in Alberta, 156; Rhopalosiphum padi on, in Wisconsin,

Poa secunda, relation of mites and thrips to silver top of, in Alberta, 156.

poae, Myzus (Rhopalosiphum).

Podacanthus wilkinsoni, on Eucalyptus in Australia, 32, 33, 231, 232; bionomics of, 32, 33; parasites of, 32; other factors affecting populations of, 33, 231, 232; effects of sprays on, 33.

Podagrica malvae semirufa, on Hibiscus esculentus in Yugoslavia, 181.

Podagrica pallida, on cotton in Sudan, 52; effects of sowing date on populations of, 52, 53.

Podagrica puncticollis, on cotton in Sudan, 52; effects of sowing date on populations of, 52, 53.

poeciloptera, Platyparea.

poephaga, Sesamia.

Poinciana regia, Pericyma cruegeri on, in Papua, 300.

Poinsettia (see Euphorbia pulcherrima). Poland, pests of crucifers in, 61, 545, 599; pests of potato in, 366, 544, 681, 683; virus diseases of potato in, 544; Thysanoptera on flax in, 366; Ostrinia nubilalis on maize in, 365; orchard pests in, 61, 188, 189, 364, 365, 544, 676, 677, 678; Steneotarsonemus fragariae on strawberry in, 545; Ptilinus fuscus on willows and poplars in, 189, 190; mites infesting stored seed of flax and hemp in, 364; Anobiids infesting timber in, 142; Melolontha in, 678; natural enemies of pests in, 189, 364, 365, 379, 545, 677, 683.

Polia dissimilis (see Hadena suasa). Polia oleracea (see Diataraxia).

Poliaspis argentosus, on Leptospermum in New Zealand, 588.

polita, Microgaster.

Pollination, and bees, 234, 411, 565; of carrots, 341; cage for insects studies of, 620.

Polyblastus flavicauda, parasitising Pristiphora abietina in Germany, 682; effects of BHC against Melolontha on populations of, 682.

Polybutene, in dormant spray against Psylla pyricola, 412.

Polychrosis botrana (see Lobesia).

polyctena, Formica.

Polynema eutettixi, parasitising leafhoppers in California, 630.

Polyphylla alba, injuring pines in U.S.S.R.,

Polyphylla decemlineata, on Mentha spp. in Washington, 324; soil treatments against larvae of, 324.

Polysulphide, 288.

Polytetrafluorethylene, use of, as barrier against mites, 518.

polytoma, Gilpinia.

Polyvinyl Acetate, efficiency of endrin sprays not increased by addition of,

Polyvinyl Pyrrolidone, ineffective in mixture with iodine against larvae of Agrotis ypsilon, 211.

Pomarsol, 381.

Pomegranate, insects reared from fruits of, in S. Africa, 531; Planococcus citri on, in Israel, 614.

pometaria, Alsophila.

pomi, Aphis.

pomonella, Cydia (Carpocapsa, Laspeyresia); Rhagoletis.

pomorum, Anthonomus.

Ponderosa Pine (see Pinus ponderosa).

ponderosae, Dendroctonus.

Popillia japonica, in U.S.A., 418, 577; factors affecting attraction of, to Ginkgo biloba and cherry, 577; apples as food for, 418; dispersal of, by aircraft, 418; survival of, in nitrogen atmospheres, 416; insecticides against, 226, 418; question of resistance to DDT in strains of, 418.

Poplar (Populus), Hyponomeuta gigas on, in Canary Islands, 659; Saperda spp. on, in Czechoslovakia, 196, 600; Cryptorhynchus lapathi on, in Hungary, 307; Tortrix sp. on, in New Zealand, 75; Ptilinus fuscus on, in Poland, 190; pests of, in Quebec, 392, 529; pests of, in Spain, 300, 353, 355, 659; flea-beetles on, in Yugoslavia, 181; varieties of, in relation to S. populnea, 196. (See also Populus spp.)

Poplar, Black (see *Populus nigra*).

Poplar, Canadian (see Populus deltoides). Poplar, Lombardy (see *Populus italica*).

Poplar, White (see *Populus alba*). Poppy (see Papaver somniferum).

Population Studies, of insects, 33, 656; sampling methods for, 33, 69, 83, 84, 211: review of mathematics in, 90.

populicola, Phytagromyza.

populnea, Saperda. Populus (see Poplar).

Populus alba, Stilpnotia salicis on, in British Columbia, 126.

Populus balsamifera, Trichiocampus viminalis on, in Quebec, 391.

Populus deltoides, Mordwilkoja vagabunda on, in Minnesota, 552, 619; Saperda carcharias on, in U.S.S.R., 114.

grandidentata, Trichiocampus viminalis on, in Quebec, 391.

Populus italica, Stilpnotia salicis on, in British Columbia, 126.

Populus monilifera, Saperda populnea on, in Czechoslovakia, 600; larvae of S. populnea killed by wound tissue of,

Populus nigra, Trichiocampus viminalis on, in Quebec, 392; Saperda carcharias on, in U.S.S.R., 114.

Populus tremuloides, pests of, in Canada, 126, 392.

Populus trichocarpa, Stilpnotia salicis on, in British Columbia, 126.

Porizon (see Tersilochus). Porthetria (see Lymantria).

portola, Draeculacephala.

Porto Rico (see Puerto Rico).

Portugal, insects associated with elm in, 298; Hyalopterus amygdali on peach in, 617.

postica, Hypera (see H. variabilis).

(Austrotortrix, postvittana, Epiphyas Tortrix).

Potamyia flava, nuisance value of, in Iowa, 262.

Potassium, effects of, in insects, 62.

Potassium Bifluoride, treatment of timber with, against Hylotrupes bajulus, 253.

Potassium Chloride, attraction of insects to, 635.

otassium Cyanide, susceptibility of Galleria mellonella to virus disease Potassium unaffected by, 427; in experiments with Tettigonia viridissima and tobacco mosaic virus, 443.

Potassium Dichromate, as constituent

of Celcure (q.v.), 485.

Potassium Myronate, oviposition by Pieris brassicae stimulated by, 290.

Potassium Nitrate, susceptibility of Galleria mellonella to virus disease un-

affected by, 427.

Potato, Ptericoptus sp. on, in Argentina, 552; Leptinotarsa decemlineata on, in Belgium, 681, 683; pests of, in Brazil, 554, 556; pests of, in Britain, 320, 321; pests of, in Bulgaria, 236, 346, 366, 539; pests of, in Canada, 73, 207, 377, 424, 526, 529, 633; pests of, in Czechoslovakia, 236, 247, 681; virus diseases of, in Finland, 101; pests of, in Germany, 252, 255, 377, 440, 543, 546, 547, 680, 684; pests of, in Hungary, 305, 307; Epilachna vigintioctopunctata on, in India, 497; Myzus persicae on, in Israel, 614; L. decemlineata on, in Italy, 533, 681; L. decemlineata on, in Majorca, 681; pests of, in Poland, 366, 544, 681, 683; L. decemlineata on, in Rumania, 681; pests of, in U.S.S.R., 15, 543, 681; pests of, in U.S.A., 325, 333, 377, 412, 460, 473, 474, 510, 511, 569, 570, 627; pests of, in Yugoslavia, 240, 241, 681; distribution of L. decemlineata on, in Europe, 681; development of Agrotis ypsilon on, 346; other insects on, 377, 378, 500, 689; aphids and virus diseases of, 240, 241, 250, 255, 320, 321, 544, 546, 547, 614, 680, 688, 689, 690, 691; experiments on transmission of viruses of, by Tettigonia viridissima, 443; Phytophthora infestans on, 569; treatment of, with nematode against L. decemlineata, 207; systemic activity of insecticides in, 325, 510, 511; question of heptachlor residues in, 479; toxicity of insecticides to, 556, 633; detection of growth inhibitors in, 22; effect of spray machinery on yields of, 333; (tubers), Phthorimaea operculella infesting, 448, 499; Coccids on, 119, 437; use of, in rearing Coccids, 229; treatment of, with insecticides, 120, 448, 499; effects of azinphos-methyl on setting of, 397. (See also Solanum polyadenium.)

Potato Wireworm, Southern (see Cono-

derus falli).

Potentilla, Capitophorus spp. on species of, in Canada and U.S.A., 23.

potentillae, Capitophorus (Pentatrichopus). P.P. 175 (see Menazon).

P.P.C. 3 (see o-Isopropylphenyl Methylcarbamate).

praefica, Prodenia.

praetiosa, auct., Bryobia (see B. rubrioculus).

praetiosa Koch, Bryobia.

Praon, parasitising Myzus persicae in New York, 401.

Praon occidentalis, parasitising Macrosiphum pisum in California, 66.

Praon simulans, parasitising Macrosiphum pisum in California, 66.

Praon unicus, parasitising Chromaphis juglandicola in California, 151.

pratensis, auct., Formica (see F. nigricans);

pratensoides, Formica minor (see F. nigricans)

pratti, Neodiprion.

Prays oleellus (on olive), in Italy, 533; in Spain, 99; in Yugoslavia, 609; insecticides against, 99, 533, 609.

precationis, Autographa.

Premnotrypes, characteristics and world distribution of species of, 507.

Prince Edward Island, pests of conifers in, 121, 529.

Priobium carpini (see Trypopitys).

Pristiphora abietina (on spruce in Germany), natural enemies of, 141, 142, 682; other factors affecting development and populations of, 141, 253, 299.

Pristiphora erichsonii (on larch), in Canada, 124, 125, 210, 341, 456; in U.S.A., 210; influence of temperature on development of, 125; natural enemies of, 125, 210; fungi infesting, 210; pathogenicity of Bacillus spp. to, 283; strain of B. thuringiensis thuringiensis isolated from, 341; comparison of survey methods for, 456.

Pristiphora monogyniae, on plum in

Finland, 103.

Proceras venosatus, on maize in China, 502; cold-hardiness of, 502.

processionea, Thaumetopoea.

proclia, Aphytis.

Procris pruni (see Adscita).

Prodenia litura, in Egypt, 149; in Israel, 351; in Sweden, 503; in Turkey, 82; on Citrus, 82; on cotton, 149, 351; on 503; characteristics tobacco, world distribution of, 507; effects of larval crowding on colour and size of, 149; insecticides against, 351, 503; resistance to endrin in strains of. 351.

Prodenia ornithogalli, parasitised by Hyposoter exiguae in California, 66.

Prodenia praefica, parasitised by Hyposoter exiguae in California, 66.

prognathana, Spilonota (Tmetocera).

Prolan, metabolism of, in Musca domestica, 64; mixture of Bulan and (see Dilan).

Prolasioptera berlesiana, destroying Dacus: oleae in Italy, 308, 438, 532.

prolixus, Rhodnius.

Promecotheca papuana, measures against, on coconut in New Britain, 166, 167; natural enemies of, 166; survey of data on, 166.

Pronematus, saprophagous species of, on soy bean in Delaware, 414.

Prontaspis yanonensis (see Unaspis).

propinqua, Calligypona.

Propyl Isome, as synergist with pyrethrins, 63; toxicity of phosphorus compounds reduced by, 63.

Prospaltella, species of Encarsia referred

to as, in Peru, 560.

Prospaltella berlesei, parasitising Pseudaulacaspis pentagona in Italy, 616.

Prospaltella fasciata (parasitising Quadraspidiotus perniciosus), bionomics of, in

France, 380; in U.S.A., 379.

Prospaltella perniciosi (parasitising Quadraspidiotus perniciosus), in China, 664; in France, 379, 380, 433, 435, 681; established in Germany, 681; in Ontario, 664; established in Pakistan, 170; established in Switzerland, 664, 681; in U.S.S.R., 664; in U.S.A., 379, 664; bionomics of, 379, 380; Aphytis proclia in competition with, 380; effects of sprays against Q. perniciosus on populations of, 433, 434, 435; spray schedule developed for use with, 664; mass breeding of, 681.

Protein Hydrolysate, in baits for Trypetids, 439, 533, 590; and malathion,

533.

proteus, Omophlus.

Protoparce quinquemaculata, parasitised by Hyposoter exiguae in California, 66. Protoparce sexta, in S. Carolina, 465, 635; on tobacco, 465; experiments on food-plant selection by, 552; recovery of female sex attractant from, 635;

baits for, 465; measures against, 465. Protozoa, infecting insects, 89, 105, 187, 283, 343, 428, 511, 582, 583, 638, 658; hot-water treatment of eggs of *Ostrinia nubilalis* against, 343; (Microsporidia), book on, 89.

Proxenus mindara, characters of stages of, damaging strawberries in California,

563.

pruinosa, Metcalfa (Ormenis); Sericesthis. pruinosum, Eulecanium (Lecanium).

pruni, Hyalopterus; Adscita (Procris). prunifoliae, Rhopalosiphum (see R. padi). Prunus, Hyponomeuta padellus on species of, in Bulgaria, 110; association of Scaphytopius acutus with species of, in New York, 23; Caliroa limacina on, in Quebec, 26.

Prunus cerasus (see Cherry).

Prunus japonica, Cydia funebrana on, in U.S.S.R., 112.

Prunus mahaleb, Neurotoma nemoralis on, in Bulgaria, 541.

Prunus maritima, Psorosina hammondi on, in Canada, 529.

Prunus nigra, primary food-plant of Myzus persicae in Maine, 461; as winter food-plant of Rhopalosiphum padi in New Brunswick, 585; experiments with R. fitchii on, 585.

Prunus padus, Hyponomeuta euonymellus

on, in Finland, 194.

Prunus pensylvanica, as winter food-plant of Rhopalosiphum padi in New Brunswick, 585; experiments with R. fitchii on, 585.

Prunus virginiana, pests of, in Canada, 343, 585; pests of, in New York, 19, 23; experiments with Rhopalosiphum fitchii on, 585; activity of systemic insecticides in, 19.

Psallus, species of, predacious on apple pests in Nova Scotia, 159; effects of sprays on populations of, 159.

Psallus ambiguus, predacious on Panonychus ulmi in England, 672; effects of fungicidal sprays on populations of, 672.

Psallus seriatus, insecticides against, on cotton in U.S.A., 408, 467.

Psammotermes assuanensis, infesting building timbers in Egypt, 180.

Psammotermes fuscofemoralis, attacking building timbers in Egypt, 180; resistance of Eucalyptus to infestation by, 180; fumigants against, 181.

Pselactus spadix (see Codiosoma).

Pseudaletia separata, on wheat in China, 625.

Pseudaletia unipuncta, on cereals and grasses in N. America, 66, 129, 393, 394; in Hawaii, 283, 428; in Pakistan, 174, 175, 693; bionomics and larval habits of, on rice, 174, 175; question of winter survival of, 394; study of frass production by, as sampling method, 69; natural enemies of, 66, 129, 175; virus disease of, 175, 283, 428, 693; infected by Protozoa, 428.

Pseudaonidia duplex, on Citrus in China, 491; new parasite of, 491.

Pseudaphycus malinus, destroying Pseudococcus comstocki, 164; introduced into

Ontario from Virginia, 164.

Pseudaulacaspis pentagona, in Bermuda, 229; in France, 666, 667; in Italy, 437, 616; in Ontario, 529; on mulberry, 437, 529; on oleander, 229; on peach, 437, 529, 616, 666, 667; development

of, on potato tubers, 437; parthenogenesis in, 437; ant associated with, 229; natural enemies of, 229, 616; insecticides against, 616, 666, 667; method of rearing, 229; use of, for rearing Coccinellids, 229.

pseudobrassicae, Rhopalosiphum.

Pseudococcus, on mulberry in India, 626. Pseudococcus brevipes (see Dysmicoccus). Pseudococcus citri (see Planococcus).

Pseudococcus citriculus, association of Cryptoblabes gnidiella with, on Citrus

fruits in Israel, 612.

Pseudococcus comstocki, on Catalpa, peach and Japanese plum in Ontario, 164; natural enemies of, 164; parathion sprays against, 164.

Pseudococcus maritimus, on stored potatoes in Brazil, 119, 120; BHC fumigant

against, 120.

pseudoconspersa, Euproctis.

Pseudodoniella laensis, on cacao in Dutch New Guinea, 76. pseudoeceticola, Tetrastichus.

pseudomagnoliarum, Coccus.

Pseudomonas, wireworms destroyed by toxins of, 582.

Pseudomonas fluorescens, wireworms destroyed by toxins of, 582.

experiments Pseudomonas pyocyanea, with, against Eurygaster integriceps, 115.

Pseudomonas solanacearum, causing bacterial wilt of banana and Heliconia, 520; insects as vectors of, 520; mechanical transmission of, 520.

pseudoobscura, Drosophila.

Pseudoperichaeta roseanae (insidiosa, auct.), parasitising Ostrinia nubilalis in U.S.S.R., 245.

Pseudosarcophaga mamillata (see Agria). pseudospretella, Hofmannophila.

Pseudotheraptus wayi, relation of ants to, on coconut in Zanzibar, 28, 29;

insecticides against, 29.

Pseudotsuga menziesii (taxifolia), pests of, in Canada, 385, 457, 458, 459, 529; Aphelia viburniana on, in Germany, 375; Dendroctonus obesus infesting logs of, imported into Italy, 532; pests of, in U.S.A., 24, 259, 355, 385, 461, 478, 532; attraction of Trypodendron lineatum to logs of, 459; X-ray detection of D. pseudotsugae in, 259.

Pseudotsuga menziesii glauca, Cydia fletcherana on, in Montana, 460.

pseudotsugae, Dendroctonus.

pseudotsugella, Dioryctria.

psi, Apatele.

Psila rosae (on carrot), in Britain, 432; in Holland, 100, 299, 608; in New York,

226; bionomics of, 299; comparison of injury by Phytomyza lateralis and, 674; insecticides against, 100, 101, 226, 299, 432, 608.

Psilodryinus typhlocybae, parasitising Metcalfa pruinosa in U.S.A., 465.

Psoloessa delicatula, on rangelands in Arizona, 224.

Psorosina hammondi, on apple, pear and Prunus maritima in Canada, 529.

Psychidosmicra, parasitising Oiketicus kirbyi in Costa Rica, 638; parasite of,

Psychidosmicra brethesi, 638.

Psylla mali (on apple), in Canada, 159, 429, 523; in Poland, 676; characteristics and world distribution of, 507; predators of, 429, 523; sprays against, 159, 523, 676.

Psylla pyri, bionomics and predators of,

on pear in Poland, 365.

Psylla pyricola (on pear in British Columbia), Anthocoris melanocerus predacious on, 158; insecticides against, 158, 395, 411, 412; resistance to insecticides in strains of, 284.

Psylla pyrisuga, on pear in Poland, 365. Psylliodes, on cruciferous oil-seed crops in France, 373; natural enemies of, 373. Psylliodes attenuatus, characteristics and

world distribution of, 507.

Psylliodes chrysocephalus, characteristics and world distribution of, 507; factors affecting development and feeding activity of, 205.

Psylliodes elliptica, insecticides against,

on cereals in Turkey, 297.

Psylliodes punctulata, relation of diversity of plant species to populations of, in New York, 69.

Pterandrus rosa (see Ceratitis). ptericoptophagus, Hormiopus.

Ptericoptus, on potato in Argentina, 552; new parasite of, 552.

pteridis, Empoasca.

Pterocarpus marsupium, Coleoptera in logs of, in India, 448; DDT treatment of timber of, 448.

Pterocarya stenoptera, Anomala corpulenta on, in China, 501.

Pteromalus puparum, parasitising Pieris rapae in Japan, 78.

pterophori, Ephialtes (Scambus).

Pteroptrix dimidiatus, parasitising Quadraspidiotus perniciosus in Switzerland, 664.

pterostichus, Thisoicetrinus.

Ptilinus fuscus, bionomics of, on willows and poplars in Poland, 189, 190; as

precursor of fungal disease, 190; insecticides against, 190.

Ptinus fur, in stored foodstuffs in U.S.S.R., 344; experiments on cold-hardiness of stages of, 344.

Ptinus raptor, in stored cereals in U.S.S.R., 344; experiments on cold-hardiness of

stages of, 344.

Ptinus villiger, infesting flour in Manitoba, 225; treatment of bagged flour against, 225.

Ptycholoma peritana, toxicity of Zectran to, 516.

Ptychomyia selecta (parasitising Pristiphora erichsonii), in Canada, 125; in Michigan, 210.

pubescens, Harpalus (see H. rufipes).

pudibunda, Dasychira.

Puerto Rico, *Chloriona orizicola* on rice in, **257**; review of plants infested by *Heliothrips haemorrhoidalis* in, **257**.

pujoli, Paremydica.

pulchellana, Argyrotaenia.

pulchellus, Coccophagus; Ephedrus.

pulicaria, Chaetocnema. pulicarius, Brachypterolus. Pullus (see Scymnus).

Pulvinaria betulae, on vines in Austria, 85.

pumilionis, Chlorops.

Pumpkin Beetle, Red (see Aulacophora foveicollis).

punctatum, Anobium.
punctatus, Dendrolimus.
puncticollis, Podagrica.
punctiferalis, Dichocrocis.
punctigera, Heliothis.

punctigera, Heliothis. punctillum, Stethorus.

punctipes, Aphycus (Metaphycus).

punctiventris, Cleonus (Bothynoderes). punctoria, Nythobia (Angitia, Horogenes). punctulata, Pheidole; Psylliodes.

punctulatus, Phetaole, Psyli punctulatus, Deraeocoris.

puparum, Pteromalus.

purchasi, Icerya.

pusilla, Oscinella frit; Phyllotreta.

pusillus, Cryptolestes.

pusio, Nythobia (Meloboris).

Putoniella marsupialis, bionomics of, on plum in Yugoslavia, 32.

putrescentiae, Tyrophagus.

Pyemotes ventricosus, infesting honey bees in Queensland, 296.

pygmaeoides, Megaselia.

pygmaeus, Ćephus.

Pygostolus falcatus (parasitising Sitona spp.), established in Manitoba, 414; in Sweden, 414.

pyralis, Sacadodes.
pyrastri, Lasiophthicus.
Pyraysta pubilalis (200 Ostr

Pyrausta nubilalis (see Ostrinia).

pyraustae, Perezia.

Pyrazothion, toxicity of, to Tetranychus telarius, 478.

Pyrenone, 500.

Pyrenone Grain Protectant, 402, 403.

Pyrethrum (including pyrethrins), against aphids, 633; toxicity of, to eggs of Bombyx mori, 171; toxicity of, to Musca domestica, 2, 3, 63, 687, 688; against Coleoptera, 180, 418; toxicity of, to Periplaneta americana, 2, 117; against Spodoptera exempta, 530; tests and uses of, against pests of stored products, 2, 3, 4, 225, 270, 402, 403, 500, 560, 561, 587, 591, 687, 688; question of metabolism of, 64; in aerosols, 418; in dusts, 2, 633; in sprays, 180, 530, 633; applied from aircraft, 530; repellent action of, 270; residues of, on wheat and wheat products, 270; with piperonyl butoxide, 3, 63, 225, 270, 402, 500, 560, 587, 591, 687, 688; mechanism of action of piperonyl butoxide with, 688; chromatographic separation of piperonyl butoxide and components of, in mixtures, 562; with other synergists, 3, 63, 687, 688; with other insecticides, 418, 587, 633, 687; toxicity of, unaffected by 2,2diphenylpentanoic acid and 2-diethylaminoethanol, 688; question of effect of perfumery chemicals on efficiency of, 496; determination of constituents of,

pyri, Eriophyes; Psylla; Stephanitis; Ty-phlodromus.

pyricola, Psylla.
pyrisuga, Psylla.

pyrivora, Cydia (Laspeyresia).

Pyroderces rileyi, toxicity of Zectran to, 516.

pyrogramma, Agonoxena.

Pyrolan, colorimetric determination of, 227.

Pyrophyllite, as carrier for insecticide dusts, 560.

Pyrus, Rhopalosiphum fitchii overwintering on, in New Brunswick, 585; experiments with R. padi on, 585. (See also Pear.)

Pyrus prunifolia, Anomala corpulenta on, in China, 501.

Pythium, effects of seed treatments of cotton on incidence of, 398.

Q

Quack Grass (see Agropyrum). Quadraspidiotus juglansregiae, on walnut in California, 491; new parasite of, 491. Quadraspidiotus perniciosus (on fruit trees), in France, 379, 433, 434, 435, 681; in Germany, 681; in Italy, 616; in New Zealand, 167; in Pakistan, 169, 170; in Switzerland, 664, 681; in U.S.S.R., 543; in U.S.A., 379, 491; on rose, 170; reared on water melon, 379; experiments on infestation of currant by, 664; physiological effects of feeding by, 45, 186; natural enemies of, 170, 379, 433, 434, 435, 491, 664, 681; method of rearing, 169; sprays against, 167, 433, 434, 435, 543, 616.

quadratum, Piesma. quadridens, Ceutorhynchus. quadriimpressum, Ambrostoma. quadrimaculatus, Anopheles. quadripunctaria, Callimorpha. quadripustulata, Winthemia. quadrisignatus, Enoclerus; Glischrochilus. quadrituberculatus, Amnemus. quaestor, Coccophagus.

Ouarantine Measures, against cotton pests in U.S.S.R., 113.

quatuordecimcostata, Sternocolaspis. quaylei, Homalotylus.

Quebec, pests of apple in, 26, 159, 454; Caliroa limacina on hawthorn and wild cherry in, 26; forest pests in, 73, 121, 122, 123, 157, 258, 341, 392, 455, 459, 529; Hylastes rufipes transmitting Dutch elm disease in, 124; Epilachna varivestis on beans in, 454; pests of crucifers in 161, 529; Hylemya antiqua on onion in, 455; pests of leguminous forage crops in, 453, 454, 529, 686; phyllody virus of clover in, 686; Ostrinia nubilalis on maize in, 27, 258, 514: Acarapis externus infesting honey bees in, 295; diseases of insects in, 341, 392, 455; natural enemies of pests in, 26, 27, 73, 157, 454, 459.

Queensland, banana pests in, 503; new Agromyzids on pea, Indigofera suffruticosa and sea-kale beet in, 295; Heliothis spp. on tomato, flax and other plants in, 588, 589; H. assulta on Physalis spp. in, 589; Lepidoptera on tobacco in, 503, 588; ants preventing establishment of pastures in, 589; Lyctus brunneus on species of Eucalyptus and Tristania in, 589; mites infesting honey bees in, 295, 296; new parasite reared from Agonoxena sp. in, 52; food-plants of Dacine Trypetids in, 40; key to species of Melanagromyza in, 295.

quercetorum, Haltica. Quercus (see Oak).

Quercus alba, Ennomos subsignarius on, in U.S.A., 418.

Quercus falcata, Ennomos subsignarius on, in U.S.A., 418.

Quercus ilex, pests of, in Spain, 659, 660. Quercus mongolica, new Lithocolletid on, in U.S.S.R., 243.

Quercus rubra, Xyleborus spp. infesting, in Germany, 11; Bucculatrix ainsliella on, in Michigan, 327.

Ouercus suber, insect fauna of, in Sardinia, 660; Lymantria dispar on, in Spain,

Quince, Eriophyes pyri on, in S. Africa, 83; Cydia molesta on, in Brazil, 555; C. molesta on, in Ontario, 163; Lithocolletis spp. on, in Switzerland, 86; Tetranychus viennensis on, in Turkey, 177; Tetranychids on, in U.S.S.R., 242; toxicity of toxaphene to, 555.

quinquefasciatus, auct., Culex (see C. pipiens fatigans).

quinquelineata, Megachile. quinquemaculata, Protoparce.

R

radians, Euxoa.

radiatae, Ips (see I. concinnus); Tarsonemoides.

radicana, Griselda. radicis, Hylobius.

Radioactive Isotopes, effects on insects of γ-radiation from, 472, 568, 607, 686; review of use of y-radiation from. against insects, 607; as tracers (see next entry).

Radioactive-tracer Methods, in studying feeding habits of aphids, 45, 93; in studying oviposition habits of Scaphoideus luteolus, 478; in studying feeding habits of predators, 338, 393; for labelling insects, 29, 265, 356, 429, 471; in investigations on insecticides, 146, 147, 218, 224, 263, 321, 467, 472, 473, 475, 497, 562.

Radish, Phyllotreta spp. on, in Austria, 184; pests of, in Germany, 303; P. cruciferae on, in India, 495; pests of, in U.S.A., 131, 510, 568; in relation to oviposition by Hylemya brassicae, 516, 517; Tetranychus multisetis feeding on, 67; new virus disease of, 131; seed treatment of, 568; insecticide residues in, 303, 479.

Rabbits, enzyme activity of plasma of, on phosphorus compounds, 213.

Rachiplusia ou, on mint in Indiana, 284; virus disease of, 284.

Ragi (see Eleusine coracana).

Ragwort (see Senecio jacobaea). Rainbow Trout (see Trout, Rainbow). Rainfall, in relation to insects, 34, 93, 97, 98, 165, 166, 224, 298, 403, 455, 462, 481, 502; effects of: on diseases of

insects, 107, 335, 341; on electrical charges on plants, 123; on insecticides, 16, 526.

Raisins, effects of, on fecundity of Aptesis basizona, 155; in diets for Itoplectis conquisitor, 208.

Ranunculus, Eurygaster spp. on, in Cali-

fornia, 520.

Rangelands (see Grasses).

rapae, Diaeretus (Aphidius); Pieris; Trybliographa (Cothonaspis).

rapax, Hemiberlesia.

Rape, pests of, in Britain, 318, 430; Omophlus spp. on, in Bulgaria, 348; pests of, in Denmark, 139; pests of, in Finland, 104, 674; Coleoptera on, in France, 373; Ceutorhynchus assimilis on, in Germany, 440; C. pleurostigma on, in Holland, 673; pests of, in Hungary, 308; Autographa nigrisigna on, in Japan, 79; Brevicoryne brassicae on, in New Zealand, 167; Meligethes aeneus on, in Poland, 61, 599; varietal susceptibility of, to B. brassicae and mosaic virus, 167; susceptibility of, to Coleoptera, 318; Myzus persicae reared on, 691; seed treatment of, 104, 674; effects of seed treatments on germination and development of, 675.

Raphatox (see DNC).

Raphimetopus ablutellus, on sugar-cane and grasses in India, 119, 553; bionomics of, 119; parasite of, 119; insecticides against, 553.

rapidus, Adelphocoris. raptor, Ptinus.

Raspberry, Byturus urbanus on, in Britain, 671; Thomasiniana theobaldi on, in Switzerland, 552; Anthonomus rubi on, in U.S.S.R., 360; wild hosts of mosaic virus of, in U.S.A., 132; malathion residues on, 671.

Raspberry, Black (see Rubus occidentalis). Raspberry, Red (see Rubus strigosus).

Rats, metabolism of fenthion in, 475; enzyme activity of homogenates of liver of, on phosphorus compounds,

ratzeburgiana, Enarmonia (Zeiraphera). Rauwolfia serpentina, Saissetia coffeae on, in E. Pakistan, 176.

RE 5305 (see m-sec.-Butylphenyl Methylcarbamate).

recurvalis, Hymenia.

Recurvaria, status of N. American species of, 121.

Recurvaria canusella (see Evagora). Recurvaria condignella (see Evagora). Recurvaria milleri (see Evagora). Recurvaria moreonella (see Evagora). Recurvaria nanella, 121. Recurvaria starki (see Evagora).

redikorzevi, Bryobia.

Reeds, Locusta migratoria breeding in areas of: in Sinkiang, 13; in U.S.S.R., 15.

regina, Phormia.

reniculella, Dioryctria.

repanda, Mocis.

Repellents, against Dacus oleae, 47; discharge of, by Tribolium castaneum, 621: review on production of, by arthropods,

République Soudanaise (see Mali). Rescue-grass (see Bromus unioloides). resinator, Apechthis.

resinella, Rhyacionia (Evetria).

resinosae, Conophthorus; Matsucoccus. Resins, infestation by bark-beetles in relation to exudation of, by pines, 340; fumigant toxicity of, to Dendroctonus spp., 223.

Resins, Synthetic, efficiency of endrin not increased by addition of, 287.

reticulatus, Asolcus simoni; Physcus; Typhlodromus.

Reticulitermes flavipes, in Germany, 649; infesting buildings in Wisconsin, 416; determination of resistance of timber to, 649; effects of fungus on, 428; measures against, 416, 576.

Reticulitermes lucifugus, in Italy, 649; bionomics of, in Spain, 300; determination of resistance of timber to, 86, 649;

measures against, 300.

Reticulitermes lucifugus santonensis, in France, 649; method of testing resistance of timbers to, 649.

Reticulitermes virginicus, effects of fungus on, 428.

Réunion, pests of crops in, 88.

Review of Applied Entomology, completion of fiftieth volume of, 621.

Reviews: Avidov (Z.), Pests of cultivated plants in Israel, 37; Balachowsky (A. S.), Ed., Entomologie appliquée à l'agriculture (vol. 1), 491; Bally (W.), Ferwerda (J. D.) & Morettini (A.), und subtropische Welt-Tropische wirtschaftspflanzen . . . Ölpflanzen, 693; Bremer (H.), Krankheiten und Beschädigungen der Gemüse und Küchenkräuter, 425; Brown (E. S.), The African army worm . . ., 530; Browne (F. G.), The biology of Malyan Scolytidae and Platypodidae, 296; Caswell (G. H.), Agricultural

entomology in the tropics, 75; Dastur (R. H.) & others, Cotton in India (vol. II), 139; Evans (G. O.), Sheals (J. G.) & Macfarlane (D.), The terrestrial Acari of the British Isles . . . (vol. 1), 41; Fulmek (L.), Parasitinsekten der Blattminierer Europas, 492; Haarer (A.E.), Modern coffee production (2nd edn.), 196; Kennedy (J. S.), Day (M. F.) & Eastop (V. F.), A conspectus of aphids as vectors of plant viruses, 357; Kevan (D. K. McE.), Soil animals, 89; Kosov (V. V.) & Polyakov (I. Ya.), Ed., Forecasting . . . pests and diseases of agricultural crops, 15; Paikin (D. M.), Eurygaster integriceps, 543; Péreau-Leroy (P.), Le palmier dattier au Maroc, 54; Polyakov (I. Ya.), Ed., Distribution of pests and diseases of agricultural crops in the U.S.S.R..., 15; Ramsfjell (T.) & Fjelddalen (J.), Sjukdommer og skadedyr på jordbruksvekster . . ., 622; Rupérez Cuéllar (A.), Virosis de insectos, 550; Steinhaus (E. A.) & Smith (R. F.), Ed., Annual review of entomology (vol. 7), 90; Vashkov (V. I.) & Schnaider (É. V.), Khlorofos . . ., 541; Weiser (J.), Die Mikrosporidien als Parasiten der Insekten, 89; An atlas of coffee pests and diseases . . . in Kenya, 490; A of the biological control attempts against insects and weeds in Canada, 692.

Rhabdepyris rhizoperthae, sp.n., parasitising Rhyzopertha dominica in India, 196. Rhabdocline pseudotsugae, treatment of Douglas-fir logs against, 532.

Rhadinaphelenchus cocophilus, Rhynchophorus palmarum as vector of, causing red ring disease of coconut in Trinidad,

Rhadinoceraea micans, bionomics of, on Iris spp. in Yugoslavia, 32; DDT dusts

against, 32.

Rhagoletis cerasi (on cherry), in Germany, 679, 680; in Italy, 137, 311, 535; in Poland, 188, 676; bionomics of, 679; varietal susceptibility of cherries to, 188; migration of, 311, 535; forecasting emergence of, 679; characteristics and world distribution of, 507; insecticides against, 137, 188, 311, 535, 676, 680.

Rhagoletis pomonella (on apple), in Canada, 159, 273; in U.S.A., 18, 268, 400, 403, 417; bionomics of, 18; effects of sprays against Lepidoptera on populations of, 18; favoured by fungicides, 273; killed by controlled-atmosphere storage of apples, 403; insecticides against, 159, 268, 273, 274, 417; question of resistance to DDT in strains of, 400.

rhamni, Erythroneura.

Rhamnus cathartica, distribution of, as primary food-plant of Aphis nasturtii in Lower Saxony, 255.

Rhaphidopalpa (see Aulacophora).

rhenanus, Typhlodromus.

Rhinacloa, predacious on Empoasca in Peru, 560; effects of thiometon sprays on populations of, 560.

Rhipidothrips gratiosus, on oats Ontario, 529; predacious on other

thrips, 529.

Rhizopertha (see Rhyzopertha). rhizoperthae, Rhabdepyris.

Rhodes Grass (see Chloris gayana).

Rhodesia, Northern, predacious ant in,

Rhodesia, Southern, pests of lucerne in, 48, 88; Busseola fusca on maize in, 48; Lamellicorn larvae infesting pastures in, 38; pests of pines in, 81.

Rhodnius prolixus, metabolism of phorate

in, 512, 513.

Rhoecocoris sulciventris (see Musgraveia). Rhopalicus tutela, parasitising Ips typographus in Czechoslovakia, 142; method of rearing, 142.

Rhopalomyzus ascalonicus (see Myzus). Rhopalophorus clavicornis, parasitising Ips typographus in Czechoslovakia, 142; method of rearing, 142.

Rhopalosiphoninus staphyleae tulipaellus (on beet), in France, 605; in Germany, 545, 546; overwintering of, 605; transmission of viruses by, 545, 546, 605.

Rhopalosiphum abietinum, on spruce in Germany, 105; relation of feeding habits to population fluctuations of, 105; insecticides against, 105.

Rhopalosiphum fitchii (Sand.), in New Brunswick, 584; in Pennsylvania, 639; trap for, 639; on cereals and grasses, 584, 585, 639; winter food-plants of, 585; trap for, 639; question of identity of, 584, 585.

Rhopalosiphum fitchii, auct. (see R. padi). Rhopalosiphum insertum, on cereals, 584; on Mespilus germanica, 585; question of status of, 584, 585.

Rhopalosiphum maidis (see Aphis).

Rhopalosiphum padi (on cereals grasses), in China, 625; in New Brunswick, 584, 585; in U.S.A., 95, 160, 401; on Prunus, 585; food-plant preferences of, 95; as vector of barley yellow-dwarf virus, 95, 133, 160, 401; migration of, 95; injection of, with plant viruses, 133; seed treatment against, 401; question of status and identity of, 584, 585.

Rhopalosiphum padi-fitchii, complex referred to as. 585.

Rhopalosiphum poae (see Myzus).

Rhopalosiphum prunifoliae (see R. padi). Rhopalosiphum pseudobrassicae, on cole in New York, 401; relation of, to 689; transmission, enemies and fungus disease of, 401.

Rhopalosiphum viridis, sp.n., as member of R. padi-fitchii complex in Canada,

585.

Rhopalotus substrigosus, parasitic Leucoptera scitella in Italy, 136.

Rhopalus parumpunctatus, on cotton in Turkey, 552. Rhothane (see DDD).

Rhyacionia, species of, from region of

Gulf of Mexico. 19.

Rhyacionia buoliana (on pines), in Argentina, 144, 230; in Belgium, 606; in Canada, 24, 73, 74, 158, 391, 392, 393; in U.S.S.R., 116; in U.S.A., 73, 74, 129, 410, 411, 413, 481, 495, 632; in Uruguay, 230; pine vigour related to damage by, 481; bionomics and adult habits of, 393, 606; sex ratio of, 606; female sex attractant of, 393; natural enemies of, 24, 73, 74, 144, 158, 230, 391, 393; factors affecting parasitisation of, by Eurytoma pini, 157, 158; estimation of populations of, 129, 495; labelled with 60Co, 393; insecticides against, 411, 413, 632.

Rhyacionia duplana, on pines in U.S.S.R.,

116.

Rhyacionia frustrana (on pines in U.S.A.), review of, 495; soil treatment with phorate against, 19; characters distinguishing R. rigidana from, 19.

Rhyacionia pinicolana, U.S.S.R., 116. on pines

Rhyacionia resinella, on pines in U.S.S.R.,

Rhyacionia rigidana, bionomics of, on pines in U.S.A., 19, 20; parasites of, 20; characters distinguishing, 19.

rhyacioniae, Hyssopus.

Rhynchites aequatus, on apple in Finland, 103.

Rhynchites cupreus, characteristics and world distribution of, 507.

Rhvnchites heros, characteristics and world distribution of, 507.

Rhynchocoris humeralis, on Citrus in Formosa, 591; insecticides against,

Rhynchophorus palmarum, as vector of nematode causing red ring disease of coconut in Trinidad, 575; cage for collecting excreta of, 575.

Rhyncolus culinaris, infesting building

timber in U.S.S.R., 112. Rhyparobia maderae (see Leucophaea).

Rhyzopertha dominica (in stored cereals), in Germany, 11; in India, 196; in Sicily, 300; new parasite of, 196; Dibrom ineffective against, 573.

Ribautiana tenerrima (see Typhlocyba).

ribeana, Pandemis.

ribesii, Nematus; Syrphus.

ribis, Capitophorus (Cryptomyzus); Cecidophyopsis (Cecidophyes, Eriophyes, Phytoptus).

ribisnigri, Myzus (Nasonovia).

Rice, pests of, in China, 501, 625; Tryporyza incertulas on, in Formosa, 328; Rupela albinella on, in Dutch Guiana, 93; pests of, in India, 315; Lepidopterous stem borers of, in Israel, 350; pests of, in Italy, 182, 350; pests of, in Japan, 78, 350; Chilo suppressalis on, in Malaya, 78; Chloriona spp. on, in Mexico, 518; pests of, in E. Pakistan, 174; Cicadulina bipunctella on, in Philippines, 96; Chloriona orizicola on, in Puerto Rico, 257; grasshopper on, Turkey, 351; pests of, in U.S.A., 132, 329, 405, 421, 508; *Blissus leucopterus* on, in Venezuela, 98; systematic list of pests of, in South-East Asia and Pacific area, 423; not attacked by Laelia exclamationis, 79; effects of stage of development of, on larval growth of Chlorops oryzae, 78; comparative susceptibility of Oryza ridleyi and, to attack by Chilo suppressalis, 78; insects transmitting virus diseases to, 132, 257, 350, 405, 508, 518, 689, 690; transmission of virus of, to other cereals, 132; in baits for Gryllotalpa, 442; translocation of systemic insecticides in, 405; seed treatment of, 329, 421, 625; effect of seed treatments on

germination and growth of, 65, 625. Rice (Stored), Coleoptera in, 571, 591, 626; systematic list of pests of, in South-East Asia, and Pacific area, 423; treatment of, with malathion, 571; malathion residues in, 571; effects of fumigants on germination of, 579. Rice Stem Borer (see Chilo suppressalis).

Rice Weevil (see Sitophilus oryzae).

richteri, Solenopsis saevissima.

ricini, Attacus cynthia.

Ricinus communis, Eutetranychus banksi on, in India, 627; E. lewisi on, in U.S.A., 517.

riehmi, Therioaphis (Myzocallidium). rigidana, Rhyacionia.

rilevi. Pyroderces.

Ripersia sacchari, new parasite reared from, in India, 91.

Risella 17, 687.

ritchiei, Achrysocharella.

Rivellia, root nodules of groundnut damaged by, in Congo, 609; experiments with other leguminous plants and, 609.

Robinia pseudacacia, Aphis craccivora on, in Bulgaria, 59; Eulecanium corni on, in U.S.S.R., 13.

robinsoni, Kakimia.

robusta, Sphaerophoria.

robustum, Heterostylum.

robustus, Kakothrips (see K. pisivorus).

Rockmelon (see Cucumis melo).

Rodolia cardinalis, introduced into Puerto Rico against Icerya purchasi, 257.

roddi, Eurytoma (Bruchophagus).

rogersi, Macrosiphum (Acyrthosiphon) malvae.

Rogas, parasitising Thymelicus lineola in Ontario, 162.

Rogas dendrolimi (parasitising larvae of Dendrolimus superans), in Mongolia, 116; in Siberia, 13.

Rogas dimidiatus, parasitising Apamea anceps in U.S.S.R., 363.

Rogas eurinus, parasitising Apamea anceps in U.S.S.R., 363.

Rogas hyphantriae, parasitising Hyphantria cunea in Yugoslavia, 682; competition between other parasites and, 682.

Rogas kanpurensis, parasitising Earias spp. in India, 37.

Rogor (and Rogor L) (see Dimethoate).

Ronnel (see Fenchlorphos).

Rooks, destroying Apamea anceps in U.S.S.R., 363.

Root Nodules, destroyed by insects, 134, 609.

Roptrocerus xylophagorum, parasitising Ips typographus in Czechoslovakia, 142; method of rearing, 142.

rorellus, Hyponomeuta.

Rosa palustris, primary food-plant of Macrosiphum euphorbiae in Maine, 461.

rosa, Ceratitis (Pterandrus).

rosae, Athalia; Aulacaspis; Eurytoma; Macrosiphum; Psila; Typhlocyba (Edwardsiana).

rosanus, Archips (Cacoecia).

Rose, aphids on, in Canada, 23; Bracon brevicornis recorded on, in Czechoslovakia, 302; Tetranychus telarius on, in Holland, 193; Coraebus rubi on, in Italy, 299; Quadraspidiotus perniciosus on, in Pakistan, 170; pests of, in U.S.S.R., 113, 355, 361; pests of, in

U.S.A., 23, 212, 264, 272, 324, 477, 633; fungus diseases of, 264, 324; systemic treatments of, 212; damage by cellulose polymers to, 272; effects of combined sprays on winter survival of, 324.

roseanae, Pseudoperichaeta.

roseus, Anuraphis (see A. plantaginea). rossi, Dioryctria; Lindingaspis.

rostrata, Aelia.

rostratum, Eutrombidium.

Rotenone, tests and uses of, against Coleoptera, 202, 279; against Empoasca, 560; against Lepidoptera, 337, 383, 525; toxicity of, to Musca domestica, 202; effects of temperature on toxicity of, 202; in dusts, 279, 560; in sprays, 337, 383, 525; effects of drugs used in combination with, 279; and DDT, 337, 525; and sulphur, 560. (See also Derris.)

rothi, Campoplex (Omorga).

rotundața, Megachile.

rouxi, Habrolepis.

Royal Jelly, in diet for Dusmetia sangwani, 406.

Rubber, baited strips of, for control of *Piophila casei*, **100.** (See *Hevea*.)

rubens, Ceroplastes.

rubi, Amphorophora; Anthonomus; Coraebus.

rubidus, Laricobius.

rubiella, Incurvaria.

rubifolii, Aphis.

rubrescens, Heliothis.

rubrioculus, Bryobia.

rubrocinctus, Selenothrips. rubropilosa, Atta sexdens.

Rubus, Byturus urbanus on species of, in Britain, 671; aphids as potential vectors of viruses of, 529.

Rubus fruticosus, Erythroneura rhamni overwintering on, in Italy, 661.

Rubus occidentalis, transmission of wineberry virus to, by aphid and grafting, 132.

Rubus parviflorus, Amphorophora parviflori on, in British Columbia, 529; Tetranychus multisetis feeding on, 67.

Rubus phoenicolasius, Amphorophora rubi and mosaic virus of, in U.S.A., 132; transmission of mosaic virus of, to raspberry, 132.

Rubus procerus, Macrosiphum fragariae on, in British Columbia, **529**.

Rubus spectabilis, Amphorophora forbesi on, in British Columbia, 529.

Rubus strigosus, use of, in experiments with Amphorophora rubi and raspberry mosaic, 132.

Rubus vitifolius, Otiorhynchus cribricollis on, in California, 509.

Ruelene (see O-Methyl O-4-tert.-Butyl-2chlorophenyl Methylphosphoramidate). rufa, Formica; Plastophora; Vespula. rufa rufopratensis minor, Formica (see F.

polyctena).

rufata, Apechthis.

ruficornis, Trigonotylus.

rufidorsus, Lygus (Liocoris).

rufimitrana, Enarmonia (Zeiraphera). rufipes, Harpalus; Hylastes (Hylurgopinus); Necrobia; Phytomyza.

rufiventris, Apate monachus; Asolcus.

rufobrunnea, Oncopera.

rufofascialis, Noctuelia.

rufopratensis minor, Formica rufa (see F. polyctena).

rufovillosum, Xestobium. rugicollis, Plesiocoris. rugulipennis, Lygus.

Ruguloscolytus rugulosus (see Scolytus). rugulosus, Scolytus (Ruguloscolytus).

Rumania, Eurydema ornatum on crucifers in, 57; Aphthona euphorbiae on flax in, 608; Ostrinia nubilalis on maize in, 56; Apion spp. on red clover in, 56; Lepidoptera on fruit trees in, 56, 57; Leptinotarsa decemlineata on potato in, **366, 681**; parasites of *Apion* in, **56.**

Rumex, Aphis fabae on, in California, 419.

Rumex obtusifolius, Pegomyia sp. on, in Italy, **662.**

rumicis, auct., Aphis (see A. fabae).

Rungsia kurdjumovi (on grasses), in New Brunswick, 585; in Wisconsin, 160.

Rupela albinella, bionomics of, on rice in Dutch Guiana, 93; effects of rainfall on diapause in, 93.

rusci, Ceroplastes.

ruskini, Xenufens.

rustica, Buprestis; Dexia.

rutilla, Eucarcelia (see Carcelia obesa).

rutilus, Perilitus.

Ryania, against aphids, 148; diverse effects of, against Cydia pomonella, **159**, **466**, **664**; toxicity of, to *Rhagoletis* pomonella, 268; effects of, on beneficial arthropods, 36; in sprays, 36, 159, 466;

and glyodin, 36.

Rye, pests of, in Bulgaria, 109, 348; Calligypona pellucida on, in Finland, 102; Meromyza saltatrix on, in Japan, 96; overwintering of Macrosteles fascifrons on, in Ontario, 70; Lepidoptera on, in U.S.S.R., 362, 596; survival of *Chloriona orizicola* on, 132; transmission of rice virus to, 132; experiments on feeding response of wireworms to extracts of germinating seed of, 528; effects on vetch fauna of sprays against pests of, 211; (stored), development of Sitophilus granarius on, 206.

Rye-grass (see *Lolium* spp.).

S

S-421 (see Octachlorodipropyl Ether). Sabadilla, mixture of sulphur and, against Rhynchocoris humeralis, 591.

sabulosus, Otiorhynchus.

Sacadodes pyralis, distribution of, 357, 507; survey of data on, 507.

saccharalis, Diatraea.

sacchari, Aphis; Ripersia; Xanthoencyrtus. saccharina, Aeneolamia varia; Eldana; Lepisma.

saccharivora, Saccharosydne.

Saccharosydne saccharivora, malathion spray against, on sugar-cane in Jamaica,

Sacks, types of, in relation to penetration by Callosobruchus maculatus, 500, 501; treatments of, against Coleoptera in stored products, 499, 500, 685. saevissima, Solenopsis.

BHC Sahlbergella, sprays ineffective against, on cacao in Ivory Coast, 7.

Sahlbergella singularis (on cacao), in Ghana, 651; in Nigeria, 652; relation of, to swollen-shoot virus and dieback, 652; insecticides against, 651.

Saissetia, new parasite of species of, in

Zanzibar, 491.

Saissetia coffeae (hemisphaerica), in California, 151, 491; bionomics of, on Rauwolfia serpentina in E. Pakistan, 176, 177; production of toxin by, 176; parasites of, 151, 491.

Saissetia nigra, question of new parasite

of, in S. Africa, 491.

Saissetia oleae, in California, 150, 491, **519**; in Pakistan, **491**; on *Citrus*, **519**; on oleander, 491; parasites of, 150, 491, 519.

Saissetia perseae (see S. nigra).

saliceti, Melasoma.

Salicicola davatchi, on pistachio Turkey, 609.

salicis, Stilpnotia.

saliens, Criocoris.

Salix, Cryptorhynchus lapathi Hungary, 307.

Salix alba, pests of, in U.S.S.R., 354.

Salix americana, Cryptorhynchus lapathi on, in Hungary, 307; Agrilus acutangulus on, in Yugoslavia, 138.

Salix amygdalina, Agrilus acutangulus on, in Yugoslavia, 138.

Salix daphnoides, Melasoma saliceti on, in Poland. 40.

Salix purpurea, Phyllodecta tibialis on, in Poland, 40; Agrilus acutangulus on,

in Yugoslavia, 138.

Salix viminalis, Chrysomeloids on, in Poland, 40; Agrilus acutangulus on, in Yugoslavia, 138.

saltatrix, Meromyza.

Sand, volatility of insecticides in, 416. sanguinolenta, Cercopis.

sangwani, Dusmetia.

Sanninoidea exitiosa (see Aegeria). santonensis, Reticulitermes lucifugus.

Saperda carcharias (on poplar), in Czechoslovakia, 600; in U.S.S.R., 114;

measures against, 114, 600.

Saperda populnea (on poplar), in Czechoslovakia, 196, 600; bionomics of, in Spain, 355; varieties of poplar in relation to, 196; mortality factors of, 600; measures against, 600.

sarcitorius, Apanteles; Ichneumon. Sarcodexia lambens (see Sarcophaga).

Sarcophaga, parasitising larvae of Caligo eurilochus in Ecuador, 274.

Sarcophaga aldrichi, attacking Stilpnotia salicis in British Columbia, 126.

Sarcophaga lambens, parasitising Oiketicus kirbyi in Costa Rica, 638; parasitising Caligo eurilochus in Ecuador, 274. sarothamni, Anthocoris.

Sarothamnus scoparius, Coleoptera on, in

England, 30, 656. sasakii, Carposina; Sitophilus (see S.

oryzae).

Saskatchewan, Pristiphora erichsonii on conifers in, 124, 125, 456; Mirids on lucerne in, 72; Etiella zinckenella on other leguminous plants in, 529; Anuraphis tulipae on carrots in, 529; pests of stored grain in, 154, 155, 529; Elaterids in, 155, 156; forecasting outbreaks of grasshoppers in, 575, 631; Coleoptera controlling Linaria vulgaris in, 386; other beneficial insects in, 125, 161.

Satellites, Artificial, experiments with various atmospheres in relation to use of cockroach in, 325.

saucia, Peridroma.

Sawdust, as carrier for insecticides, 402. sayi, Netelia.

Scab, Apple (see Venturia inaequalis). scabriculus, Norbanus; Trematopygus.

Scale Insects, ants associated with, 229, 519, 531; relation of, to *Cryptoblabes gnidiella*, 612; natural enemies of, 53, 91, 150, 151, 164, 170, 229, 257, 266, 379, 380, 382, 433, 434, 435, 490, 491, 519, 529, 564, 613, 614, 683, 688;

fungus diseases of, 54, 587, 588; sooty mould associated with, 176, 587, 613; control of noxious plants by, 587, 588; parthenogenesis in, 437; dispersal of, 35; effect of infestation by, on oxygen consumption of plants, 45; estimation of populations of, 83, 84; methods of rearing, 169, 229; new species of, 234.

Scambus (see Ephialtes).

Scaphoideus littoralis, on vines in France, 372; question of introduction of, from N. America, 372; light-trap catches of, 372.

Scaphoideus luteolus, transmitting phloem necrosis of elm in U.S.A., 478; oviposition habits of, 478; labelled with

³²P, 478.

Scaphytopius acutus, and virus diseases of peach and cherry in New York, 23; other *Prunus* spp. in relation to, 23; fluctuations in populations of, 23; traps for, 23.

Scapsipedus micado, in Alabama, 567; in

Japan, 567.

Scaptocoris divergens, on banana in Honduras, 565, 566; soil and moisture relations of, 566; soil fungi inhibited by secretions of, 565.

Scaptomyzella flava, characters distinguishing injury to rape by, 139.

Schedius telenomicidus (see Ooencyrtus). Schistocerca gregaria, in Aden, 286; in Eritrea, 286; in India, 116, 500; in Morocco, 54, 383; in Somalia, 286; characteristics and world distribution of, 506; factors affecting development and survival of, 5, 6, 285, 286; sensory equipment of, 423; flight habits of, 423, 500; nervous control of flight in, 620; biochemistry of maturation in males of, 423; measurement of carbondioxide output of, 40; cannibalism among hoppers of, 172; infected by Nosema locustae, 582; phases of, 116, 286; colour patterns of, 5, 116; tests of sprays against, 383; pick-up of spray droplets by, 291; factors affecting toxicity of insecticides to, 164, 170, 608, 658; factors affecting resistance to insecticides in, 201; metabolism and excretion of aldrin and dieldrin by, 321.

Schizaphis graminum (see Toxoptera). Schizolachnus piniradiatae, on pines in Canada and U.S.A., 458; bionomics and population distribution of, 458, 459; natural enemies and fungus disease of, 459.

Schizotetranychus andropogoni, on sugarcane in India, 628; toxicity of acaricides to, 628.

schlechtendali, Vasates.

schneideri, Chermes nordmannianae. Schoenobius incertulas (see Tryporyza).

Schradan, against aphids, 148, 188, 240; against Cerococcus catenarius, 558; study on uptake of, by Dysdercus koenigii, 497; against Phyllotreta pusilla, 511; against mites, 191, 678; effects of, on parasite of Eriosoma lanigerum, 312; metabolism of, in insects, 63, 64; in sprays, 188, 191, 240, 312, 511, 558, 678; systemic action and other methods of application of, 188, 558; labelled with ³²P, 497; and phosphamidon, 511; and sesamex, 63.

Sciara, species of, infesting mushrooms in Brazil, 644; sprays against, 644.

Sciara amoena, on carnations in Germany,
254; parathion drench against, 254.
Sciara tritici, on cereals in Ontario, 529.
Scirpophaga nivella, treatment of sugarcane setts against, in India, 297.

Scirtothrips aurantii, map of distribution

of, 1.

Scirtothrips bispinosus (see Dendrothrips). Scirtothrips citri, map of distribution of, 1. Scirtothrips signipennis (see Chaetana-phothrips orchidii).

scissifrons, Sitona. scitaria, Striglina. scitella, Leucoptera.

scoliopi, Macrosiphum. Scolytus multistriatus (vector of Cerato-

cystis ulmi on elm), introduced into Ontario, 124; in U.S.A., 217, 619, 641; nematodes associated with, 619; method of rearing, 217; measures against, 217, 218, 641.

Scolytus rugulosus, estimation of popula-

tions of, on willow in France, 83.

Scolytus tsugae, on conifers in British Columbia, 458.

Scolytus unispinosus, bionomics of, on conifers in British Columbia, 458; Dendroctonus pseudotsugae associated with, 458.

Scopiorus (see Polyblastus).

Scotinophara lurida, characteristics and world distribution of, 507.

Scots Pine (see *Pinus sylvestris*). Scrobipalpa, type and scope of, 235.

Scrobipalpa atriplicella, characters distinguishing Phthorimaea operculella from, 236.

Scrobipalpa heliopa, bionomics of, on tobacco in India, 236; characteristics and world distribution of, 507; eggparasite of, 236; as type species, 235.

Scrobipalpa instabilella, nomenclature of, 235.

Scrobipalpa ocellatella, in Bulgaria, 347; in Czechoslovakia, 236; in Germany, 440, 441; characteristics and world distribution of, 507; on beet, 347, 440, 441; effects of temperature on development of, 236; measures against, 347, 441; nomenclature of, 235, 236.

scutellaris, Asolcus; Erythrocera (Eurys-

thaea).

scutellata, Sturmia (Blepharipoda).

scutellatus, Pachynematus.

Scutigerella immaculata, damaging maize in France, 433; tests and uses of insecticides against, 433; favoured by soil treatments with chlorinated hydrocarbons, 433.

Scymnus, predacious on Planococcus citri in Israel, 614; parasite of, 614.

Scymnus hilaris, effect of insectide sprays on populations of, in Japan, 36.

Scymnus impexus (predacious on Chermes piceae), introduced into Canada from Europe, 394, 456; bionomics and winter mortality of, 394.

SD (see also Shell).

SD-2966 (see Methyl 3-(Ethoxy(p-dimethylaminophenyl)phosphinyloxy)crotonate).

SD 3348 (see 5,6,7,8,9,9-Hexachloro-1,2,-3,4,4a,5,8,8a-octahydro-1,4,5,8-dimethanophthalazine Sulphate).

SD 3450 (see 5,6,7,8,9,9-Hexachloro-1,4,-4a,5,8,8a-hexahydro-1,4,5,8-dimethanophthalazine N-Oxide).

SD-3562 (see Dimethyl 1-Dimethylcarbamoyl-1-propen-2-yl Phosphate).

SD-4402 (see 1,3,4,5,6,7,8,8-Octachloro-3a,4,7,7a-tetrahydro-4,7-methanoph-thalan).

SD-5539 (see Dimethyl 1-(m-Nitrobenzyloxycarbonyl)-1-propen-2-yl Phosphate).

SD-5656 (see N,N-Diallyl-3-(dimethoxy-phosphinyloxy)crotonamide).

Sea-kale Beet (see Chard).

Sedges, overwintering of Sphenophorus callosus on, in Georgia, 152.

segmentator, Phytodietus.

segetum, Agrotis (Euxoa).

seitneri, Tomicobia.

Seius, saprophagous species of, on soy bean in Delaware, 414.

selecta, Ptychomyia (Bessa).

Selenothrips rubrocinctus, on cacao in Dutch New Guinea, 76; map of distribution of, 1.

Selinon, containing DNC (q.v.), 111. Semasia diniana (see Enarmonia griseana). semipunctatum, Graphosoma. semirufa, Podagrica malvae.

semistriatus, Asolcus.

senatoria, Anisota.

Senecio, development of Tyria jacobaeae on species of, 260.

Senecio cineraria, 260.

Senecio jacobaea, use of Tyria jacobaeae against: in Canada, 260; in Victoria, 452.

Senecio pauperculus, 260.

Senecio vulgaris, 260; and virus diseases

of crop plants, 17, 304.

Senegal, pests of field and stored groundnuts in, 298; virus disease of groundnuts, in 298.

senex, Lathromeris.

seniculum, Apion.

separata, Pseudaletia.

separatella, Maliarpha.

septemfasciata, Nomadacris.

septempunctata, Coccinella.

seriatus, Psallus.

Serica (see Maladera).

sericaria, Ephestia (Anagasta) (see A. kuehniella).

sericea, Blastothrix.

Sericesthis pruinosa, characteristics and world distribution of, 507.

sericeus, Apanteles.

Serine, in saliva of Aphis pomi, 95.

l-Serine, in relation to feeding response of wireworms, 528.

serinopa, Nephantis.

Serratia marcescens, infecting cultures of Ips confusus, 300.

serricorne, Lasioderma.

sertifer, Neodiprion (Diprion).

Sesame (see Sesamum).

Sesamex, as synergist: with phosphorus compounds, 63; with pyrethrins, 63; mode of action of, with Sevin, 263; effects of, on oxidation of other insecticides, 63, 64.

Sesamia, characters of pupae of, 292.

Sesamia calamistis (on cereals), in Nigeria, 292; in Kenya, 179; measures against, 293.

Sesamia cretica, on maize in France, 372; characteristics and world distribution

of, 507.

Sesamia nonagrioides, in France, 372, 666; in Morocco, 383; in Sardinia, 182; on maize, 372, 383, 666; on rice, 182; bionomics of, 182, 383; parasite of, 182; insecticides against, 383; sowing date in relation to infestation by, 666; nomenclature of, 182.

Sesamia nonagrioides botanephaga, on

cereals in Nigeria, 292.

Sesamia penniseti, on cereals in Nigeria, 292.

Sesamia poephaga, on cereals in Nigeria, 292.

sesamiae, Apanteles.

Sesamum orientale (indicum), Antigastra catalaunalis on, in India, 316; review of pests and diseases of, 693.

Sesbania bispinosa (aculeata), Lepidoptera

on, in India, 316.

setariae, Carolinaia (Hysteroneura).

setarius, Calomycterus.

setifer, Bustomus.

setulosus, Bustomus; Eremnus.

Sevin, against, aphids, 333, 394, 468, 498, 508, 515, 637, 643, 667, 668; against other Hemiptera, 168, 395, 478, 515, 559, 591; against Coleoptera, 16, 148, 184, 252, 408, 515, 637; against Hoplocampa testudinea, 666; against Rhagoletis pomonella, 268, 273, 274; cross-resistance to, in parathion-resistant strain of Musca domestica, 263; against Lepidoptera, 9, 21, 26, 92, 145, 211, 220, 269, 276, 333, 335, 337, 351, 394, 398, 406, 408, 409, 411, 466, 469, 471, 476, 480, 487, 508, 514, 515, 516, 557, 558, 637, 641, 644, 645, 646, 663, 669; against Frankliniella, 408; Tetrany-chids favoured by, 394, 408, 666; effects of, on beneficial arthropods, 159, 378, 508; toxicity of, to bees, 276, 394, 412, 515; factors affecting toxicity of, 16, 274; mammalian toxicity of, 146; metabolism of, 263, 635; in dusts, 145, 184, 351, 409, 471, 508, 557, 591, 637, 644; in granules, 469; in sprays, 9, 16, 21, 26, 159, 168, 184, 213, 220, 252, 268, 269, 276, 223, 235 213, 220, 252, 268, 269, 276, 333, 335, 337, 351, 394, 395, 398, 408, 411, 466, 468, 478, 487, 508, 514, 515, 526, 557, 558, 591, 637, 641, 643, 645, 646, 663, 666, 669; as deposit, 273, 394; aircraft application of, 220, 276; seed treatment with, 65, 66, 184; question of vapour toxicity of, in soils, 416; tests of, against pests of stored products, 213; effects of, on growth or respiration of plants, 65, 562; residues of, in milk, 92, 266; residues of, in plants and plant products, 228, 398, 526, 561; labelled with ¹⁴C, 263; with piperonyl butoxide, 476, 508; effects of piperonyl butoxide on development of resistance to, 148; with other synergists, 263, 476; and azinphos-methyl, 268; and Bacillus thuringiensis, 514; and lead arsenate. 268; and toxaphene, 478.

sexdens, Atta.

sexta, Protoparce.

shawiana, Hendecaneura.

Sheep, transport of Eriococcus orariensis

by, 587; insecticide residues in tissues of, 280, 384.

sheldoni, Aceria.

Shell 1836 (see Diethyl 2-Chlorovinyl Phosphate).

Shell 3562 (see Dimethyl 1-(Dimethylcarbamoyl)-1-propen-2-yl Phosphate).

Shell BAS-4294 (see Dimethyl Methylbenzyloxycarbonyl)-1-propen-2yl Phosphate).

Shell BAS-4475 (see Dimethyl 1-(2,4-Dichlorobenzyloxycarbonyl)-1-propen-2-yl Phosphate).

Shell Ondina 17, 2, 3.

Shell OS-1808 (see Diethyl 2-Ethoxycarbonyl-1-methylvinyl Phosphate).

Shell Risella 17, 3.

Shell SD-2244 (see Diethyl 1-(Dimethylcarbamoyl)-1-propen-2-yl Phosphate). Shell SD-3562 (see Dimethyl 1-(Dimethylcarbamoyl)-1-propen-2-yl Phosphate).

Shell SD-4092 (see Dimethyl 1-Benzyloxycarbonyl-1-propen-2-yl Phosphate).

Shell SD-4329 (see Dimethyl 1-(p-Chlorobenzyloxycarbonyl)-1-propen-2-yl Phosphate).

Shell SD-4402 (see 1.3.4.5.6.7.8.8-Octachloro-3a,4,7,7a-tetrahydro-4,7-metha-

nophthalan).

Shell SD-5533 (see Dimethyl 1-(p-Nitrobenzyloxycarbonyl)-1-propen-2-yl Phosphate).

Shell SD-5539 (see Dimethyl 1-(m-Nitrobenzyloxycarbonyl)-1-propen-2-yl Phosphate).

Shg 1496, in sprays against Dacus oleae,

shikhae, Palorus.

Shortleaf Pine (see Pinus echinata).

Siam, Aonidiella aurantii on Citrus in, 491. SIB No. 7, in bait-sprays for Anastrepha ludens, 280; constituents of, 280.

Siberian Cedar (see Pinus cembra).

Siberian Larch (see *Larix sibirica*). sibirica, Aelia.

sibiricus, Dendrolimus (see D. superans). Sida carpinifolia, Earias on, in India, 37. Sida cordifolia, Earias on, in India, 37.

Sierra Leone, first record of Contarinia sorghicola on sorghum in, 685.

Sigalphus thoracicus (see Triaspis). Siglure, isomers of, as attractants for Ceratitis capitata, 383.

signipennis, Scirtothrips (see Chaetanaphothrips orchidii).

Silkworms (see Antheraea pernyi and Bombyx mori).

silvaticus, Tetrastichus.

silvestrii, Coccophagus.

silvestris, Phorocera (Parasetigena). Simaethis pariana (see Anthophila). similalis, Loxostege.

similis, Cicadulina; Diprion; Tachinomyia. simoni, Asolcus.

simplex, Erythroneura; Oligonychus (Paratetranychus).

simulans, Praon.

Sinapis alba, susceptibility of, to Coleoptera in Britain, 318.

Sindar Corp. A-1634 (see Butyl 1,3-Dimethylbutyl Acetal of Acetaldehyde). Sinea, predacious on cotton pests in

Arizona, 639. sinensis, Ceroplastes.

Singapore, Nacoleia octasema in, 354.

singularis, Sahlbergella.

sinicus, Adoretus.

Sinigrin (see Potassium Myronate).

sinuatus, Agrilus.

Sipha agropyrella (see Rungsia kurdjumovi).

siro, Acarus.

Sisymbrium orientale, use of, in experiments with aphids and cauliflower mosaic virus, 197.

Siteroptes graminum, in Alberta, 156; relation of, to silver top of grasses, 156. Sitka Spruce (see *Picea sitchensis*).

Sitobion fragariae (see Macrosiphum). Sitodiplosis mosellana (on wheat), in Britain, 288,420; in Canada and U.S.A., 420; susceptibility of varieties of wheat to Mayetiola destructor and, 420;

delayed emergence of, 288. Sitona, on lucerne in Bulgaria, 60; in Italy, 391; on clover in U.S.S.R., 114;

parasites of, 391, 414.

Sitona crinitus, parasite of, in France, 390.

Sitona cylindricollis (on Melilotus in Canada), adult habits of, 319, 414; termination of diapause in, 93, 94; factors affecting dispersal of, 318; parasites of, 390, 391, 414; traps for, 319; method of rearing, 94.

Sitona decipiens, on leguminous forage crops in Finland, 41, 102; bionomics

of. 102.

Sitona flavescens, bionomics of, on leguminous forage crops in Finland, 102.

Sitona hispidulus, bionomics of, on leguminous forage crops in Finland, 102; on alsike clover in Oregon, 275; parasites of, in Sweden, 390, 391; relation of root-rot fungi to, 275; soil treat-ments against, 275; larval characters of S. lineatus and, 496.

Sitona humeralis, parasites of, in France,

390, 391.

Sitona lineatus, in British Columbia, 134; in Finland, 41, 102; in Sweden, 391; in U.S.A., 134; on leguminous forage crops, 41, 102, 134; on peas, 102, 134; root nodules destroyed by, 134; bionomics of, 102, 134; parasite of, 391; measures against, 134; larval characters of *S. hispidulus* and, 496.

Sitona scissifrons, parasitised by Perilitus

rutilus, 391.

Sitona sulcifrons, in Finland, 41, 102; on leguminous forage crops, 102; on Phleum pratense, 41; bionomics of, 102.

Sitona suturalis, bionomics of, on leguminous forage crops in Finland, 102.

Sitophilus, study on heating of stored

grain in relation to, 548.

Sitophilus granarius (in stored cereals), in Bulgaria, 347; in China, 693; in Germany, 11; in U.S.S.R., 344; effects of type of grain on development of, 206; experiments on cold-hardiness of stages of, 344; recognition of larval instars of, 424; in diet for Ahasverus advena, 657; effects of physical factors on stages of, 607, 632; tests and uses of insecticides against, 3, 4, 212, 347, 516, 603; fumigant action of insecticides against, 9, 267, 347, 573; effects of hydrogen cyanide on activity and respiration of, 584; effect of concentration on toxicity of insecticides to, 117, 118; resistance to insecticides in strains of, 82, 128, 129.

Sitophilus oryzae (in stored cereals), in Brazil, 558; in Colombia, 560; in Germany, 11; in India, 499, 500, 685; in Kenya, 51; in Louisiana, 403, 571; effects of type of grain on development of, 206; recognition of larval instars of, 424; effects of γ-radiation on stages of, 607; repellency of insecticides to, 270; tests and uses of insecticides against, 2, 3, 51, 118, 212, 215, 270, 402, 403, 499, 500, 554, 558, 560, 561, 571, 573, 591, 684; cuticular structure in relation to contact toxicity of insecticides to, 314; characters and synonymy of, 1, 2; S. zeamais misidentified as, 2.

Sitophilus oryzae minor (see S. oryzae). Sitophilus oryzae platensis (see S. zeamais). Sitophilus sasakii (see S. oryzae).

Sitophilus zeamais, in Argentina, Brazil, Chile and Uruguay, 2; intercepted in stored grain in Germany, 11; characters and synonymy of, 2; misidentified as S. oryzae, 2.

Sitotroga cerealella (in stored cereals), in Colombia, 560; in Illinois, 269, 414; relation of, to infestation of grain by Aspergillus spp., 414, 415; parasitism of eggs of, by Trichogramma spp., 128, 381, 681; predacious mites in cultures

of, 376; disinfection of, 415; treatments of stored grain against, 269, 560, 561.

Skatole (see β -Methylindole).

SKF 525A (see 2-Diethylaminoethyl 2,2-Diphenylpentanoate).

Skins, Dermestes infesting, in S. Africa, 49; treatment of, with insecticides, 49. smaragdina, Oecophylla.

smaragdula, Nezara viridula.

Sminthurus viridis, bionomics and factors affecting populations of, on clover and pastures in Victoria, 34; characteristics and world distribution of, 507; mite predacious on, 34; sprays against, 34. smithi, Aphidius; Eotetranychus.

Snails, biology of Sciomyzids attacking, in N. America and Europe, 91.

Sodium Arsenite, in baits and bait-sprays against *Dacus oleae*, 308, 533.

Sodium Carboxymethyl Cellulose, toxicity of, to *Tetranychus telarius*, 272.

Sodium Chloride, attraction of insects to, 635.

Sodium Fluoride, susceptibility of Galleria mellonella to virus disease unaffected

by, 427. Sodium Fluoroacetamide, in sprays against *Brevicoryne brassicae*, 74.

Sodium Fluosilicate, against Coleoptera, 49, 360; treatment of rubber trees with, against termite, 77; in dusts, 49, 360; treatment of skins with, 49; toxicity of, to strawberry, 360.

Sodium Hydroxide, treatment of storerooms with, against *Lasioderma serricorne*, **58**; decomposition of insecticide residues in olives by, **238**, **563**.

Sodium Hypochlorite, disinfection of Sitotroga cerealella with, 415.

Sodium Pentachlorophenate, soil treatment with, against termites, 577.

Sodium Propionate, use of, against moulds in rearing media for insects, 258.

Sogata cubana (see Chloriona). Sogata orizicola (see Chloriona).

Sogatelenchus mexicanus, gen. et sp.n., parasitising Chloriona spp. in Mexico, 518.

Soil, water content of, in relation to hatching of Anomala corpulenta, 502; method for extraction of sawfly cocoons from, 258, 259; wireworm populations in relation to particle size of, 73; types of, in relation to insects, 140, 214, 566, 683; effects of treatment of, with fertilisers against pests of conifers, 253; monograph on fauna of, 89; question of latency of insect virus in, 531; types of, in relation to infection of potato by viruses, 544; nitrogen

content of, unaffected by BHC, 678; types of, in relation to insecticide treatments, 277, 326, 416, 508, 572, 675; persistence of insecticides in, 77, 83, 119, 272, 293, 432, 483, 526, 527, 567, 572, 573, 577, 578; factors affecting distribution of insecticides in, 255, 475, 572, 573; determination of insecticide residues in, 145, 146, 339.

solana, Empoasca.

solani, Macrosiphum (Aulacorthum, Myzus).

solanifolii, Macrosiphum (see M. euphorbiae).

Solanum, experiments with Leptinotarsa decemlineata and alkaloids of, 318. Solanum bonariense, new Tortricid on, in

Argentina, 230.

Solanum chacoense, rôle of alkaloids in resistance of, to strains of Leptinotarsa

decemlineata, 318.

Solanum melongena, Corythaica cyathicollis on, in Brazil, 554; Leptinotarsa decemlineata on, in Bulgaria, 366; pests of, in India, 79, 497; Tetranychus desertorum on, in Japan, 452; grasshopper on, in Turkey, 351; Ostrinia nubilalis on, in U.S.S.R., 12; Micrutalis malleifera reared on, 636; new virus disease of, 345.

Solanum nigrum, Micrutalis malleifera transmitting curly-top virus of, in

Florida, 636.

Solanum polyadenium, rôle of chemical factors in resistance of, to Coleoptera, 608.

Solanin, experiments with, on resistance of Solanum to Leptinotarsa decemlineata, 318.

Solenopsis saevissima richteri (in U.S.A.), in pastures, 464; baits for, 464, 639; insecticides against, 145, 464, 483.

Solenopsis xyloni, in N. Carolina, 517. solitaria, Thaumetopoea.

solitarius, Apanteles.

Solomon Islands, new parasite of Agonoxena pyrogramma in, 293.

solstitiale, Amphimallon.

somniaria, Lambdina fiscellaria.

Sool, containing chlorfenson (q.v.), 440. Sooty Mould, associated with Hemiptera, 176, 233, 415, 587, 613.

sophiae, Colaphellus.

Sophora japonica, Eurytoma sophorae infesting seeds of, in Hungary, 306.

sophorae, Eurytoma.

Sorbus aucuparia, form of Hyponomeuta padellus associated with, in Finland, 194; Argyresthia conjugella on, in Germany, 9, 10, 441.

sordida, Hadena.

sorghicola, Contarinia.

Sorghum, investigation on first records of Contarinia sorghicola on, in African territories, 684, 685; Aphis sacchari on, in China, 624; aphids and virus disease of, in Italy, 80; pests of, in Kenya, 179; Lepidopterous stem borers of, in Nigeria, 292, 293; pests of, in U.S.A., 219, 421; Blissus leucopterus on, in Venezuela, 98; aphids and virus disease of, in Yugoslavia, 80; survival of Chloriona orizicola on, 132; in diet for Corcyra cephalonica, 626; recovery of, from insect attack by tillering, 179; destruction of débris of, in control of stem borers, 293; unaffected by insecticide residues in soil, 578; (stored), insects in, 554, 685; measures against pests in, 685; sorption of methyl bromide by, 554; effects of fumigants on germination of, 579.

Sorghum halepense, aphids and red-stripe disease of, in Yugoslavia, 80.

Sorghum vulgare, aphids and red-stripe disease of, in Yugoslavia, 80.

soudanensis, Hyperchalcidia.

Soy Bean, pests of, in China, 593, 594; mite fauna of, in Delaware, 414; pests of, in Japan, 452, 552; resistance to Epilachna varivestis in strains of, 413; experiments with Rivellia and, 609; review of pests and diseases of, 693. Soy Bean Pod Borer (see Leguminivora

glycinivorella).

Soy-bean Oil, in baits for fire ants, 464. Space Vehicles, experiments with various atmospheres in relation to use of cockroach in, 325.

spadix, Codiosoma (Pselactus).

Spain, Ceratitis capitata on apricot in, 98; Brevipalpus phoenicis on Citrus in, 299; virus disease of maize in 350; pests of olive in, 99; Leptinotarsa decemlineata on potato in, 681; forest pests in, 300, 355, 381, 551, 659, 660; insects attacking shade trees in, 353; Anobiids in furniture and building timber in, 355; Reticulitermes lucifugus in, 300; insect viruses in, 300, 550, 660; natural enemies of insects in, 313, 381.

Spanogonicus albofasciatus, on cotton in New Mexico, 409; soil treatments

against, 409.

Spanworm, Elm (see Ennomos subsignarius).

Sparganothis pilleriana, on vines in Austria, 248.

Sparganothis sulfureana, bionomics of, on cranberry in U.S.A., 337; sprays against, 337.

spartii, Cryptolestes.

Spearmint (see Mentha spicata).

speciosa, Diabrotica,

Spectrophotometry, use of, in determination of insecticides, 92, 145, 561.

spermotrophus, Megastigmus.

destroyed Sphaerophoria, aphids by species of, in New York, 401.

Sphagnum, use of, for protection of pupae of Tyria jacobaeae, 452.

sphaleropa, Eulia.

Sphenophorus callosus, bionomics of, on maize and Cyperus in Georgia, 152;

soil treatment against, 152.

Sphenoptera gossypii, bionomics of, on cotton in W. Pakistan, 176; dispersal of hibernating adults of, 176; tests of systemic insecticides against, 176.

Spiders, high-altitude collection of, 578; predacious on noxious insects, 24, 29, 153, 393, 402, 537, 638, 639.

Sphilochalcis, hyperparasite of Oiketicus kirbyi in Costa Rica, 638.

Spilochalcis albifrons, hyperparasite of Lepidoptera in California, 67.

Spilonota ocellana (on fruit trees), Quebec, 159; bionomics of, in U.S.S.R.,

Spilonota prognathana, on apple and hawthorn in U.S.S.R., 112; parasites of, 112.

Spilosoma lubricipeda (see Diacrisia). Spilosoma menthastri (see Diacrisia lubricipeda).

Spinach, aphids on, in California, 131, 480; virus diseases of, 17, 131, 480.

spiniferus, Aleurocanthus; Aphytis.

Spintherus linearis (parasitising Apion spp.), in Bulgaria, 540; in Rumania,

spiraecola, Aphis.

splendana, Cydia (Laspeyresia).

Spodoptera, species of, in Africa, 530.

Spodoptera exempta, survey of data on, in Africa, 530; in Kenya, 179, 530, 531, 686; in Tanganyika, 530, 531; on cereals, 530; on grasses, 179, 530, 686; losses of cattle attributed to, 530; phase differences and question of migration in, 530; characteristics and world distribution of, 507; parasites of, 531; occurrence and use of diseases of, 531, 686; insecticides against, 179, 530, 686.

Spodoptera exigua, on tomato in British Columbia, 529; in California, 66, 509; on celery, 509; parasitised by Hyposoter exiguae, 66; insecticides against, 476, 509.

Spodoptera frugiperda, on jute in British Guiana, 27; on maize in U.S.A., 275, 277, 512, 563; study of factors affecting adult life-span of, 563; parasitised by Hyposoter exiguae, 66; insecticides against, 27, 275, 277, 512.

Spodoptera mauritia (Lawn Armyworm), virus disease of, in Hawaii, 196.

Spondyliaspis, Glycaspis spp. erroneously placed in, 232.

Spotted Alfalfa Aphid (see Therioaphis

maculata).

Sprays, in relation to electrical charges on plants, 123; factors affecting toxicity of, to locusts, 657; pick-up of droplets of, by flying locusts, 291; effects of, on fauna, 12, 694; evaluation of programmes of, on arthropod fauna of apple, 376; timing of applications of, in relation to honey bees, 550; comparative effects of, on pests and their natural enemies, 8, 35, 36, 53, 55, 108, 158, 159, 222, 227, 279, 289, 312, 313, 434, 523, 603, 682; method of application in relation to effectiveness of, 383, 670; factors affecting patterns persistence of deposits of, 16, 21, 389, 574; residues in relation to pressure of, 570; evaporation losses from droplets in, 571; ground equipment for applying, 47, 85, 300, 413, 449, 630, 666, 667, 686; effects of equipment for applying, on potato yields, 333; equipment for applying in ships' holds, 195; applied from aircraft, 7, 55, 121, 122, 211, 220, 222, 276, 280, 336, 383, 395, 411, 413, 425, 455, 462, 530, 531, 540, 574, 659, 680, 683; viscosity of oils applied in, 470; question of phytotoxicity of wetting agents in, 463; portable chamber for experiments with, 248.

Spruce, pests of, in Canada, 24, 25, 107, 157, 163, 208, 457, 529; pests of, in Czechoslovakia, 142, 196, 246, 283, 601; Ips typographus on, in France, 606; pests of, in Germany, 44, 105, 107, 141, 251, 253, 682; Ernobius abietis on, in Italy, 136; Chermes spp. on, in U.S.S.R., 112; pests of, in U.S.A., 150, 208, 474; Formica spp. in relation to, 105, 615; relation of nitrogen content of, to feeding habits of Rhopalosiphum abietinum, 105; effects of fertilisers on pests of, 253; Coleoptera infesting timber of, 112, 253, 647, 648; penetration of timber of, by wood

preservatives, 647, 648.

Spruce, Engelmann (see *Picea engelmanni*). Spruce, Sitka (see Picea sitchensis).

Spruce, White (see Picea glauca and P. alba).

Spruce Budworm (see Choristoneura fumiferana).

spumarius, Philaenus (see P. leucophthalmus).

spuria, Pimpla.

sputator, Agriotes.

Squash, Diaphania nitidalis on, in N. Carolina, 471; varietal susceptibility of, to D. nitidalis, 471.

Squash, Banana, Tetranychus multisetis feeding on, 67.

stabilis, Orthosia (Taeniocampa).

stabulans, Lydella.

stabularis, Eulaelaps.

stali, Idiocerus.

stanleyi, Aphycus (Metaphycus).

staphyleae, Rhopalosiphoninus (Hyperomyzus, Myzotoxoptera).

Starch, in relation to infestation of timber

by Lyctus, 589, 622.

starki, Apanteles; Evagora (Recurvaria). Statistical Methods, for sampling Antestiopsis on coffee, 490; for morphometry of phases of locusts, 286; for describing distribution of stages of insects, 385; for analysis of experiments on insect control, 399; for estimation of toxicity of insecticides, 221.

Stauffer 1504 (see O,O-Dimethyl S-Phthalimidomethyl Phosphorodithio-

ate).

Stauffer R-1448 (see O,O-Diethyl S-Phthalimidomethyl Phosphorodithioate).

Stauffer R-1504 (see O,O-Dimethyl S-Phthalimidomethyl Phosphorodithio-

Stauffer R-1505 (see O,O-Diethyl S-Phthalimidomethyl Phosphorothioate).

Stauffer R-1571 (see O,O-Dimethyl S-Phthalimidomethyl Phosphorothioate) Stauffer R-2968 (see Diethyl Dithiopho-

sphorylethylcarbamate).

Staurochaeta albocingulata, parasitising Monoctenus juniperi in Holland, 302.

Stellaria media, as food-plant of stages of Calligypona pellucida in Finland, 102; Myzus ascalonicus overwintering on, in France, 605; arthropods on, in U.S.A., 477, 636; and virus diseases, 605, 636.

Steneotarsonemus fragariae (on strawberry), in Belgium, 445; in England, 431; in Poland, 545; in U.S.S.R., 675; bionomics of, 445, 545; Typhlodromus spp. predacious on, 545; measures against, 431, 545, 675; stimulation of development of, by parathion, 545.

Steneotarsonemus pallidus, auct. (see S.

fragariae).

Stenobracon deesae (see Glyptomorpha).

Stenodema laevigatum, on cereals and cotton in Turkey, 352.

Stenodiplosis bromicola, sp.n., bionomics of, infesting Bromus spp. in U.S.S.R., 542.

Stephanitis pyri, characteristics world distribution of, 507.

Stephanoderes hampei (on coffee), in Brazil, 555, 643; in Dutch New Guinea, 76; insecticides against, 555, 643, 644.

stercorosus, Geotrupes.

Sternocolaspis quatuordecimcostata, cotton and sugar-cane in Brazil, 646; toxicity of insecticides to, 646.

Sterols, utilisation of, by locusts, 5.

Stethorus pauperculus, predacious on Tetranychids in India, 627; toxicity of insecticides to, 627.

Stethorus punctillum, predacious on Tetranychus telarius in Holland, 682; effects of sprays on populations of, 682.

Sthenias grisator, bionomics of, on mango and other plants in India, 170.

sticticalis, Loxostege.

Stigmella malella, bionomics of, on apple in Italy, 309, 533, 534; measures against, 309, 534.

Stilpnotia salicis, in British Columbia, 126; in Spain, 300; in U.S.S.R., 115; on poplar, 126, 300; virus disease of, 300; natural enemies of, 126; Apanteles glomeratus not developing in, 115.

Straw, oviposition of Ochsenheimeria vacculella in, 596; termite attacking, in unbaked bricks, 180; treatment of,

against termites, 181.

Strawberry, Steneotarsonemus fragariae on, in Belgium, 445; pests of, in Britain 430, 431, 670, 671, 686; *Harpalus rufipes* on, in Bulgaria, 59; Hemiptera on, in Canada, 529, 686, 692; Hepialus lupulinus on, in France, 7; aphids on, in Germany, 106, 546; aphids on, in Holland, 192; pests of, in India, 595; S. fragariae on, in Poland, 545; pests of, in U.S.S.R., 360, 361, 675; pests of, in U.S.A., 23, 563, 570; *Haltica oleracea* on, in Yugoslavia, 242; identity of aphids on, 23; Tetranychus multisetis feeding on, 67; oviposition of T. telarius on, 547; Hemiptera and virus diseases of, 23, 106, 107, 192, 546, 686, 692; comparison of viruses of Fragaria vesca and, 441; insecticide residues on, 570, 671; effects of insecticides on, **360**, **570**, **675**. (See also *Fragaria vesca*.)

Strawberry Tarsonemid (see Steneotarsonemus fragariae).

Straws, Drinking, use of, as artificial nesting sites for bees, 411.

Streptococcus faecalis, larvae of Galleria mellonella infected by, 620.

striatella, Calligypona (Delphacodes).

striatigenas, Chelonus.

Striglina scitaria, bionomics of, on Sesbania bispinosa in India, 316; insecticides against, 317.

Strobane, against Anthonomus grandis, 408, 412, 515; against Heliothis zea, 406, 408, 412, 515; against Hemiptera, 408, 515; against Rhagoletis pomonella, 268; against Frankliniella, 408, 515; Tetranychus desertorum favoured by, 408; in dusts, 412; in sprays, 268, 408, 412, 515; review of toxicology of, 140; and DDT, 406, 412, 515.

strobi, Chermes (Pineus); Pissodes.

Strymon melinus, on cotton in Texas, 518; reared from capsules of Cienfuegosia sulphurea, 518.

Sturmia, parasitising Apamea anceps in

U.S.S.R., 363.

Sturmia bella, reared from Lozotaeniodes cupressana in Italy, 137.

Sturmia bohemica, established in Ontario, as parasite of Gilpinia hercyniae, 24.

Sturmia scutellata (parasitising Lymantria dispar), in Massachusetts, 400; in Yugoslavia, 371; persistence of, in sprayed areas, 400.

Sturmia vanessae (see S. bella).

Sturmiopsis parasitica, parasitic on Coniesta ignefusalis in Nigeria, 292.

Stylocryptus erythrogaster (see Endasys). Stylopalpia costalimai, sp.n., on groundnuts in Brazil, 145, 646; tests of dusts against, 145.

suasa, Hadena.

Subalpine Fir (see Abies lasiocarpa). Subcoccinella vigintiquatuorpunctata, on lucerne in Bulgaria, 60.

subfasciatus, Zabrotes.

subflava, Abbella.

subfusca, Hypsauchenia,

subfuscus, Athous; Troctes (Liposcelis).

subramanii, Euphysothrips. subrufa, Zygina (Empoasca).

subsignarius, Ennomos.

substrigosus, Rhopalotus.

subterranea, Feltia. Subteranean Clover (see Trifolium subterraneum).

subtilicornis, Diadromus.

succincta, Patanga.

Sucrase, rôle of, in nutrition of Agrotis orthogonia, 69.

Sucrose, in relation to feeding response of wireworms, 528; in diets for insects, 5, 52, 155, 208, 214, 290, 391, 404.

Suction Apparatus, for sampling insect populations, 224, 291.

Sudan, cotton pests in, 52, 94; first record of Contarinia sorghicola in, 684.

Sugar, excretion of, by aphids, 322; in bait for Piophila casei, 100; in baits and bait-sprays for Trypetids, 98, 99, 643; relation of, to oviposition by Tetranychus telarius, 547; in diets for insects, 633; effects of, in diet for Pseudotheraptus wayi, 29.

Sugar-cane, Aceria merwei on, in S. Africa, 83; Eumargarodes laingi on, in Australia, 265; Sternocolaspis quatuordecimcostata on, in Brazil, 646; pests of, in China, 623; pests of, in India, 119, 171, 297, 497, 498, 553, 628; pests of, in Jamaica, 289; pests of, in Louisiana, 216, 219, 280, 469; Aeneolamia varia saccharina on, in Trinidad, 332: pests of, in Venezuela, 98, 357; systematic list of pests of, in South-East Asia and Pacific area, 143; stool size of, in relation to sampling for A. v. saccharina, 332; effects of Aleurolobus barodensis on sugar content of, 498; estimation of losses of, caused by Diatraea, 357; rearing of D. saccharalis on, 214; virus diseases of, 81, 684; cultural practices in relation to biological control of pests of, 199, 200; treatment of setts of, 119, 297; toxicity of insecticides to, 297; effects of soil insecticides on injurious microfauna associated with, 553.

Sugar-cane Froghopper (see Aeneolamia varia saccharina).

Sugar-cane Termite (see Microtermes obesi).

Sugar Pine (see *Pinus lambertiana*).

sulcatus, Chelonus (Chelonella); Otiorhynchus.

sulcifrons, Sitona.

sulciventris, Musgraveia (Rhoecocoris).

Sulfotep, against Eotetranychus lewisi, 517; effects of, on colour of poinsettia, 517; in aerosols, 517.

Sulfoxide, as synergist; with phosphorus compounds, 63; with pyrethrins, 2, 3, 63.

sulfureana, Sparganothis,

Sulphenone (see p-Chlorophenyl Phenyl

Sulphone).

Sulphur, against Cydia pomonella, 20; premature spread of Gypsonoma dealbana caused by, 83; against Hemiptera, 560, 591, 645; against termite, 181; against Eriophyids, 83, 287, 298, 396, 409, 410, 560, 644, 645; against other mites, 28, 108, 120, 439; effects on beneficial Mirids of sprays containing, 672; as fumigant, 181; dusting with, 28, 83, 120, 298, 396, 439, 560, 644,

645; spraying with, 28, 120, 560, 644, 645; in mixed sprays, 20, 108, 287, 672; determination of particle size ot, 88; persistence of, in soil, 83; hazards of use of, 28; and arsenious oxide, 181; and rotenone, 560; and sabadilla, 591.

Sulphur Dioxide (as fumigant), toxicity of, to Coleoptera, 267; effects of, on seed germination, 579; in mixtures

with other fumigants, 579.

Sulphuric Acid, experiments with dieldrin and, in decomposition of froth masses

of Cercopids, 321.

Sulphur Pentafluoride, effects of, on toxicity and anticholinesterase activity of phosphorus and carbamate compounds, 634.

m-Sulphurpentafluorophenyl Methylcarbamate, toxicity of, to Musca domes-

tica, 634.

Sulphuryl Fluoride, fumigant toxicity of,

to Coleoptera, 267, 274.

O,O-Dimethyl Sumithion (see Methyl-4-nitrophenyl Phosphorothioate).

Sunflower (see Helianthus annuus).

Sunlight, effects of, on toxicity of BHC and DDT, 356. (See also Light.)

Supella supellectilium, infesting insect cultures in California, 339; baits for, 339.

supellectilium, Supella.

superans, Dendrolimus.

Superovokil, in sprays against Eotetranychus carpini, 183; constituents of, 183. suppressalis, Chilo.

Surinam (see Guiana, Dutch).

surinamensis, Oryzaephilus.

Survey Pegs, termite attacking, 76.

sutor, Monochamus.

suturalis, Araecerus; Sitona. swainei, Neodiprion.

Swede, Plutella maculipennis on, in Britain, 198; pests of, in Canada, 162, 529; pests of, in Finland, 104, 674; Hylemya floralis on, in Germany, 254; Ceutorhynchus pleurostigma on, in Holland, 673; pests of, in New Zealand, **75, 505, 506;** *H. brassicae* on, in U.S.A., 510, 568; resistance to Brevicoryne brassicae in varieties of, 167; in relation to oviposition by H. brassicae, 516; survey of insects on, 75, 506; treatment of, with nematode against pests, 162; seed treatment of, 104, 568, 674; effects of seed treatments on germination and development of, 675.

Sweden, pests of cereals in, 101, 354; Haplodiplosis equestris on Agropyrum repens in, 101; Tricholaba trifolii on red clover in, 85; forest pests in, 185, 426; Plutella maculipennis in, 198, 430; question of introduction of Tyria jacobaeae into Canada from, 260; virus disease of T. jacobaeae in, 260; beneficial insects in, 160, 161, 390, 391, 414, 493.

Sweet Clover (see Melilotus). Sweet Pepper (see Capsicum).

Sweet Potato, pests of, in Israel, 130; pests of, in U.S.A., 131, 133, 134; vectors of virus diseases of, 130, 131, 133, 134; experiments on transmission of internal cork virus of, 131.

Swietenia, Xyleborus morigerus on, in

Java, 625.

Swietenia macrophylla, attacked by Coptotermes niger in British Honduras, 76.

Swingfog Machine, 27, 536.

Switzerland, orchard pests in, 46, 86, 100, 663, 664, 682; Thomasiniana theobaldi on raspberry in, 552; Argyrotaenia pulchellana on vines in, 100; forest pests in, 46, 100, 186; Piophila casei infesting cheese-stores in, 100; native and introduced parasites of Quadraspidiotus perniciosus in, 664, 681; beneficial mites in, 682.

Sycamore (see Acer pseudoplatanus).

Sylepta derogata, characteristics and world distribution of, 507.

sylvatica, Tomosvaryella.

sylvaticus, Geotrupes (see G. stercorosus). Symplesis, parasitising Phyllocnistis in Aden, 80; parasitising Evagora stark in Canada, 586.

Synanthedon (see Aegeria).

Syngamia abruptalis, on mint in India, 424.

Syntomaspis druparum (see Torymus varians).

Syntomosphyrum, classification and new species of, 493.

Syria, Chloropsalta viridissima on vines in, 608; new egg-parasite of Eurygaster integriceps in, 655.

Syringopais temperatella, characteristics and world distribution of, 507.

Syritta pipiens, pollinating carrot in Utah, 341.

Syrphus, predacious on Aphis fabae in Bulgaria, 537.

Syrphus balteatus, predacious on aphids in Yugoslavia, 240.

Syrphus ribesii, predacious on aphids in Yugoslavia, 240.

Syrphus viridiceps, predacious on nymphs of Glycaspis baileyi in New South Wales, 233.

Systox (see Demeton).

Sytam (see Schradan).

T

tabaci, Bemisia; Thrips. tabaniformis. Paranthrene.

tabogae, Kalotermes.

Tachina larvarum (parasitising Lepidoptera), in Massachusetts, 400: U.S.S.R., 244, 245; persistence of, in sprayed areas, 400.

Tachina mella, parasitising Stilpnotia salicis in British Columbia, 126.

Tachinomyia similis, parasitising Stilpnotia salicis in British Columbia, 126.

Taeniocampa (see Orthosia).

Taeniothrips laricivorus, characteristics and world distribution of, 507.

Taeniothrips lefroyi, bionomics of, on broad beans in India, 499; insecticides against, 499.

Taeniothrips vaccinophilus, on blueberry in New Brunswick, 71; soil treatment

against, 71.

Tahiti, Brontispa longissima on coconut and other palms introduced into, from New Caledonia, 81.

Taiwan (see Formosa).

Talc, as carrier in dusts, 42; effect of, on toxicity of insecticide dusts, 118.

Tamarack (see Larix laricina).

Tamarind (Tamarindus indica), insects infesting dry fruits or seeds of, in India, 499.

Tanganyika, pests of coffee in, 652, 653; first record of Contarinia sorghicola in, 684; Nomadacris septemfasciata in, 98: Pheidole punctulata in. 87: Spodoptera exempta in, 530; beneficial insects in, 28, 652, 653.

Tantalum, Radioactive, Eurygaster inte-

griceps labelled with, 429.

Tanyrhynchus brevis, bionomics of, on vines and grasses in S. Africa, 650.

Tar Distillates (in sprays and dormant sprays), against aphids, 84, 677; against Coccids, 434, 676; against Psylla mali, 676, 677; against Coleoptera, 600, 647; against Lepidoptera, 15, 534; against mites, 60, 189, 676; effects of, on beneficial insects, 434, 678; penetration of timber by, 647; and DNC, 434, 676, 677; and lime-sulphur, **189**; and oil, **434**, **534**; creolin (q.v.)derived from, 675.

Tar Oil, treatment of timber with preparation containing, against Hylotrupes

bajulus, 254.

tardoides, Chermes (Adelges). tardus, Chermes (Adelges). Tarpaulins, use of, in fumigation, 215. tarsalis, Cephalonomia; Ctenicera.

Tarsonemoides confusus, sp.n., bionomics of, attacking Ips confusus in U.S.A., 386, 387.

Tarsonemoides gaebleri, attacking Ips typographus in Germany, 386.

Tarsonemoides plastographus, attacking Ips plastographus in California, 386, 387.

Tarsonemoides radiatae, sp.n., attacking

Ips spp. in U.S.A., 386, 387.

Tarsonemoides truncatus, attacking eggs of Ips in California, 386; characters of, 386; members of group of, 386.

Tarsonemus, saprophagous species of, on

soy bean in Delaware, 414.

Tarsonemus pallidus, auct. (see Steneotar-

sonemus fragariae).

Tarsonemus truncatus (see Tarsonemoides). Tasmania, Oncopera spp. on pastures in, 424.

tasmaniae, Aphodius.

tauricus, Agriotes litigiosus.

taxus, Aonidiella.

TDE (see DDD).

Tea, pests of, in Ceylon, 448, 449, 553, 554; Ceroplastes spp. on, in China, 362; Oligonychus coffeae on, in India, 288; Tetranychids on, in Japan, 452; Xyleborus morigerus on, in Java, 625; Ceroplastes spp. on, in U.S.S.R., 362; effects of cultural practices on beneficial insects and pests of, 553; question of use of DDT on, 449.

Tea-tussock Moth (see Euproctis pseudo-

conspersa).

Teak, pests of, in India, 620. (See also next entry.)

Tectona grandis (Teak), Xyleborus morigerus on, in Java, 625; resistance of timber of, to infestation by termites, 649.

tectonae, Perissopneumon.

Tedion, against Cydia pomonella, 21; against Tetranychids, 288, 289, 324, 445, 477, 550, 614, 633, 672, 677, 678, 685; effects of, on developing eggs of Oligonychus coffeae, 288, 289; resistance to, in strains of Tetranychids, 266, 510; toxicity of, to parathion-resistant strain of Tetranychus telarius, 278; against other mites, 439; effects of, on beneficial arthropods, 159; relation of structure to toxicity of, 445; in sprays, 159, 266, 288, 289, 324, 439, 477, 633, 672, 677, 678; as deposit, 288; persistence of, on apple leaves, 672; and Aramite, 477, 633; and DDT, 21; and malathion, 678; and oil emulsion, 510; and 2,4,5,4'-tetrachlorodiphenyl sulphide, 685.

Tedion V-18 (see Tedion).

Tedion V-18 Extra. 678.

Teepol, ineffective in sprays against Natada nararia, 448.

Tegea atropicta, predacious on termite in New South Wales, 33.

telarius, Tetranychus (Eotetranychus). telenomicidus, Ooencyrtus (Schedius).

Telenomus, egg-parasite of Dendrolimus superans in Mongolia, 116.

Telenomus dendrolimusi, bionomics of, parasitising Dendrolimus punctatus in China, 624; field use and large-scale rearing of, 624.

Telenomus dignus, parasitising Chilo sup-

pressalis in Japan, 451.

Telenomus gracilis, egg-parasite of Dendrolimus superans in Siberia, 13.

teleturga, Pansepta.

Telodrin (see 1,3,4,5,6,7,8,8-Octachloro-3a, 4, 7, 7a-tetrahydro-4, 7-methanoph-thalan).

Temelucha (see Cremastus).

Temnochila virescens, predacious on Dendroctonus barberi in New Mexico, 460.

Temnorhinus (see Cleonus). temperatella, Syringopais.

Temperature, in relation to use of electrical radiation against pests of stored cereals, 374; effects of: on aphids, 305, 320, 455, 462, 481, 482; on other Hemiptera, 173, 174, 601; on Coleoptera, 58, 94, 117, 127, 206, 294, 318, 319, 323, 440, 454, 486; on Diptera, 18, 30, 46, 97, 135, 139, 446, 494, 542; on Hymenoptera, 87, 125, 299, 551, 565; on Lepidoptera, 4, 52, 54, 55, 69, 78, 79, 149, 154, 190, 208, 236, 241, 276, 290, 291, 313, 368, 451, 452, 453, 465, 501, 597, 598, 599, 662, 665, 678; on development of Camnula pellucida, 158; on beneficial insects, 209, 257, 258, 299, 317, 394, 601, 678; on mites, 91, 361, 549, 583; on susceptibility of *Pectino*phora gossypiella to Bacillus thuringiensis, 427; on virus diseases of Lepidoptera, 335, 427; on plant viruses, 197; on toxicity and action of insecticides, 21, 202, 203, 218, 274, 356, 466, 471, 472, 579, 633, 676. (See also Coldhardiness and Heat.)

Tenebrio molitor, in cereal products in California, 420; effects of parental age on offspring of, 462; used in tests of toxicity and action of insecticides, 202, 584; larvae of, in diets for Formica polyctena, 105.

Tenebrio obscurus, in cereal products in California, 420.

tenebrioides, Zabrus. tenebrionis, Capnodis. Tenebroides mauritanicus, intercepted on oil-seed cake in Germany, 11; toxicity of fumigants to, 267.

tenella, Galerucella.

tenellus, Circulifer; Gelis.

tenerrima, Typhlocyba (Ribautiana).

Tennessee, Ennomos subsignarius on oak and hickory in, 418; Acarapis infesting honey bees in, 295.

Tent Caterpillar, Eastern (see *Malacosoma* americanum).

tenthredinis, Mesoleius (see M. aulicus).

tephragramma, Diparopsis.

Tephrina arenacearia, bionomics of, on lucerne in Bulgaria, 538, 539; insecticides against, 539.

TEPP, against aphids, 217, 324, 532; toxicity of, to Locusta migratoria, 172; against Tetranychus telarius, 217, 324; relation between diapause and resistance to, in Tetranychus telarius, 428; metabolism and mode of action of, 63, 213; in dusts, 217; in sprays, 217, 324, 532; HETP containing, 172; and Kelthane, 532.

terebella, Bracon.

Termite, Sugar-cane (see Microtermes obesi).

Termites, in W. Africa, 293, 294, 485, 495, 649, 650; in Australia, 33; accidental introduction of, into Britain, 321; in British Honduras, 76; in Canary Islands, 300; in Egypt, 180; in France, 649; in Germany, 649; in India, 119, 553, 627; in Italy, 649; in Java, 296; in Malaya, 76, 77; in Spain, 300; in Trinidad, 321; in Wisconsin, 416; in Zanzibar, 29; attacking sugar-cane, 119, 553; infesting other crops, 293, 294, 627; damaging trees, 76, 77, 180, 293, 294; attacking timber, 33, 76, 180, 296, 416, 485, 649, 650; resistance of timbers to attack by, 86, 180, 649, 650; attacking furniture and musical instruments, 296, 321; attacking straw in baked bricks, 180; review of damage to manufactured materials caused by, 489; habits of, 90, 495; effects of fungus on, 428; natural enemies of, 29, 33; treatment of concrete against, 416; nest and soil treatments against, 77, 119, 180, 576, 627; treatment of sugarcane setts against, 119; other measures against, 180, 293, 294, 296, 300, 485, 486, 553, 627; factors affecting toxicity of Thiodan to, 356; supplement to bibliography of, 300.

Tersilochus argentinensis, parasitising Listroderes in S. America, 353; characters of, 353.

Tersilochus parkeri, parasitising Listroderes in S. America, 353; introduced into Australia, 353; characters, redescription and forms of, 353.

tessulatus, Ctenomorphodes.

testudinea, Hoplocampa.

Tetanops myopaeformis (on beet), in Alberta, 402; in N. Dakota, 330; insecticides against, 330, 402.

2,4,5,4'-Tetrachlorodiphenyl Sulphide. toxicity of, to Tetranychids, 445, 685. 2,4,5,4'-Tetrachlorodiphenyl Sulphone (see Tedion).

2,4,5,4'-Tetrachlorodiphenyl Sulphoxide. toxicity of, to Tetranychids, 445, 685.

Tetrachloroethylene, repellency of, to Coleoptera, 270; as solvent for malathion, 270.

Tetrachloromethane, trunk injection with, against Saperda carcharias, 601.

6,7,8,9-Tetrahydro-5-H Tetrazoleazepine, effects of, in mixture with rotenone against Epilachna varivestis, 279.

Tetram (see Amiton Oxalate).

Tetramethyl Pyrophosphate, effects of other insecticides on anticholinesterase activity of, in cockroaches, 62.

Tetramorium caespitum, on beet in California, 419; aphids associated with, 419; chlordane spray against, 419.

Tetraneura ulmi (see Byrsocrypta). Tetranychus atlanticus, on soy bean in Delaware, 414; in France, 550; in Turkey, 444; on fruit trees, 444, 550;

acaricides against, 550.

Tetranychus bimaculatus (see T. telarius). Tetranychus cinnabarinus, on carnation in France, 550; in U.S.A., 147; bionomics of, 67; acaricides against, 550; genetic basis of resistance to acaricides in strains of, 147; use of, in tests of systemic insecticides, 277; T. multisetis treated as form of, 67.

Tetranychus desertorum, on cucurbits, Solanum melongena and soy bean in Japan, 452; on cotton in Texas, 408; demeton sprays against, 408; favoured

by insecticides, 408.

Tetranychus kanzawai, on tea and other

plants in Japan, 452.

Tetranychus lobosus (in U.S.A.), on cotton, 515; on soy bean, 414; acaricides against, 515.

Tetranychus marianae, new species allied to, 452.

Tetranychus mcdanieli, on apple in British Columbia, 394; sprays against, 394.

Tetranychus multisetis, bionomics of, on cucurbits in California, 67; experiments with other food-plants and, 67; treated as form of T. cinnabarinus, 67.

Tetranychus pacificus, in rearing media for predacious mites, 641; susceptibility or resistance to acaricides in strains of, 511, 512.

Tetranychus phaselus, sp.n., on bean in Japan, 452; T. marianae allied to, 452.

Tetranychus telarius, in Belgium, 688; in Brazil, 643, 645; in Britain, 431, 432; in British Guiana, 28; in Canada, 125; in France, 255, 256, 604; in Germany, 547, 688; in Greece, 352; in Holland, 193, 682, 688; in Hungary, 307; in Israel, 613, 684; in Italy, 663; in Japan, 36; in U.S.S.R., 242; in U.S.A., 19, 147, 217, 264, 272, 281, 324, 338, 477, 570, 573, 633; on beans, 125, 410, 547, 603, 604; morphological study of injury to bean leaves by, 107; on beet, 613; on cotton, 352, 410, 613, 643, 645; on fruit trees, 36, 242, 255, 256, 307, 410, 684; nature of damage to vines by, 663; on black currant, 432; on strawberry, 217, 570; on groundnuts, 613; on jute, 28; on maize, 281; on Mentha spp., 324; on tomato, 573; on carnation, 431; on *Ilex crenata*, 19; on roses, 193, 264, 272, 324, 477, 633; on greenhouse plants, 338, 547, 682; overwintering of, on annual weeds, 477; food-plant preferences of, 67; metabolism of plant pigments in, 108; effects of food-plant treatments on populations of, 256, 410, 516; effects of photoperiod on diapause in, 688; regulation of water loss by, 564; acetylcholine in, 40; esterase activity in eggs of strains of, 278; natural enemies of, 125, 522, 528, 682; tests and uses of acaricides against, 19, 28, 217, 264, 272, 281, 324, 352, 431, 445, 477, 478, 570, 603, 604, 613, 633, 643, 645, 684; susceptibility or resistance to acaricides in strains of, 193, 194, 272, 278, 516, 548, 613, 688; rôle of food-plant in resistance to acaricides in strains of, 548; relation between resistance to TEPP and diapause in, 428; ovicidal action of acaricides on parathionresistant strains of, 278; genetic basis of resistance to acaricides in strains of, 147, 602; effects of insecticide sprays on populations of, 36, 633; vapour toxicity of Dibrom to, 573; diverse effects of sesamex on toxicity of insecticides to, 64; question of use of endrin against, 432; used in tests of acaricides, 395, 444; weeding as measure against, 477; colour forms of, 548.

Tetranychus tumidus, sprays against, on cotton in Texas, 16.

Tetranychus urticae (see T. telarius).

Tetranychus viennensis (on fruit trees), in France, 550; in Greece, 239; in Hungary, 307; in Turkey, 177; in U.S.S.R., 242; bionomics of, 177; acaricides against, 550; favoured by DDT, 239. tetrarhodus, Capitophorus (Pentatrichopus).

tetrarhodus, Capitophorus (Pentatrichopus). Tetrastichus, egg-parasite of Anisota senatoria in Connecticut, 271; parasitising Laelia exclamationis in India, 79; as primary or secondary parasites of Planococcus citri in Israel, 614; classification and new species of, 493.

Tetrastichus amethystinus, parasitic on Leucoptera scitella in Italy, 136.

Tetrastichus atriclavus, parasitic on Lepidopterous stem borers in Nigeria, 292. Tetrastichus eurytus, parasitising Apion in

Bulgaria, 540.

Tetrastichus inunctus, parasitising Agevillea abietis in Germany, 252, 375; effects of insecticides on populations of, 375.

Tetrastichus migrator, parasitic on Leuco-

ptera scitella in Italy, 136.

Tetrastichus pseudoeceticola, parasitising Rhyacionia buoliana in Argentina, 230.

Tetrastichus silvaticus, relation of oviposition habits of Neodiprion swainei to egg-parasitism by, in Quebec, 459.

Tetrastichus tompanus, parasitising Apion in Rumania, 56.

Tetrastichus triozae, parasitising Trioza nigricornis in Italy, 436.

Tetrastichus turionum, established in Ontario as parasite of Rhyacionia buoliana,

Tettigella viridis, on wheat in China, 625. Tettigonia viridissima, experiments on transmission of tobacco and potato viruses by, 443; virus inactivated by homogenates of salivary glands of, 443.

texana, Anomis.

Texas, pests of Citrus in, 22, 221, 465, 564; cotton pests in, 16, 263, 269, 330, 334, 338, 403, 408, 409, 412, 413, 417, 473, 477, 517, 518, 567, 568, 576, 577, 629; development of cotton pests on Cienfuegosia in, 518; experiments with Pectinophora gossypiella and Gossypium hirsutum in, 269; Rhyacionia rigidana on pine in, 19; rice pests in, 329, 405; pests of other cereals and grasses in, 219, 281, 406; Aphis gossypii on melons in, 279; Thysanoptera on onion in, 477; pests of potato in, 510, 511; other insects in, 128, 219, 264, 406; natural enemies of pests in, 22, 279, 281, 338, 406, 564; aircraft collection of arthropods in, 577, 578.

Textiles, pests of, in Germany, 377; treatments of, against *Hofmannophila*

pseudospretella, 290; attraction of insects to stains of, 635; comparison of methods for measurement of effectiveness of treatments of, 553; use of, in rearing insects, 339. (See also Woollen Fabrics.)

textor, Hyphantria cunea.

Thanasimus formicarius, predacious on Ips typographus in Czechoslovakia, 142; method of rearing, 142.

Thanite, not toxic to Locusta migratoria,

173.

Thaumatomyia glabra, predacious on Pemphigus betae in Alberta, 469; effects of insecticides on populations of, 469.

Thaumetopoea pityocampa (on forest trees), in France, 423; in Italy, 615; in Spain, 381, 659; in Yugoslavia, 536; characteristics and world distribution of, 507; natural enemies of, 381, 615; use of bacterial and virus diseases against, 423, 536; endrin spray against, 659.

Thaumetopoea processionea, in France, 84; estimation of populations of parasites of, 84.

Thaumetopoea solitaria, on pistachio in Turkey, 609.

Thecodiplosis brachyntera, on pines in U.S.S.R., 116.

Thelairosoma gracile, parasitising Nephantis serinopa in India, 169.

Thelohania weiseri, sp.n., infecting Tortrix viridana in Germany, 105; experiments with other Lepidoptera and, 105.

theobaldi, Thomasiniana. theobroma, Distantiella.

Theocolax formiciformis, parasitising Ano-

bium punctatum in Britain, 489.

Therioaphis maculata (on lucerne in U.S.A.), varietal susceptibility of lucerne to, 629, 642; toxic and other effects of, 322; overwintering of, 631; factors affecting population changes in, 462; auxins in honeydew of, 629; auxin content of plants in relation to resistance to, 551; parasite of, 67; diseases of, 462, 570; insecticides against, 468; characters distinguishing Aphidius smithi from parasites of, 66.

Therioaphis riehmi (on Melilotus), in Canada, 529; in U.S.A., 469; tests of species and varieties of Melilotus for

resistance to, 469.

Theronia atalantae (parasitising Lymantria dispar), in Massachusetts, 400; in Yugoslavia, 54; persistence of, in sprayed areas, 400.

Thimbleberry (see Rubus parviflorus).

Thimet (see Phorate).

Thiodan, against aphids, 216, 333, 388, 412, 468, 631, 633, 637; ineffective against Eriosoma lanigerum, against other Hemiptera, 325, 570: against Coleoptera, 247, 325, 493, 539, 637, 641; against Diptera, 50, 101, 268; against Lepidoptera, 21, 154, 222, 269, 326, 333, 335, 337, 465, 503, 514, 538, 539, 631, 641, 644; factors affecting toxicity of, to termites, 356; against Thysanoptera, 477, 503; against Smin-thurus viridis, 34; against mites, 287, 431, 432, 628, 643, 671, 672; ineffective against Tetranychus telarius, 633; effects of, on beneficial insects, 312, 378, 627; ovicidal action of, 326; toxicity of, to fish, 304; in aerosols, 247; in dusts, 101, 216, 539, 637, 644; in sprays, 21, 34, 154, 216, 222, 269, 287, 325, 326, 333, 335, 336, 337, 388, 412, 431, 432, 465, 468, 477, 503, 514, 538, 539, 570, 631, 633, 641, 643, 671, 672; lack of persistence of deposits of, 389; seed treatment with, 50, 66; soil treatment with, 493; trunk and root treatments of peach trees with, 631; effects of, on plants, 503, 570, 672; residues of, on strawberry, 570; tolerance limit for, 570; analysis of residues of, 92; and fungicides, 50; and DDT, 631, 637.

Thiolutin, treatment of flannel with, ineffective against Anthrenocerus austra-

lis, 587.

Thiometon, against aphids, 59, 74, 188, 249, 312, 608; against Empoasca, 560; against Dacus oleae, 237, 238; against Heliothrips haemorrhoidalis, 257; against Neurotoma nemoralis, 541; against Saperda spp., 600; against mites, 60, 288, 289, 545, 628; effects of, on beneficial insects, 312, 560; in aerosols, 600; in sprays, 59, 60, 74, 188, 237, 238, 249, 288, 289, 312, 541, 545, 560, 600, 608; tests of systemic action and other methods of application of, 249, 257, 600; residues of, in olives, 238.

Thiram, seed treatments with mixtures of insecticides and, 103, 674.

Thisoicetrinus pterostichus, damaging grasses and field crops in Turkey, 351; BHC against, 351.

thomasi, Capitophorus (Pentatrichopus). Thomasiniana lavandulae, on lavender in France, 436, 606.

Thomasiniana theobaldi, bionomics of, on raspberry in Switzerland, 552; measures against, 552.

thompsoni, Aphidoletes.

thoracicus, Triaspis (Sigalphus).

Thorictodes heydeni, in stored wheat and flour in India, 315.

Threonine, in honeydew of Aphis pomi, 95.

l-Threonine, in relation to feeding response of wireworms, 528.

Thrips, on wheat in China, 625.

Thrips imaginis, characteristics and world distribution of, 507.

Thrips lini, bionomics of, on flax in Poland, 366.

Thrips nilgiriensis, associated with abnormal growth of coffee in India, 448.

Thrips tabaci, on tobacco in Bulgaria, 345; on onion in Texas, 477; transmitting virus diseases, 345; insecticides against, 345, 477.

Thuja, Thyridopteryx ephemeraeformis on, in New York, 19; activity of systemic insecticides in, 19.

thurberiella, Bucculatrix,

Thuricide, spore-preparation of Bacillus thuringiensis (q.v.), 269, 514.

thuringiensis, Bacillus.

Thymelicus lineola, introduced into Ontario, 161, 389; on grasses, 162, 389; on oats, 162; bionomics of, 162; parasites of, 162; diseases of, 389, 390.

Thyridopteryx ephemeraeformis, on ornamental evergreens in U.S.A., 19, 421; Bacillus thuringiensis against, 421; insecticides against, 19, 421.

Thysanus, species of, as hyperparasites of Planococcus citri in Israel, 614.

tibialis, Adelencyrtus; Chaetocnema; Phyllodecta.

TIFA, 550.

tiliae, Erythroneura; Typhlodromus (see T. pyri).

Timber, Coleoptera in, 31, 42, 90, 112, 142, 194, 356, 448, 489, 622, 647, 648; starch content of, in relation to infestation by Lyctus, 622; X-ray detection of Coleoptera in, 259; Drosophila virilis breeding beneath bark of, 623; termites attacking, 76, 296, 649, 650; resistance of, to infestation by termites, 649, 650; determination of resistance of, to termites, 86; method for determination of attraction of insects to, 340, 341; treatments of, 31, 42, 296, 448, 489, 647, 648, 649. (See also Building Timber.)

Timothy (see Phleum pratense).

tineivorus, Blattisocius.

Tineola bisselliella, damaging textiles in Germany, 377; basis of attractiveness of textile stains to, 635; use of, in determination of effectiveness of moth-proofing agents, 553; insects infesting cultures of, 339.

Tipula oleracea, characteristics and world distribution of, 507.

Tipula paludosa, cross-transmission studies with iridescent virus of, 690. tipuliformis, Aegeria.

Tmetocera (see Spilonota).

TMPP (see Tetramethyl Pyrophosphate).

Toadflax (see Linaria vulgaris).

Tobacco, Feltia subterranea on, in Brazil, 357; new virus disease of, in Bulgaria, 345; Myzus persicae on, in Ceylon, 81; Scrobipalpa heliopa on, in India, 236; aphids on, in Germany, 303; pests of, in Ontario, 388; Lepidoptera on, in Queensland, 503, 588; grasshopper on, in Turkey, 351; pests of, in U.S.A., 152, 406, 465, 566, 636; systematic list of pests of, in South-East Asia and Pacific area, 143; effects of, on development of Agrotis ypsilon, 346; Phthori-maea operculella on, 236; varietal susceptibility of, to pests, 366, 406; nicotine content of, in relation to feeding by *M. persicae*, 566; insects and virus diseases of, 81, 271, 303, 345, 443, 636, 690; salivary-gland homogenates of Tettigonia viridissima inactivating virus of, 443; injured by preparation of Bacillus thuringiensis, distribution of aircraft-applied insecticides on leaves of, 574; determination of insecticide residues in, 228.

Tobacco (Stored), pests of, 57, 58, 152, 236, 650; effects of redrying of, on populations of *Lasioderma serricorne*, 117; effect on symbiont of *L. serricorne* of nicotine in, 650; tainted by insecti-

cides, 503.

Tobacco Hornworm (see *Protoparce sexta*).

α-Tocopherol Acetate, in diet for Anthonomus grandis, 404.

Tomatin, experiments with, on resistance of Solanum to Leptinotarsa, 318.

Tomato, Aculus lycopersici on, in S. Africa, 83; pests of, in Brazil, 554, 556; Trialeurodes vaporariorum on, Britain, 198; Spodoptera exigua on, in British Columbia, 529; pests of, in Bulgaria, 345, 346; Heliothis spp. on, in Queensland, 588, 589; grasshoppers on, in Turkey, 351; pests and viruses of, in U.S.A., 271, 508, 573, 636, 640; Leptinotarsa decemlineata feeding on, 366; effects of ageing of foliage of, on fecundity of T. vaporariorum, 198; Thrips tabaci not surviving on, 345; and development oviposition Tetranychus telarius on, 547; insects and virus diseases of, 345, 636;

influence of, on resistance of T. telarius to acaricides, 548.

Tomato Juice, attraction of insects to fabric stains caused by, 635.

tomentosus, auct., Byturus (see B. urbanus). Tomicobia seitneri, parasitising Ips typographus in Czechoslovakia, 142; method of rearing, 142.

Tomosvaryella sylvatica, parasitising Macrosteles fascifrons in Manitoba,

551.

tompanus, Tetrastichus (Geniocerus).

tomyris, Pimpla.

Tortrix, species of, on cabbage and poplar in New Zealand, 75; tests of Bacillus thuringiensis against, 75.

Tortrix loeflingiana, on oak in Czechoslovakia, 683; parasites of, 683.

Tortrix postvittana (see Epiphyas). Tortrix viburniana (see Aphelia).

Tortix viridana (on oak), in Czechoslovakia, 105, 683; in Germany, 105,
106, 140, 248, 250; in Spain, 355, 381,
659; in U.S.S.R., 597, 598, 599;
bionomics of, 598, 599; temperature
in relation to egg development of, 599;
factors affecting populations of, 106;
Protozoa infecting, 105; natural
enemies of, 251, 381, 598, 683;
measures against, 355, 659; characteristics and world distribution of, 248,
507.

Torymus, parasitising Contarinia oregonensis in British Columbia, 386.

Torymus varians, on apple in Finland, 103.

Toumeyella numismatica, Coccophagus quaestor reared from, in Manitoba, 529.

Toxaphene, tests and uses of, against aphids, 498, 569, 633, 637; unsatisfactory against Therioaphis maculata, 468; against Mirids, 213, 408, 467, 478, 515; against other Hemiptera, 94, 554; against Curculionids, 16, 202, 275, 317, 338, 408, 412, 467, 515, 609, 637; resistance to, in strains of Anthonomus grandis, 218, 219; against other Coleoptera, 27, 173, 202, 316, 349, 644; against Noctuids, 16, 211, 269, 275, 277, 406, 408, 412, 514, 515, 569; against other Lepidoptera, 16, 269, 273, 353, 425, 555; resistance to, in strains of *Estig*mene acraea, 480; ineffective against Laelia exclamationis, 79; toxicity of, to Musca domestica, 202; toxicity of, to Orthoptera, 172, 174, 498; against termites, 77, 577, 627; against Thysanoptera, 151, 408, 409, 467, 515, 569;

against Vespula germanica, 422; toxicity of, to Schizotetranychus andropogoni, 628; Tetranychus desertorum favoured by, 408; bees unharmed by, 349; toxicity of, to predator of Tetranychids, 627; toxicity of, to fish, 304; review of toxicology of, 140; effect of temperature on toxicity of, 174, 202, 203; metabolism of, 64, 218; in aerosols, 27; in dusts, 64, 151, 349, 412, 514, 569, 637; in sprays, 16, 64, 79, 94, 174, 213, 220, 269, 273, 275, 338, 353, 408, 409, 412, 425, 468, 478, 515, 554, 555, 609, 632, 633, 644; applied from aircraft, 425, 569; as deposit, 316; seed treatment with, 65; soil treatment with, 77, 577, 578; persistence of, in soils, 578; treatment of nests with, 422, 627; treatment of selfing bags with, 277; tainting of vegetables and fruit by, 229; effects of, on plants, 65, 220, 555, 632, 633; residues of, in plants, 64, 578; residues of, in fat of cattle, 64; factors affecting persistence of residues of, 16; determination of, 88, 91, 620; and DDT, 220, 406, 408, 412, 467, 468, 478, 480, 514, 515, 569, 632, 633; question of synergism of, with DDT, 218, 219; and methylparathion, 409; and Sevin, 478.

Toxares deltiger, parasitising Anuraphis plantaginea in Holland, 673.

Toxoptera aurantii, on cacao in Dominican Republic, 144; on Citrus in Gaza Strip, 97; map of distribution of, 1; transmitting tristeza virus, 97.

Toxoptera citricidus, not reported from Aden, 80; and tristeza disease of Citrus in India, 80; world distribution of, 1,

507; survey of data on, **507**.

Toxoptera graminum (on cereals in U.S.A.), 160; varietal susceptibility of cereals to, 22, 219, 470, 551, 575, 629; comparison of oats and Phleum pratense as food-plants for, 585; plantgrowth regulators in relation to, 551, 629; transmitting yellow-dwarf virus, 134; destroyed by *Hippodamia convergens*, 281; trap for, 639; systemic insecticides against, 281; new biotype of, 470.

Toxoptera graminum phlei, subsp.n., bionomics of, on Phleum pratense in New Brunswick, 585.

transitans, Larvacarus. transitella, Paramyelois.

transversa, Drosophila.

transversoguttata, Coccinella.

Trapa bispinosa, pests of, in India, 595.

trapezalis, Marasmia.

Traps, for aphids, 29, 160, 547, 614, 639; for other Hemiptera, 23, 372; for Cephalcia, 601; for Coleoptera, 30, 155, 156, 341, 606, 671; for Drosophila, 265, 326, 327; for Trypetids, 280, 643; for other Diptera, 30, 319; for Lepidoptera, 55, 165, 368, 369; for other insects, 29, 30; effects of wind-speed catches in, 30; high-altitude collection of insects in, 199; use of female sex attractant in, 55, 368, 369. (See also Adhesive Traps and Light-traps.)

Trefoil, Birdsfoot (see Lotus corniculatus). Trematopygus scabriculus, parasitising Pachynematus scutellatus in Czecho-

slovakia, 246.

Trembling Aspen (see Populus tremuloides).

Trialeurodes, systemic insecticides against, on potato in Texas, 510.

Trialeurodes abutilonea, transmitting virus of sweet potato in Maryland, 133.

Trialeurodes vaporariorum (on tomato), in Britain, 198; in Ohio, 573; effects of ageing of tomato leaves on fecundity of, 198; vapour toxicity of Dibrom to, 573.

Triaspis caudatus, parasitising Apion in Bulgaria, 540.

Triaspis thoracicus, parasitising larvae of Bruchus pisorum in Bulgaria, 58.

Triboliocystis garnhami, parasitising laboratory cultures of Tribolium spp.,

Tribolium, in China, Kenya and Jamaica, 583.

Tribolium castaneum, in Gambia, 178, 179; in India, 685; in Kenya, 51; in Louisiana, 571; in stored cereals, 51, 118, 571, 685; in stored groundnuts, 178, 179, 583; dispersal of, 620; factors affecting fecundity and egg cannibalism of, 486, 621; discharge of repellent by, 621; Gregarines infesting cultures of, 582, 583; method of marking eggs of, 621; reared on wheatfeed, 486; tests and uses of insecticides against, 2, 3, 51, 118, 179, 316, 571, 685; factors affecting toxicity of insecticides to, 118, 314, 466; used in tests of insecticides, 202, 265, 470.

Tribolium confusum, in California, 339, 420; in Gambia, 178; in India, 500; in stored cereals and cereal products, 212, 270, 420, 500; in stored groundnuts, 178; infesting insect cultures, 339; survival of, in nitrogen atmospheres, 416; experiments on competition between Latheticus oryzae and, 89; Gregarines infesting cultures of, 582; bait for, 339; tests and uses of insecticides against, 4, 212, 213, 267,

270, 500, 554, 573, 603; factors affecting toxicity of fumigants to, 274, 466; repellence of insecticides to, 270.

Tribolium destructor, effects of crowding on larval development of, 184; diet for, 184.

Tribolium madens, Gregarines infesting cultures of, 583.

1,2,3-Tribromopropane, vapour toxicity of, to *Phylloxera vitifoliae*, **572**.

Trichiocampus viminalis, bionomics of, on poplars in Quebec, 392; virus and other mortality factors of, 392.

Trichlorobenzene, trunk applications of, against Aegeria exitiosa, 642; toxicity

of, to peach, 642.

2,4,5-Trichlorodiphenyl Sulphide, Sulphone and Sulphoxide, toxicity of, to winter eggs of *Panonychus ulmi*, 685.

1,1,1-Trichloroethane, factors affecting fumigant toxicity of, to *Tribolium*

confusum, 274.

Trichlorphon, against aphids, 279, 549, 559, 667, 668; ineffective against Eumargarodes laingi, 265; against other Hemiptera, 382, 639; against Coleoptera, 176, 493, 600; against Noctuids, 46, 211, 530, 538, 539, 559, 644; against Tortricids (sens.lat.), 9, 337, 558, 646; against other Lepidoptera, 46, 309, 503, 514, 534, 539; against *Pegomyia betae*, 443, 549; against Trypetids, 47, 98, 99, 135, 237, 238, 240, 268, 309, 311, 668, 676; against Sminthurus viridis, 34; against Thysanoptera, 348, 569; question of toxicity of, to Schizotetranychus andropogoni, 628; beneficial insects destroyed by, 627, 639; metabolism of, 64, 213; in dusts, 348, 644; in sprays, 9, 34, 46, 47, 98, 135, 176, 237, 240, 279, 309, 311, 337, 503, 514, 530, 534, 538, 539, 549, 558, 559, 569, 600, 646, 668, 676; in bait-sprays, 98, 99; applied from aircraft, 530; soil treatment with, 493; tests of systemic action and other methods of application of, 176, 265, 600; residues of, in olives, 238; and DDT, 676; and methyl-demeton, 549; booklet on properties and use of, 541. trichodactyla, Hylemya.

Trichogramma, egg-parasite of Cydia

pomonella in China, 592.

Trichogramma cacoeciae, parasitising Lepidoptera in Germany, 381, 681; spatial distribution of, in apple trees, 681; method of rearing, 381.

Trichogramma evanescens, in China, 623, 624; parasitising sugar-cane borers,

623; parasitising *Telenomus dendrolimusi*, 624, 625; experiments on parasitism of eggs of *Sitotroga cerealella* by, 128; method of rearing, 623.

Trichogramma japonicum, egg-parasite of Chilo suppressalis in Japan, 451.

Trichogramma minutum (egg-parasite of Lepidoptera), in Argentina, 144; in Canada, 26, 27, 74, 457; in Connecticut, 271.

Tricholaba trifolii, on red clover in

Sweden, 85.

Tricholoma matsutake, Diptera on, in Japan, 623.

Trichomalus cristatus, parasitising Oscin-

ella frit in U.S.S.R., 12.

Trichoplusia ni, in Canada, 161, 163; in India, 424; in U.S.A., 66, 172, 219, 222, 269, 284, 328, 335, 404, 413, 477, 508, 514, 569, 630, 639; on Capsicum, 630; on cotton, 413, 477, 639; on crucifers, 161, 163, 172, 222, 269, 328, 335, 404, 424, 508; relation of populations of, to other cabbage pests, 172; on lettuce, 514, 569; on mint, 284; processed vegetables contaminated by, 222; fluctuations in populations of, 335; light-traps for, 477; light-trap catches of, 219, 335; natural enemies of, 66, 163, 401, 639; virus diseases of, 284, 328, 335, 401, 426, 639; use of viruses against, 404; spore preparation of Bacillus thuringiensis against, 269, 328, 569; use of larvae of, in determination of activity of preparations of B. thuringiensis, 342; insecticides against, 161, 222, 269, 413, 476, 508, 514, 516, 569, 630; resistance to insecticides in larvae of, 222; diet for, 342.

Trichopria, species of, as secondary parasite of Ostrinia nubilalis in U.S.S.R.,

246.

tridens, Ips.

tridentinus, Nothroholaspis.

Triecphora mactata (see Cercopis sanguinolenta).

trifasciatus, Lounsburyia (Coccophagus); Hadrotettix.

trifenestrata, Cricula.

trifolii, Apion; Tricholaba.

Trifolium hybridum, Sitona spp. on, in Finland, 41, 102; S. hispidulus and fungi associated with root damage to, in Oregon, 275; Apion seniculum on, in U.S.S.R., 114; transmission of stolbur virus to, by Aphrodes bicinctus, 304.

Trifolium incarnatum, Bruchidius perparvulus on plants and stored seed of, in Italy, 182. Trifolium medium, Apion virens on, in U.S.S.R., 114.

Trifolium pratense, Apion spp. on, in Bulgaria, 540; pests of, in Canada, 71, 453, 454; Apion spp. on, in Czechoslovakia, 445; Sitona spp. on, in Finland, 41, 102; Eurytoma platyptera infesting seeds of, in Hungary, 306; Apion spp. on, in Rumania, 56; Tricholaba trifolii on, in Sweden, 85; A. virens on, in U.S.S.R., 114; pests of, in U.S.A., 134, 323, 417, 564; Apion spp. on, in Yugoslavia, 32; effects of infestation by Apion on seed yields of, 32; influence of time of harvest of, on populations of Dasyneura leguminicola, 71; root rot of, 323; yields of, in relation to applications of insecticides, 323; insecticide residues on, 324, 417.

Trifolium repens, Aphrodes bicinctus on, in Canada, 686; A. bicinctus on, in Czechoslovakia, 304; Sitona spp. on, in Finland, 41, 102; dodder transmission of virus to Fragaria vesca from, in Germany, 441; invasion of pasture by, under influence of Hyperodes bonariensis in New Zealand, 506; Apion seniculum on, in U.S.S.R., 114; insects and virus diseases of, 304, 441.

686.

Trifolium subterraneum, pests of, in Australia, 34, 504; experimental infection of, by wilt virus of broad bean, 34.

Trigona, habits of, damaging crops in Dutch Guiana, 143.

Trigona amalthea, 143.

Trigona hyalinata branneri, 143.

Trigonotylus ruficornis, on cereals in Sweden, 354.

trigonum, Eutrombidium (see E. rostratum).

Trimedlure, attractiveness of, to *Ceratitis* capitata, 384.

Trinervitermes, in W. Africa, 495; foraging and feeding habits of, 495.

Trinervitermes ebenerianus, soil treatment against, in Nigeria, 293, 294.

Trinervitermes heimi, treatment of sugarcane setts against, in India, 119.

Trinervitermes indicus, treatments of nests of, in India, 627.

Trinidad, *Rhynchophorus palmarum* as vector of nematode causing red ring disease of coconut in, 575; *Aeneolamia varia saccharina* on sugar-cane in, 332; *Cryptotermes brevis* in, 321; predacious bird introduced into Bermuda from, 229.

Trioxys pallidus (parasitising aphids), established in California, 151; in

France and Persia, 151; bionomics of, 151; question of identity of, 151.

Trioxys utilis, introduced into California, 67; bionomics and sex ratio of, parasitising Therioaphis maculata, 67; parasitism of, 68; question of identity of, 151.

Trioza nigricornis, bionomics and parasite of, on onion in Italy, 436; sprays against, 436, 437.

triozae, Tetrastichus.

tripunctata, Drosophila.

Trisetacus pini, on Pinus strobus in Canada, 529.

Tristania, starch content related to infestation of species of, by Lyctus brunneus in Queensland, 589.

Tristania conferta and suaveolens, 589.

tristis, Ambeodontus.

Trithion, against aphids, 217, 279, 468, 643, 645; ineffective against *Phylloxera vitifoliae* in galls, 668; against other Hemiptera, 395, 515; against Coleoptera, 338, 515, 644; against Diptera, 268, 402; against Lepidoptera, 21, 421, 466, 515; against Thysanoptera, 477; against mites, 217, 281, 515, 643, 645; cross-resistance to, in strain of *Tetranychus pacificus*, 512; toxicity of, to bees, 412; effects of, on beneficial arthropods, 159; metabolism of, in insects, 64; in dusts, 217; in granules, 402; in sprays, 159, 217, 268, 279, 395, 421, 466, 468, 477, 515, 643, 644, 645; soil treatment with, 402; persistence of, on apple leaves, 672; effects of, on plants, 466, 643, 645; and DDT, 268; toxicity of DDT increased by, 21.

tritici, Contarinia; Frankliniella; Haplo-

thrips; Hydrellia; Sciara.

Triticum aestivum, factors affecting resistance of varieties of, to Hemiptera, 72, 219, 527. (See also Wheat.)

Triticum durum, factors affecting resistance of varieties of, to Hemiptera, 72, 219, 527. (See also Wheat.)

Triticum vulgare (see T. aestivum).

Triton X-100, 172, 463.

Triton X-114, 412.

Triton X-116, 668.

trivialis, Calocoris.

Troctes subfuscus, in stored grain in Canada, 154.

Trogoderma, survey of data on, in stored products in Australia, Egypt and U.S.A., 18; key to larvae of, 18.

Trogoderma granarium (in stored foodstuffs), in Britain, 295; not found in Gambia, 178; in Germany, 11; in Pakistan, 295; in Sicily, 300; in U.S.A., 17, 18; studies on feeding, growth,

respiration and behaviour of stages of, 6; diapause in, 6, 294; insecticides against, 212, 213.

Trogoderma inclusum, distribution of, in U.S.A., 18.

Trogoderma parabile, infesting stored cereals and feed mills in Canada, 529; distribution of, in U.S.A., 18.

Tropaeolum, Aphis fabae on, 603.

Tropidophryne palestinensis, insect described as, parasitising Planococcus citri in Israel, 614.

Trout, Rainbow, toxicity of insecticides

to, 304.

truncatella, Litomastix (Copidosoma). truncatus, Tarsonemoides (Tarsonemus). truncorum, Formica.

Trybliographa rapae, parasitising Hylemya brassicae in Holland, 39.

tryoni, Dacus.

Trypodendron lineatum (infesting conifers), in British Columbia, 207, 423, 424, 459; in Germany, 44; flight behaviour and log selection by, 424, 459, 460; overwintering habits of, 207, 423; traps for, 459, 460; tests of attractants for, 44.

Trypopitys carpini, in building timber in

Poland, 142.

Tryporyza incertulas (on rice), in China, 501; in Formosa, 328; influence of temperature on development of, 501; characteristics and world distribution of, 507.

Tsuga heterophylla, pests of, in British Columbia, 395, 459; attraction of Trypodendron lineatum to logs of, 459.

tsugae, Scolytus.

tsuneonis, Dacus. tuberis, Epitrix.

Tubes, Micro-capillary, topical applica-

tion of insecticides with, 3.

Tulip, Anuraphis tulipae on, in British Columbia, 529; Rhopalosiphoninus staphyleae tulipaellus overwintering on, in France, 605; breaking virus of, 529. tulipae, Aceria; Anuraphis (Dysaphis).

tulipaellus, Rhopalosiphoninus (Hyperomyzus, Myzotoxoptera) staphyleae.

tumidus, Asolcus; Tetranychus.

Tung (see Aleurites spp.).

Tunisia, Bruchidius incarnatus in, 249; parasite of Anacridium aegyptium in, 313.

turcicus, Cryptolestes. turionellae, Pimpla. turionum, Tetrastichus.

Turkey, cereal pests in, 43, 297, 351, 352, 485, 655, 656; pests of cotton in, 82, 177, 237, 352, 495, 552; orchard pests in, 82, 177, 237, 300, 444, 495; pests of

olive in, 81, 177; pests of hazel in, 82, 83; pests of pistachio in, 81, 609; pests of vegetables in, 82, 351, 352; pests of cedar in, 355; grasshopper on grasses and tobacco in, 351; pests of stored products in, 236, 609; Galleria mellonella injurious in bee-hives in, 551; beneficial insects in, 43, 352, 485, 655.

Turnip, Plutella maculipennis on, in Britain, 198; pests of, in Finland, 674; survey of insects on, in New Zealand, 75, 506; Dalopius pallidus infesting, in Nova Scotia, 73; Hylemya spp. on, in U.S.A., 510, 568; in relation to oviposition and rearing of H. brassicae, 39, 516; experiments with aphids and mosaic virus of, 197, 250; insecticidal properties of constituent of, 483; seed treatment of, 568, 674; toxicity of seed treatments to, 104, 675; question of heptachlor residues in, 479.

Turpentine, fumigant toxicity of, to

Dendroctonus spp., 223.

tutela, Rhopalicus.

Two-spotted Mite (see Tetranychus telarius).

Tydeus caudatus, on vines in Italy, 183. Tydeus interruptus, in stored grain in Canada, 154, 529.

Typhlocyba comes (see Erythroneura). Typhlocyba rosae, insecticides against, on loganberry in British Columbia,

395.

Typhlocyba tenerrima, insecticides against, on loganberry in British Columbia, 395.

typhlocybae, Psilodryinus.

Typhlodromus (predacious on other mites), in Delaware, 414; in Nova Scotia, 259; in Switzerland, 682; unaffected by nutrition of host food-plant, 259.

Typhlodromus aberrans, predacious on Eotetranychus carpini in Italy, 183.

Typhlodromus andersoni, predacious on other mites in Ontario, 522.

Typhlodromus bakeri, characters distinguishing T. caudiglans from, 522.

Typhlodromus bellinus, cultures of Panonychus ulmi destroyed by, 522.

Typhlodromus caudiglans, sp.n., bionomics of, predacious on other mites in California, 521, 522; in Canada, 522; factors affecting development of, 522; characters of, 522; misidentified as T. rhenanus, 522.

Typhlodromus conspicuus, predacious on other mites in Ontario, 522.

Typhlodromus fallacis, cultures of Tetranychus telarius destroyed by, 522. Typhlodromus finlandicus, predacious on

apple pests in Nova Scotia, 159; effects of sprays on populations of, 159. Typhlodromus graminis, species related to, predacious on Steneotarsonemus fragariae in Poland, 545.

Typhlodromus guatemalensis, 522.

Typhlodromus hibisci, in California, 378, 641: predacious on other mites, effects of diet on reproduction of, 641; method of rearing, 641.

Typhlodromus limonicus, predacious on

other mites in California, 378.

Typhlodromus longipilus, in Holland, 682; in Ontario, 522; predacious on other mites, 522, 682; effects of sprays on populations of, 682.

Typhlodromus occidentalis, predacious on Tetranychus telarius, 528; effects of prey density on oviposition by, 528.

Typhlodromus pyri (predacious on other mites), in Austria, 302; in Britain, 379; in Germany, 319, 439; in Ontario, 522; factors affecting development and oviposition by, 122, 319; apple pollen as food for, 319; effects of sprays on populations of, 439.

Typhlodromus reticulatus, predacious on Steneotarsonemus fragariae in Poland,

Typhlodromus rhenanus (predacious on other mites), in Britain, 522; question of occurrence of, in California, 522; in Canada, 159, 454, 522; effect of sprays on populations of, 159; T. caudiglans misidentified as, 522.

Typhlodromus tiliae (see T. pyri). Typhlodromus vitis (see T. aberrans). typica, Chermes (Dreyfusia) piceae.

typographus, Ips.

Tyria jacobaeae, introduced into Australia, 452; in Britain, 452; introduced into Canada, 260; in Ireland, 260; in Italy, 452; in Sweden, 260; control of ragwort by, 260, 452; food-plant range of, 260; effects of cold treatment on diapause and viability of, 452, 453; diseases of, 260; question of strains of, 453.

Tyroglyphus farinae (see Acarus siro).

Tyrophagus, saprophagous species of, on soy bean in Delaware, 414; in parasitised eggs of Caligo eurilochus in Ecuador, 268.

Tyrophagus longior, on cucumber in British Columbia, 529.

Tyrophagus putrescentiae, in U.S.A., 518, 573; in stored grain, 518; on mushrooms, 573; food preferences of, 518; factors affecting development of, 583; reared on Aspergillus, 583; Dibrom ineffective against, 573.

1-Tyrosine, in relation to feeding response of wireworms, 528.

UCC 8305 (see P-Chloro-2,4-dioxa-5methyl-P-thiono-3-phosphabicyclo-[4.4.0]decane).

Udea ferrugalis, bionomics of, on cabbage in Bulgaria, 60; sprays against, 61.

Ufens, species of, parasitising leafhoppers in California, 630.

Uganda, Platypodid attacking Baikaea insignis minor in. 496; Antestiopsis spp. on coffee in, 490; cotton pests in, 652; first record of Contarinia sorghicola in, 684.

ugandae, Cryptolestes.

uhleri, Pilophorus.

ulmi, Byrsocrypta (Tetraneura); Panonychus (Metatetranychus).

Ulmus americana, Scolytus multistriatus on, in Connecticut, 641.

Ulmus pumila, Ambrostoma quadriimpressum on, in China, 593.

Uloma, infesting dry fruits of tamarind

in India, 499. Ultraviolet Light, types of, in rearing of Anthonomus grandis, 407; attraction of Hoplocampa testudinea to, 666; in traps for Lepidoptera, 334, 476, 477, 630; fluorescence of fractions of female sex attractant of Lymantria dispar under, 370; effect of, on susceptibility of Galleria mellonella to polyhedrosis virus, 427; inactivation of plant virus by, 133; for examination of deposits of labelled insecticides, 221.

ulula, Dyspessa.

Unaspis yanonensis, identity of parasite of, in China, 491.

unctuosus, Lygus (Liocoris). undecimpunctata, Diabrotica. undulata, Clastoptera.

unicostata, Monosteira.

unicus, Praon.

Union Carbide 10854 (see m-Isopropyl-

phenyl Methycarbamate).

Union of Soviet Socialist Republics, pests of beet in, 15, 191, 545, 609; pests of cabbage in, 114, 115, 244, 245, 362, 542; Ostrinia nubilalis on egg-plant in, 12; flax pests in, 15; potato pests in, 15, 543, 681; pests of cereals in, 12, 13, 14, 15, 113, 243, 245, 343, 344, 345, 362, 363, 446, 447, 543, 595, 596; pests of grasses in, 191, 345, 363, 447, 542, 596; pests of clovers in, 114, 686; phyllody virus of clover in, 686; cotton pests in, 15,

113; O. nubilalis on hemp in, 14, 245; orchard pests in, 13, 15, 112, 191, 242, 343, 359, 360, 361, 362, 543, 597; pests of soft fruits in, 360, 361, 362, 595, 596, 675; pests of vines in, 15; Ceroplastes spp. on tea in, 362; forest pests in, 12, 13, 111, 112, 114, 115, 116, 190, 191, 242, 243, 244, 248, 344, 354, 355, 368, 597, 598, 599, 620; Eulecanium corni on ornamental trees in, 13; rose pests in, 355, 361; Lepidoptera on rosaceous plants in, 112, 113; insect pests of field crops in region of, 40; Ptinus spp. in stored foodstuffs in, 344; weevils infesting timber in, 112; locusts in, 13, 15; Plutella maculipennis in, 87, 198, 430; Lamellicorns in, 111, 541; other insects in, 62, 192, 597; forecasting abundance of pests and diseases of agricultural crops in, 15; Acarapis woodi infesting honey bees in, 296; diseases of insects in, 115, 359, 363, 447; natural enemies of pests in, 12, 13, 15, 87, 112, 114, 115, 160, 161, 190, 192, 243, 244, 245, 246, 344, 362, 363, 592, 596, 597, 598, 664; question of introduction of parasites of Ceroplastes into, 362; suggestions on biological control in, 596; nomenclature of Eurytoma spp. in, 564; transliterated list of names of insecticides and other agricultural chemicals used in, 214.

unipuncta, Pseudaletia. unipunctata, Gynandrophthalma.

unispinosus, Scolytus. United States of America, pests of cereals and grasses in, 71, 134, 160, 219, 265, 333, 398, 405, 420, 421, 520; virus diseases of cereals in, 71, 134; cotton pests in, 147, 218, 334, 511, 568; Pectinophora gossypiella on other malin, 511; Metcalfa vaceous plants pruinosa on Citrus in, 465; pests of other orchard fruits in, 18, 71, 147, 403; virus diseases of fruits in, 71; pests of nuts in, 147; pests of vines in, 147; Amphorophora rubi on raspberry in, 132; aphids on strawberry in, 23; lucerne pests in, 92, 400; pollinators of lucerne in, 411, 565; Therioaphis riehmi on Melilotus in, 469; characters of Eurytoma spp. on leguminous forage crops in, 564; forest pests in, 19, 20, 73, 74, 121, 122, 129, 208, 209, 210, 411, 418, 458, 474, 495, 513, 532, 538; insects and diseases of elm in, 217, 478; pests of tobacco and other Nicotiana spp. in, 406, 566; disinfestation of tobacco in, 117, Tetranychids on greenhouse plants in, 147, 517; virus diseases of other plants in, 17, 23, 132, 133, 271;

list of plant pests intercepted in, 299; mites infesting stored products in, 17, 18, 518; Aphis craccivora misidentified as A. medicaginis in, 262; fire ants in, 145, 483; wild bees in, 140, 565; Orthoptera in, 406, 521, 567, 575; dispersal of Popillia japonica in, by aircraft, 418; other insects in, 92, 306, 377, 513, 565; Acarapis spp. and honey bees in, 342, 504, 505; natural enemies of insects in, 73, 74, 317, 377, 379, 400, 465, 565, 664; fish in relation to the use of insecticides in, 262; tolerance limits for insecticides in, 64, 323, 326; surveys of data on arthropods not known to occur in, 506-508. (See also under individual States.)

urbanus, Byturus.

Urea, diapause in eggs of Acheta commodus prevented by, 296.

Urena lobata, Earias on, in India, 37. Urine, attraction of insects to fabric stains caused by, 635.

urozonus, Eupelmus.

Ursolic Acid, influence of, on adherence of spray deposits to apple, 21. urticae, Tetranychus (see T. telarius).

Uruguay, new Pyralid on sunflower in, 230; Acarapis woodi infesting honey bees in, 296; parasites of Rhyacionia buoliana in, 230.

Utah, insects pollinating carrot in, 341; Acarapis sp. infesting honey bees in, 295.

utilis, Trioxys.

vaccinii, Frankliniella.

Vaccinium (Blueberry), Thysanoptera on, in New Brunswick, 71; Hendecaneura shawiana on, in New York, 567.

vaccinophilus, Taeniothrips.

vacculella, Ochsenheimeria. Vacuum Cleaner, use of, in tests of vapour toxicity of dichlorvos, 226.

vagabunda, Mordwilkoja.

valens, Dendroctonus.

Valeric Acid, as attractant for Popillia japonica, 577.

2-n-Valeryl-1,3-indandione, experiments with synergists and, against Alphitobius laevigatus, 201, 687. Valgus hemipterus, on apple in Turkey,

237.

validirostris, Pissodes.

Valine, in saliva of Aphis pomi, 95. Valone (see 2-Isovaleryl-1,3-indandione). Vanessa cardui, parasitised by Hyposoter

exiguae in California, 66.

vanessae, Sturmia (see S. bella).

vaporariorum, Trialeurodes.

Vaporisers, for laboratory tests with dichlorvos, 226.

varia, Aeneolamia.

variabilis, Hypera (Phytonomus).

varians, Torymus.

variator, Bracon.

varicornis, Leptocorisa.

variegana, Argyroploce.

variegata, Meromyza.

variegatum, Zagrammosoma (Atoposoma). varipes, Apion.

varium, Apion (Piezotrachelus).

varivestis, Epilachna.

Vasates cornutus (see Aculus).

Vasates lycopersici (see Aculus).

Vasates mckenziei, on wheat and Agropyrum repens in Ontario, 71, 523; transmission of plant viruses by, 71, 523.

Vasates schlechtendali, on apple in Nova Scotia, 259; effects of plant nutrition on populations of, 259.

vassilievi, Asolcus.

vastatrix, Phylloxera (see P. vitifoliae).V-C 13 (see O,O-Diethyl O-2,4-Dichlorophenyl Phosphorothioate).

velutinana, Argyrotaenia.

Venezuela, *Blissus leucopterus* on cereals and grasses in, 98; pests of sugar-cane in, 98, 357.

venosatus, Proceras (Diatraea).

ventralis, Pegomyia.

ventricosus, Pyemotes.

Venturia inaequalis, effects of sprays against, on predators of apple pests in Nova Scotia, 159.

venturii, Diapetimorpha.

verbasci, Anthrenus.

Vermiculite, in rearing medium for *Operophtera brumata*, **258**; as carrier for insecticides, **277**.

versicolor, Meteorus.

verticalis, Gilpinia (Diprion).

Verticillium alboatrum, associated with Sitona hispidulus on alsike clover in Oregon, 275.

vespiformis, Aegeria (Synanthedon),

Vespula arenaria, predacious on larvae of Neodiprion swainei in Quebec, 73.

Vespula germanica, history of establishment of, in New Zealand, 421-423; bionomics and adult habits of, 422; food preferences of, 422; baits for, 422; measures against, 422, 423.

Vespula maculata, predacious on larvae of Neodiprion swainei in Quebec, 73.

Vespula norwegica norvegicoides, predacious on larvae of Neodiprion swainei in Quebec, 73.

Vespula rufa consobrina, predacious on larvae of Neodiprion swainei in Quebec, 73.

Vespula vulgaris, predacious on larvae of Neodiprion swainei in Quebec, 73.

vestitus, Anthonomus; Hylesinus.

Vetch (see Vicia).

Vetch, Hairy (see Vicia villosa).

viburniana, Aphelia (Tortrix).

Viburnum opulus, as primary food-plant of Aphis fabae in Bulgaria, 536.

Vicia, Sitona lineatus on, in U.S.A., 134; root nodules of, destroyed by S. lineatus 134.

Vicia faba (see Beans, Broad).

Vicia villosa, pests of, in Oregon, 211.

viciae, Megoura.

Victoria, aphids and virus disease of broad beans in, 34; Sminthurus viridis on pastures in, 34; Didymuria violescens on Eucalyptus in, 32, 33; natural enemies of pests in, 32, 34; Tyria jacobaeae introduced into, for control of Senecio jacobaeae, 452.

viennensis, Tetranychus.

Vietnam, Xyleborus spp. on coffee in, 618.

vigintioctomaculata, Epilachna. vigintioctopunctata, Epilachna.

vigintiquatuorpunetata Subcocci

vigintiquatuorpunctata, Subcoccinella. Vigna unguiculata (sinensis) (see Cowpeas). Villa hottentota, parasitising Apamea anceps in U.S.S.R., 363.

villiger, Ptinus.

viminalis, Trichiocampus.

Vine, Grape, pests of, in S. Africa, 83, 298, 650; pests of, in Austria, 85, 248; pests of, in Bulgaria, 58, 60, 110, 348, 538, 539; Brevipalpus chilensis on, in Chile, 495; pests of, in France, 8, 9, 83, 84, 255, 372, 434, 445, 494, 667, 668; pests of, in Germany, 71, 108, 109, 255; Tetranychus telarius on, in Holland, 682; pests of, in India, 595; Planococcus citri on, in Israel, 614; pests of, in Italy, 40, 48, 183, 310, 660, 661, 663; Eotetranychus smithi on, in Japan, 452; damaged by Vespula germanica in New Zealand, Argyrotaenia pulchellana on, in Switzerland, 100; Chloropsalta viridissima on, in Syria, 608; pests of, in U.S.S.R., 15; pests of, in U.S.A., 147, 265, 398; development of Cryptoblabes gnidiella on, 612; root-infesting form of Phylloxera vitifoliae on, 572; effects of nutrition of, on populations of Eotetranychus carpini vitis, 256; mildew of, 538; transmission of virus diseases of, 71, 303, 372; comparison of damage to, by virus and mites, 663; systemic activity of insecticides in, 434;

insecticide residues on, 228, 398; effects of fungicides on, 108.

vinitor, Nysius.

violescens, Didymuria.

Violet, Tetranychus multisetis feeding on, 67.

virens, Apion.

virescens, Heliothis; Temnochila.

virgata, Ferrisiana.

Virginia, pests of crucifers in, 335, 404; Hypera variabilis on lucerne in, 400; orchard pests in, 20, 335, 577; Neodiprion pratti pratti on pines in, 336; Popillia japonica on Ginkgo biloba in, 577; natural enemies of pests in, 164, 400; virus disease of Trichoplusia ni in,

Virginia, West, Hypera variabilis on lucerne in, 636; Rhyacionia buoliana on red pine in, 632.

virginicus, Reticulitermes.

viridana, Tortrix.

viridiceps, Syrphus.

viridifasciata, Chortophaga.

viridis, Ceramidia; Chermes; Coccus; Rhopalosiphum; Sminthurus; Tettigella (Cicadella).

viridissima, Chloropsalta; Tettigonia.

viridula, Nezara. virilis, Drosophila.

Virus Diseases (of insects), in Coleoptera, 690; in Diptera, 690; in Lepidoptera, 54, 55, 78, 144, 175, 196, 260, 283, 284, 300, 328, 335, 359, 363, 367, 368, 369, 389, 390, 400, 401, 404, 407, 423, 426, 427, 428, 448, 496, 531, 580, 581, 639, 660, 683, 686, 690, 693; epizootiology and synergism of, in *Pseudaletia uni*puncta, 283; in sawflies, 24, 25, 336, 341, 342, 580; in Panonychus citri, 581; laboratory experiments with, 284, 581, 690; field use of, 25, 328, 336, 342, 404, 423, 448, 531, 551, 580; ecology of use of, compared with chemical control, 262; insecticides in relation to, 359, 368, 538; other factors affecting susceptibility of insects to, 427; factors affecting incidence of, 335, 341, 342, 426; trans-ovarian transmission of, 341, 496, 580; question of latency of, 368, 531; inclusion bodies associated with, 581; booklet on, 550.

Virus Diseases (in plants *), relation of aphids to, 16, 17, 23, 34, 35, 71, 80, 81, 95, 97, 101, 106, 107, 130, 131, 132, 133, 134, 192, 197, 249, 250, 254, 255, 264, 298, 303, 304, 320, 321, 331, 357–359, 375, 401, 408, 441, 463, 467, 480, 482, 529, 544, 545, 546, 547, 549, 605, 611, 614, 636, 669, 680, 684, 688 610, 611, 614, 636, 669, 680, 684, 688, 689, 690, 691, 692; preference of Myzus

persicae for plants infected with, 584; classification of transmission of, by aphids, 691; injection of, into aphids, 133; relation of Cicadellids to, 23, 70, 304, 307, 336, 372, 408, 470, 524, 684, 686, 688, 689, 691; transmitted by Delphacids, 132, 257, 298, 349, 350, 508, 518, 689, 690; other Hemiptera in relation to, 130, 133, 465, 635, 636, 651, 652; transmitted by Coleoptera, 240, 304; transmitted by Mamestra brassicae, 443; preference of Pegomyia betae for beet free from, 253; transmitted by Tettigonia viridissima, 443; transmitted by Thrips tabaci, 345; review of chewing insects as vectors of, 443; transmission of, by mites, 70, 71, 302, 303, 523, 684; nematodes as vectors of, 71; transmitted by dodder, 71, 441; other means of transmission of, 684; effects of, on plants, 17, 408, 409, 663; varietal susceptibility or resistance of plants to, 17, 167, 241, 350; multiplication of, 691; factors affecting distribution and virulence of, 35, 271, 544; experimental inactivation of, 133, 443; other laboratory experiments with, 250, 254, 610, 611, 684; of field crops in Norway, 622; survey of data on, affecting graminaceous plants, 684; complexes of, 133.

visci, Carulaspis (see C. minima). Vitamins, in diets for insects, 5, 41.

viteana, Paralobesia.

vitifoliae, Phylloxera (Dactylosphaera). vitis, Eotetranychus carpini; Eriophyes;

Typhlodromus (see T. aberrans).

vitripennis, Nasonia. vittata, Phyllotreta.

vittiger, Calocoris norvegicus. vittula, Phyllotreta.

Voandzeia subterranea, experiments with Rivellia damaging root nodules of, 609. vulgaris, Melolontha (see M. melolontha); Vespula; Zenillia (Phryxe).

W

wairoensis, Coelostomidia.

w-album, Cratichneumon.

Wallis Islands, Aspidiotus destructor on coconut in, 81.

Walnut, pests of, in France, 151, 194; pests of, in U.S.A., 128, 147, 151, 276, 482, 491; review of Coleoptera infesting timber of, 194; (stored), fumigation of, against Paramyelois transitella, 276. Warehouses, Ephestia elutella in, 374.

WARF Antiresistant DDT, 641.

These references are also indexed under the individual plants infected.

warreni, Neodiprion.

Washington, Cydia pomonella on apple in, 466; aphids and virus diseases of beans in, 264; yellows virus of beet in, 17; Hylemya brassicae on crucifers in, 510; pests of mint in, 324; Sitona lineatus on peas and leguminous forage crops in, 134; Myzus persicae on potato in, 412; pests and virus diseases of cereals in, 281, 401; forest pests in, 209, 259, 385, 478; mites in stored grain in, 149; wild bees in, 299, 565; natural enemies of insects in, 565.

Water, influence of supply of, on attraction of bark-beetles to pines, 340.

Water, Hot, treatment of *Narcissus* bulbs with, against *Merodon equestris*, **288**; eggs of *Ostrinia nubilalis* freed from infection with *Perezia pyraustae* by treatment with, **343**.

waterhousei, Longitarsus.

Water-nut (see Trapa bispinosa).

watersi, Diparopsis.

Waterstonia, keys to females of species of, 490; new species of, 491.

Waterstonia fuliginosa, sp.n., parasitising Phenacoccus aceris in Japan. 491.

Wax, mode of action of dusts in removal of, from insects, 496; in stable formulations of dichlorvos, 562.

Wax Bean (see Beans, Wax).

wayi, Pseudotheraptus.

Webworm, Juniper (see Dichomeris mar-

ginella).

Weeds, freeing of rose-beds from, as measure against overwintering Tetranychid, 477; review of biological control of, in Canada, 692. (See also under individual plants.)

Wesson's Salts, in diets for insects, 214,

404.

Western Hemlock (see Tsuga hetero-phylla).

Wetting Agents, evaluation of, 140. Weymouth Pine (see *Pinus strobus*).

Wheat, pests of, in Australia, 493, 684; pests of, in Britain, 50, 51, 288, 431; pests of, in Bulgaria, 109, 348; pests of, in Canada, 70, 71, 72, 160, 523, 524, 527, 529; pests of, in China, 625; pests of, in Egypt, 684; Calligypona pellucida on, in Finland, 102, 103; pests of, in India, 119, 627, 684; Calocoris spp. on, in Italy, 310; Meromyza saltatrix on, in Japan, 96, 97; pests of, in Jordan, 684; Nematocerus sp. on, in Kenya, 180; leafhoppers on, in Pakistan, 684; pests of, in Persia, 428, 429, 484, 485, 684; Haplodiplosis equestris on, in Sweden, 101; pests of, in Turkey, 297, 484, 485; pests of, in U.S.S.R.,

13, 14, 243, 244, 343, 345, 362, 363, 446, 447, 595; pests of, in U.S.A., 65. 71, 214, 219, 470, 575; varietal susceptibility or resistance of, to insects, 22, 65, 72, 219, 420, 470, 526, 527, 575, 629; chemical differences between varieties of, in relation to feeding by Agrotis orthogonia, 527; arthropods and virus diseases of, 71, 132, 523, 524, 684; not infested by Camarota curvipennis, 249; survival of Chloriona orizicola on, 132; Chlorops pumilionis on, 113; study of, in relation to feeding by Oscinella frit, 595; Pentatomids in relation to, 429, 446, 653; strain of Aceria tulipae on, 156; use of, in rearing insects, 73, 575, 654; detection of growth inhibitors in, 22; seed treatment of, 50, 51, 627; toxicity of insecticides to, 50, 627; insecticides stimulating growth of, 627; effects on vetch fauna of sprays against pests of, 211.

Wheat (Stored), Coleoptera in, 2, 3, 118, 154, 155, 178, 212, 215, 270, 315, 344, 347, 499, 500, 554, 566; disturbance of, in relation to temperature requirements of *Ptinus* spp. in, 344; development of *Sitophilus* spp. on, 205, 206; mites in, 154, 155; other insects in, 154, methods for detection of infestation of, 571; fumigation of, 118, 347; sorption of fumigants by, 215, 554; other insecticide treatments of, 2, 3, 178, 212, 270, 347; effects of cleaning processes on insecticide residues on, 270; effects of fumigants on germination of, 579; structure and use of fossae for, in

Malta, 177, 178.

Wheat Bran, in diets for locust, 170.

Wheat Bulb Fly (see *Hylemya coarctata*). Wheat Flour, in baits for *Hylemya* spp., 388; in rearing medium for Tenebrionids, 89.

Wheat Germ, in diets for insects, 465, 474,

517, 657.

Wheatfeed, in diets for Coleoptera, 486, 656, 657.

Whitewash, BHC applied in, 374. wiedemanni, Metasyrphus.

wilkinsoni, Podacanthus.

Willow, Coleoptera on, in France, 83; pests of, in Poland, 40, 189; pests of, in Spain, 353; flea-beetles on, in Yugoslavia, 181; Lymantria dispar on, 514. (See also Salix spp.)

Willow, White (see Salix alba).

Wind, loss of larvae of *Trichiocampus* viminalis caused by, 392; effects of: on dispersal of insects, 33, 305, 318, 344, 430, 501, 596; on populations of Coccids and their natural enemies, 54;

on catches of insects in traps, 30; on effectiveness of insecticides, 16.

Wineberry (see Rubus phoenicolasius).

Winter Moth (see Operophtera brumata). Winthemia, parasitising larvae of Caligo eurilochus in Ecuador, 274.

Winthemia quadripustulata (datanae) (parasitising Lepidoptera), in Connecticut, 407; in U.S.S.R., 244, 245.

Wireworm, Southern Potato (see Cono-

derus falli).

Wireworms, in Canada, 73, 156, 388; in China, 625; in France, 371, 433; in Greece, 352; in U.S.S.R., 14; in U.S.A., 324, 420, 582; effects of diet on larvae of, 495; experiments on feeding response of, to rye-seed extracts, 528; factors affecting abundance of, 73, 371; method of rearing, 575; mites infesting cultures of, 575; effects of bacterial toxins on populations of, 582; traps for, 156; insecticides against, 324, 352, 371, 388, 421, 433; effects of insecticides against, on populations of Scutigerella immaculata, 433.

Wisconsin, pests of apple in, 415, 417; pests of cereals and grasses in, 4, 95, 160, 276, 396; virus diseases of cereals in, 95, 160; Hylemya spp. on crucifers in, 516, 568; Diprion similis on white pine in, 418; nematodes associated with Scolytus multistriatus in, 619; termites infesting buildings in, 416;

beneficial insects in, 415.

WL-1650 (see 1,3,4,5,6,7,8,8-Octachloro-3a,4,7,7a-tetrahydro-4,7-methanophthalan).

Wofatox (see Methyl-parathion).

woglumi, Aleurocanthus.

woodi, Acarapis.

Woodpeckers, destroying Formica polyctena in Germany, 104, 141; predacious Dendroctonus barberi in Mexico, 460.

Woollen Fabrics, treatments of, against Anthrenocerus australis, 587; attraction of insects to stains of, 635; use of, in rearing insects, 339.

Wormwood (see Artemisia vulgaris).

379-X, 572.

Xanthoencyrtus sacchari, sp.n., reared from Ripersia sacchari in India, 91. xanthoria, Encolpotis.

xanthostigma, Apanteles.

Xenoschesis fulvipes, parasitic on Acantholyda erythrocephala in Germany, 10.

Xenufens, egg-parasite of Caligo eurilochus in Ecuador, 267, 274; parasitised by mites, 268.

Xenufens ruskini, **267.**

Xestobium rufovillosum, in timber in Poland, 142.

X-rays, effects of, on fecundity and development of *Pexicopia malvella*, 113; use of, in detection of pests in plants and stored products, 259, 495,

Xyleborus, on coffee in Ivory Coast, 7: BHC sprays ineffective against, 7.

Xyleborus dispar (infesting oak), in Germany, 11; in Yugoslavia, 31; treatment of timber against, 31.

Xyleborus fornicatus, on tea in Ceylon, 449, 554; sprays against, 449, 554.

Xyleborus germanus, on deciduous trees in Germany, 11; bionomics of, damaging persimmon in Japan, 450; insecticides against, 451.

Xyleborus monographus, infesting oak logs in Yugoslavia, 31; treatment of

timber against, 31.

Xyleborus morigerus, on orchids imported into Britain and France from New Guinea, 625; on timber trees and other plants in Java, 625; on coffee in Vietnam, 618; survey of data on, 625.

Xyleborus morstatti (on coffee), in Ivory Coast, 651; in Vietnam, 618; Colletotrichum coffeanum associated 651; spray mixtures against, 651.

Xylene, as solvent for insecticides, 226. Xylocoris galactinus, in stored grain in Canada, 154.

xyloni, Solenopsis.

xylophagorum, Roptrocerus (Pachyceras). Xylosandrus (see Xyleborus).

xylosteanus, Archips.

Xyloterus (see Trypodendron).

379-Y, 572.

Yams, new Coccid on, in New Guinea,

yanonensis, Unaspis (Prontaspis).

Yeast, relation of, to tolerance of nicotine by Lasioderma serricorne, 650; attraction of insects to, 339, 340; in diets and rearing media for insects, 89, 170, 184, 214, 339, 626, 657.

Yeast Hydrolysate, in rearing media for

mites, **641**.

Yeast Protein, in baits: for Anastrepha ludens, 279, 280; for Gryllotalpa, 442. yohoensis, Ips.

ypsilon, Agrotis.

Yugoslavia, cereal pests in, 80, 181, 249; Apion spp. on red clover in, 32; pests of fruit trees in, 32, 54, 55, 138, 239, 240, 241; Prays oleellus on olive in, 609; Haltica oleracea on strawberry in, 242; pests of basket willows in, 138, 181; forest pests in, 31, 54, 55, 181, 241, 535, 536; pests of vegetables in, 181, 240, 366, 681; flea-beetles on medicinal plants in, 181; pests of ornamental plants in, 32, 55, 181; survey of data on flea-beetles in, 181; Hyphantria cunea in, 368; Lymantria dispar in, 54, 55, 367-371; virus diseases of Lepidoptera in, 54, 55, 367, 368, 369; beneficial insects in, 31, 54, 55, 138, 240, 367, 370, 371, 682; virus diseases of plants in, 80, 240.

Z

Zabrotes subfasciatus (in stored beans), in Angola, 99; in Colombia, 560; insecticides against, 99, 560, 561.

Zabrus tenebrioides, characteristics and world distribution of, 507.

Zagrammosoma americanum (parasitising Evagora spp.), in California, 222; in Canada, 586.

Zagrammosoma nigrolineatum, egg-parasite of Caliroa limacina in Quebec, 26.

Zagrammosoma variegatum afrum, parasitising Leucoptera in Tanganyika, 652. Zantedeschia aethiopica, Calacarus citri-

folii on, in S. Africa, 83.

Zanzibar, pests of coconut in, 28, 29; ants associated with *Pseudotheraptus wayi* in, 28, 29; new parasite of *Saissetia* sp. in, 491.

Zea mays (see Maize).

Zea mays indentata, susceptibility of, to Oscinella frit, 595.

zea, Heliothis.

Zeadiatraea grandiosella on maize in Arkansas, 21, 165; insecticides against, 21; method of winter rearing, 65. zealandica, Coelostomidia.

zeamais, Sitophilus.

Zectran (see 4-Dimethylamino-3,5-xylyl Methylcarbamate).

Zeiraphera (see Enarmonia).

Zelus, predacious on cotton pests in Arizona, 639.

Zenillia caesar, parasitising larvae of Ostrinia nubilalis in Quebec, 27.

Zenillia mitis, parasitising Ostrinia nubilalis in U.S.S.R., 245.

Zenillia vulgaris, Pieris rapae attacked by, in New York, 401,

zetterstedtii, Anabrolepis.

Zeuzera coffeae, on cacao in Dutch New Guinea, 76.

zimini, Masicera.

zimmermanni, Dioryctria.

Zinc Sulphate, effects of sprays containing, on beneficial arthropods, 36.

zinckenella, Etiella.

Zineb (in sprays and spray mixtures), against Cydia pomonella, 20; against mites, 108, 264, 409, 410, 556, 560; question of resistance to, in strains of Phyllocoptruta oleivora, 409; effects of, on beneficial arthropods, 159, 672; evaluation of spray programmes of, 376; effect of, on Aramite, 265.

Zinnia, Aphis gossypii transmitting virus

disease of, in Israel, 615.

Zinophos (see O,O-Diethyl O-2-Pyrazinyl Phosphorothioate).

Ziziphus jujuba, Perissopneumon tectonae on, in India, 174; Larvacarus transitans on, in Pakistan, 176, 626.

Ziziphus nummularia, Perissopneumon tectonae on, in India, 174.

ziziphus, Parlatoria.


zonellus, Chilo.


Zonosemata electa, on Capsicum in Ontario, 529.

Zootermopsis angusticollis, resistance of timbers to infestation by, 649.

Zygina subrufa, seed treatments against, on rice in China, 625.

Zytron (see O-Methyl O-2,4-Dichlorophenyl Isopropylphosphoramidothioate).





NOTICES

The annual subscription, payable in advance, to the Review of Applied Entomology, Series A (Agricultural) is 80s., post free; Series B (Medical and Veterinary) 40s., post free. Prices of back volumes on application; complete sets are available.

Orders and Subscriptions should be sent to the Director, Commonwealth Institute of Entomology, 56, Queen's Gate, London, S.W.7, or through a bookseller.

Secretaries of Societies and Editors of Journals wishing to exchange their publications for those of the Institute are requested to communicate with the Director. Authors of papers on economic entomology, whether published in entomological journals or not, are invited to send reprints to the Director for notice in the Review.

The Executive Council of the Commonwealth Agricultural Bureaux is a signatory to the Fair Copying Declaration, details of which can be obtained from the Royal Society, Burlington House, London, W.1.

The Commonwealth Agricultural Bureaux Organisation does not accept responsibility for any trade advertisements in this Journal.